

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C12N 15/00, C07K 14/00</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/37630</b> <b>(43) International Publication Date:</b> 29 June 2000 (29.06.00)
<b>(21) International Application Number:</b> PCT/US99/31005 <b>(22) International Filing Date:</b> 22 December 1999 (22.12.99) <b>(30) Priority Data:</b> 09/220,876 23 December 1998 (23.12.98) US <b>(71) Applicant:</b> GENETICS INSTITUTE, INC. [US/US]; 87 CambridgePark Drive, Cambridge, MA 02140 (US). <b>(72) Inventors:</b> JACOBS, Kenneth; 151 Beaumont Avenue, Newton, MA 02160 (US). MCCOY, John, M.; 56 Howard Street, Reading, MA 01867 (US). LAVALLIE, Edward, R.; 113 Ann Lee Road, Harvard, MA 01451 (US). COLLINS-RACIE, Lisa, A.; 124 School Street, Acton, MA 01720 (US). EVANS, Cheryl; 18801 Bent Willow Circle, Germantown, MD 20874 (US). MERBERG, David; 2 Orchard Drive, Acton, MA 01720 (US). TREACY, Maurice; 12 Foxrock Court, Dublin 18 (IE). BOWMAN, Michael, R.; 50 Aldrich Road, Canton, MA 02021 (US). <b>(74) Agent:</b> MANDRAGOURAS, Amy, E.; Lahive & Cockfield, LLP, 28 State Street, Boston, MA 02109 (US).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> SECRETED PROTEINS  <b>(57) Abstract</b>  Novel proteins are disclosed.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## SECRETED PROTEINS

5 This application is a continuation-in-part of the following applications:

- (1) Ser. No. 08/634,325, filed April 18, 1996;
- (2) Ser. No. 08/783,520, filed January 13, 1997, which is a  
continuation-in-part of application Ser. No. 08/634,325, filed April 18,  
10 1996;
- (3) Ser. No. 08/885,610, filed June 30, 1997, which is a continuation-in-part  
of application Ser. No. 08/634,325, filed April 18, 1996;
- (4) Ser. No. 08/943,861, filed October 3, 1997, which is a continuation of  
application Ser. No. 60/080,227 (converted to a provisional application  
15 from non-provisional application 08/725,885), filed October 4, 1996;
- (5) Ser. No. 08/943,862, filed October 3, 1997, which is a continuation of  
application Ser. No. 60/093,043 (converted to a provisional application  
from non-provisional application 08/726,257), filed October 4, 1996;
- (6) Ser. No. 08/960,024, filed October 29, 1997, which is a  
20 continuation-in-part of application Ser. No. 60/077,176 (converted to a  
provisional application from non-provisional application 08/742,973),  
filed November 1, 1996; and
- (7) Ser. No. 09/137,226, filed August 20, 1998, which is a  
continuation-in-part of application Ser. No. 60/092,114 (converted to a  
25 provisional application from non-provisional application 08/916,041),  
filed August 21, 1997;

all of which are incorporated by reference herein.

30

### FIELD OF THE INVENTION

5           The present invention provides novel proteins , along with therapeutic, diagnostic and research utilities for these proteins.

### BACKGROUND OF THE INVENTION

10           Technology aimed at the discovery of protein factors (including e.g., cytokines, such as lymphokines, interferons, CSFs and interleukins) has matured rapidly over the past decade. The now routine hybridization cloning and expression cloning techniques clone novel polynucleotides "directly" in the sense that they rely on information directly related to the discovered protein (i.e., partial DNA/amino acid  
15   sequence of the protein in the case of hybridization cloning; activity of the protein in the case of expression cloning). More recent "indirect" cloning techniques such as signal sequence cloning, which isolates DNA sequences based on the presence of a now well-recognized secretory leader sequence motif, as well as various PCR-based or low stringency hybridization cloning techniques, have advanced the state of the art by making  
20   available large numbers of DNA/amino acid sequences for proteins that are known to have biological activity by virtue of their secreted nature in the case of leader sequence cloning, or by virtue of the cell or tissue source in the case of PCR-based techniques. It is to these proteins that the present invention is directed.



SUMMARY OF THE INVENTION

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- 5 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:1;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:1 from nucleotide 19 to nucleotide 561;
- (c) a polynucleotide comprising the nucleotide sequence of the  
10 full-length protein coding sequence of clone AK296\_1i deposited under accession number ATCC 98026;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AK296\_1i deposited under accession number ATCC 98026;
- 15 (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AK296\_1i deposited under accession number ATCC 98026;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AK296\_1i deposited under accession number ATCC 98026;
- 20 (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:2;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:2 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:2;
- 25 (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above; and
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ  
30 ID NO:1 from nucleotide 19 to nucleotide 561; the nucleotide sequence of the full-length protein coding sequence of clone AK296\_1i deposited under accession number ATCC 98026; or the nucleotide sequence of a mature protein coding sequence of clone AK296\_1i deposited under accession number ATCC 98026. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert

of clone AK296\_1i deposited under accession number ATCC 98026. In yet other preferred embodiments, such polynucleotide encodes a protein comprising the amino acid sequence of SEQ ID NO:2 from amino acid 3 to amino acid 181. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a  
5 fragment of the amino acid sequence of SEQ ID NO:2 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:2, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:2 having biological activity, the fragment comprising the amino acid sequence from amino acid 85 to amino  
10 acid 94 of SEQ ID NO:2.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:2;
- 15 (b) the amino acid sequence of SEQ ID NO:2 from amino acid 3 to amino acid 181;
- (c) fragments of the amino acid sequence of SEQ ID NO:2, each fragment comprising eight consecutive amino acids of SEQ ID NO:2; and
- (d) the amino acid sequence encoded by the cDNA insert of  
20 clone AK296\_1i deposited under accession number ATCC 98026;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:2 or the amino acid sequence of SEQ ID NO:2 from amino acid 3 to amino acid 181. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid  
25 sequence of SEQ ID NO:2 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:2, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:2, the fragment comprising the amino acid sequence from amino acid 85 to amino acid 94 of SEQ ID NO:2.

30 In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:3;

- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:3 from nucleotide 123 to nucleotide 1457;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AK533\_1i deposited under accession number ATCC 98026;
- 5 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AK533\_1i deposited under accession number ATCC 98026;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AK533\_1i deposited under accession number ATCC 98026;
- 10 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AK533\_1i deposited under accession number ATCC 98026;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:4;
- 15 (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:4 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:4;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above; and
- 20 (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:3 from nucleotide 123 to nucleotide 1457; the nucleotide sequence of the full-length protein coding sequence of clone AK533\_1i deposited under accession number ATCC 98026; or the nucleotide sequence of a mature protein coding sequence of clone AK533\_1i deposited under accession number ATCC 98026. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AK533\_1i deposited under accession number ATCC 98026.

30 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:4;
- (b) fragments of the amino acid sequence of SEQ ID NO:4, each

fragment comprising eight consecutive amino acids of SEQ ID NO:4; and

(c) the amino acid sequence encoded by the cDNA insert of clone AK533\_1i deposited under accession number ATCC 98026;

the protein being substantially free from other mammalian proteins. Preferably such  
5 protein comprises the amino acid sequence of SEQ ID NO:4. In further preferred  
embodiments, the present invention provides a protein comprising a fragment of the  
amino acid sequence of SEQ ID NO:4 having biological activity, the fragment preferably  
comprising eight (more preferably twenty, most preferably thirty) consecutive amino  
acids of SEQ ID NO:4, or a protein comprising a fragment of the amino acid sequence of  
10 SEQ ID NO:4, the fragment comprising the amino acid sequence from amino acid 217 to  
amino acid 226 of SEQ ID NO:4.

In one embodiment, the present invention provides a composition  
comprising an isolated protein encoded by a polynucleotide selected from the group  
consisting of:

15 (a) a polynucleotide comprising the nucleotide sequence of SEQ  
ID NO:5;

(b) a polynucleotide comprising the nucleotide sequence of  
SEQ ID NO:5 from nucleotide 258 to nucleotide 392;

(c) a polynucleotide comprising the nucleotide sequence of SEQ  
20 ID NO:5 from nucleotide 330 to nucleotide 392;

(d) a polynucleotide comprising the nucleotide sequence of the  
full-length protein coding sequence of clone AK583\_1i deposited under accession  
number ATCC 98026;

(e) a polynucleotide encoding the full-length protein encoded  
25 by the cDNA insert of clone AK583\_1i deposited under accession number ATCC  
98026;

(f) a polynucleotide comprising the nucleotide sequence of a  
mature protein coding sequence of clone AK583\_1i deposited under accession  
number ATCC 98026;

30 (g) a polynucleotide encoding a mature protein encoded by the  
cDNA insert of clone AK583\_1i deposited under accession number ATCC 98026;

(h) a polynucleotide encoding a protein comprising the amino  
acid sequence of SEQ ID NO:6;

(i) a polynucleotide encoding a protein comprising a fragment

of the amino acid sequence of SEQ ID NO:6 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:6;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above; and

5 (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:5 from nucleotide 258 to nucleotide 392; the nucleotide sequence of SEQ ID NO:5 from nucleotide 330 to nucleotide 392; the nucleotide sequence of the full-length protein  
10 coding sequence of clone AK583\_1i deposited under accession number ATCC 98026; or the nucleotide sequence of a mature protein coding sequence of clone AK583\_1i deposited under accession number ATCC 98026. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AK583\_1i deposited under accession number ATCC 98026.

15 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:6;
- (b) fragments of the amino acid sequence of SEQ ID NO:6, each  
20 fragment comprising eight consecutive amino acids of SEQ ID NO:6; and
- (c) the amino acid sequence encoded by the cDNA insert of clone AK583\_1i deposited under accession number ATCC 98026;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:6. In further preferred  
25 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:6 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:6, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:6, the fragment comprising the amino acid sequence from amino acid 17 to  
30 amino acid 26 of SEQ ID NO:6.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ

ID NO:7;

(b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:7 from nucleotide 6 to nucleotide 1424;

(c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:7 from nucleotide 78 to nucleotide 1424;

(d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AM282\_1i deposited under accession number ATCC 98026;

(e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AM282\_1i deposited under accession number ATCC 98026;

(f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AM282\_1i deposited under accession number ATCC 98026;

(g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AM282\_1i deposited under accession number ATCC 98026;

(h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:8;

(i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:8;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above; and

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:7 from nucleotide 6 to nucleotide 1424; the nucleotide sequence of SEQ ID NO:7 from nucleotide 78 to nucleotide 1424; the nucleotide sequence of the full-length protein coding sequence of clone AM282\_1i deposited under accession number ATCC 98026; or the nucleotide sequence of a mature protein coding sequence of clone AM282\_1i deposited under accession number ATCC 98026. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AM282\_1i deposited under accession number ATCC 98026. In yet other preferred embodiments, such polynucleotide encodes a protein comprising the amino acid sequence

of SEQ ID NO:8 from amino acid 1 to amino acid 91. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:8; or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8 having biological activity, the fragment comprising the amino acid sequence from amino acid 231 to amino acid 240 of SEQ ID NO:8.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:8;
- (b) the amino acid sequence of SEQ ID NO:8 from amino acid 1 to amino acid 91;
- (c) fragments of the amino acid sequence of SEQ ID NO:8, each fragment comprising eight consecutive amino acids of SEQ ID NO:8; and
- (d) the amino acid sequence encoded by the cDNA insert of clone AM282\_1i deposited under accession number ATCC 98026;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:8 or the amino acid sequence of SEQ ID NO:8 from amino acid 1 to amino acid 91. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:8, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8, the fragment comprising the amino acid sequence from amino acid 231 to amino acid 240 of SEQ ID NO:8.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:9;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:9 from nucleotide 87 to nucleotide 458;

- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:9 from nucleotide 378 to nucleotide 458;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AM340\_1i deposited under accession number ATCC 98026;
- 5 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AM340\_1i deposited under accession number ATCC 98026;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AM340\_1i deposited under accession number ATCC 98026;
- 10 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AM340\_1i deposited under accession number ATCC 98026;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:10;
- 15 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:10 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:10;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above; and
- 20 (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:9 from nucleotide 87 to nucleotide 458; the nucleotide sequence of SEQ ID NO:9 from nucleotide 378 to nucleotide 458; the nucleotide sequence of the full-length protein coding sequence of clone AM340\_1i deposited under accession number ATCC 98026; or the nucleotide sequence of a mature protein coding sequence of clone AM340\_1i deposited under accession number ATCC 98026. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AM340\_1i deposited under accession number ATCC 98026.

25 30

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:10;



- (b) fragments of the amino acid sequence of SEQ ID NO:10, each fragment comprising eight consecutive amino acids of SEQ ID NO:10; and
- (c) the amino acid sequence encoded by the cDNA insert of clone AM340\_1i deposited under accession number ATCC 98026;
- 5 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:10. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:10 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino
- 10 acids of SEQ ID NO:10, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:10, the fragment comprising the amino acid sequence from amino acid 57 to amino acid 66 of SEQ ID NO:10.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group

15 consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:11;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:11 from nucleotide 17 to nucleotide 685;
- 20 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:11 from nucleotide 86 to nucleotide 685;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AM610\_1i deposited under accession number ATCC 98026;
- 25 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AM610\_1i deposited under accession number ATCC 98026;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AM610\_1i deposited under accession
- 30 number ATCC 98026;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AM610\_1i deposited under accession number ATCC 98026;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:12;

(i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:12 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:12;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above; and

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:11 from nucleotide 17 to nucleotide 685; the nucleotide sequence of SEQ ID NO:11 from nucleotide 86 to nucleotide 685; the nucleotide sequence of the full-length protein coding sequence of clone AM610\_1i deposited under accession number ATCC 98026; or the nucleotide sequence of a mature protein coding sequence of clone AM610\_1i deposited under accession number ATCC 98026. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AM610\_1i deposited under accession number ATCC 98026.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:12;
- (b) fragments of the amino acid sequence of SEQ ID NO:12, each fragment comprising eight consecutive amino acids of SEQ ID NO:12; and
- (c) the amino acid sequence encoded by the cDNA insert of clone AM610\_1i deposited under accession number ATCC 98026;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:12. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:12 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:12, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:12, the fragment comprising the amino acid sequence from amino acid 106 to amino acid 115 of SEQ ID NO:12.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:13;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:13 from nucleotide 70 to nucleotide 504;
- 5 (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AP162\_1i deposited under accession number ATCC 98026;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AP162\_1i deposited under accession number ATCC 98026;
- 10 (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AP162\_1i deposited under accession number ATCC 98026;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AP162\_1i deposited under accession number ATCC 98026;
- 15 (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:14;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:14 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:14;
- 20 (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above; and
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above.
- 25 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:13 from nucleotide 70 to nucleotide 504; the nucleotide sequence of the full-length protein coding sequence of clone AP162\_1i deposited under accession number ATCC 98026; or the nucleotide sequence of a mature protein coding sequence of clone AP162\_1i deposited under accession number ATCC 98026. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AP162\_1i deposited under accession number ATCC 98026. In yet other preferred embodiments, such polynucleotide encodes a protein comprising the amino acid sequence of SEQ ID NO:14 from amino acid 42 to amino acid 61. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a
- 30

fragment of the amino acid sequence of SEQ ID NO:14 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:14, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:14 having biological activity, the fragment comprising the amino acid sequence from amino acid 67 to amino acid 76 of SEQ ID NO:14.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 10 (a) the amino acid sequence of SEQ ID NO:14;
- (b) the amino acid sequence of SEQ ID NO:14 from amino acid 42 to amino acid 61;
- (c) fragments of the amino acid sequence of SEQ ID NO:14, each fragment comprising eight consecutive amino acids of SEQ ID NO:14; and
- 15 (d) the amino acid sequence encoded by the cDNA insert of clone AP162\_1i deposited under accession number ATCC 98026;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:14 or the amino acid sequence of SEQ ID NO:14 from amino acid 42 to amino acid 61. In further preferred embodiments,

20 the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:14 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:14, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:14, the fragment comprising the amino acid sequence from amino acid 67 to amino acid 76

25 of SEQ ID NO:14.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- 30 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:16;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:16 from nucleotide 77 to nucleotide 694;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AR260\_1i deposited under accession

number ATCC 98026;

(d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AR260\_1i deposited under accession number ATCC 98026;

5 (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AR260\_1i deposited under accession number ATCC 98026;

(f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AR260\_1i deposited under accession number ATCC 98026;

10 (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:17;

(h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:17 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:17;

15 (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above; and

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:16 from nucleotide 77 to nucleotide 694; the nucleotide sequence of the full-length protein coding sequence of clone AR260\_1i deposited under accession number ATCC 98026; or the nucleotide sequence of a mature protein coding sequence of clone AR260\_1i deposited under accession number ATCC 98026. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AR260\_1i deposited under accession number ATCC 98026.

25 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 30 (a) the amino acid sequence of SEQ ID NO:17;
- (b) fragments of the amino acid sequence of SEQ ID NO:17, each fragment comprising eight consecutive amino acids of SEQ ID NO:17; and
- (c) the amino acid sequence encoded by the cDNA insert of clone AR260\_1i deposited under accession number ATCC 98026;

the protein being substantially free from other mammalian proteins. Preferably such

protein comprises the amino acid sequence of SEQ ID NO:17. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:17 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:17, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:17, the fragment comprising the amino acid sequence from amino acid 98 to amino acid 107 of SEQ ID NO:17.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:18;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:18 from nucleotide 23 to nucleotide 676;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AS32\_1i deposited under accession number ATCC 98026;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AS32\_1i deposited under accession number ATCC 98026;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AS32\_1i deposited under accession number ATCC 98026;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AS32\_1i deposited under accession number ATCC 98026;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:19;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:19 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:19;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above; and
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:18 from nucleotide 23 to nucleotide 676; the nucleotide sequence of the full-length protein coding sequence of clone AS32\_1i deposited under accession number ATCC 98026; or the nucleotide sequence of a mature protein coding sequence of clone AS32\_1i deposited under accession number ATCC 98026. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AS32\_1i deposited under accession number ATCC 98026. In yet other preferred embodiments, such polynucleotide encodes a protein comprising the amino acid sequence of SEQ ID NO:19 from amino acid 78 to amino acid 97. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:19 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:19, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:19 having biological activity, the fragment comprising the amino acid sequence from amino acid 102 to amino acid 111 of SEQ ID NO:19.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:19;
- (b) the amino acid sequence of SEQ ID NO:19 from amino acid 78 to amino acid 97;
- (c) fragments of the amino acid sequence of SEQ ID NO:19, each fragment comprising eight consecutive amino acids of SEQ ID NO:19; and
- (d) the amino acid sequence encoded by the cDNA insert of clone AS32\_1i deposited under accession number ATCC 98026;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:19 or the amino acid sequence of SEQ ID NO:19 from amino acid 78 to amino acid 97. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:19 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:19, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:19, the fragment comprising the amino acid sequence from amino acid 102 to amino acid 111

of SEQ ID NO:19.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- 5 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:21;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:21 from nucleotide 65 to nucleotide 490;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:21 from nucleotide 137 to nucleotide 490;
- 10 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AS34\_1i deposited under accession number ATCC 98026;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AS34\_1i deposited under accession number ATCC 98026;
- 15 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AS34\_1i deposited under accession number ATCC 98026;
- 20 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AS34\_1i deposited under accession number ATCC 98026;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:22;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:22 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:22;
- 25 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above; and
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above.
- 30

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:21 from nucleotide 65 to nucleotide 490; the nucleotide sequence of SEQ ID NO:21 from nucleotide 137 to nucleotide 490; the nucleotide sequence of the full-length protein coding sequence of clone AS34\_1i deposited under accession number ATCC 98026; or the



nucleotide sequence of a mature protein coding sequence of clone AS34\_1i deposited under accession number ATCC 98026. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AS34\_1i deposited under accession number ATCC 98026.

5 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:22;
- (b) fragments of the amino acid sequence of SEQ ID NO:22,
- 10 each fragment comprising eight consecutive amino acids of SEQ ID NO:22; and
- (c) the amino acid sequence encoded by the cDNA insert of clone AS34\_1i deposited under accession number ATCC 98026;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:22. In further preferred  
15 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:22 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:22, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:22, the fragment comprising the amino acid sequence from amino acid 66 to  
20 amino acid 75 of SEQ ID NO:22.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ  
25 ID NO:23;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:23 from nucleotide 225 to nucleotide 677;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:23 from nucleotide 387 to nucleotide 677;
- 30 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AT205\_1i deposited under accession number ATCC 98026;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AT205\_1i deposited under accession number ATCC

98026;

(f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AT205\_1i deposited under accession number ATCC 98026;

5 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AT205\_1i deposited under accession number ATCC 98026;

(h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:24;

10 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:24 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:24;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above; and

15 (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:23 from nucleotide 225 to nucleotide 677; the nucleotide sequence of SEQ ID NO:23 from nucleotide 387 to nucleotide 677; the nucleotide sequence of the full-length protein coding sequence of clone AT205\_1i deposited under accession number ATCC 98026; or  
20 the nucleotide sequence of a mature protein coding sequence of clone AT205\_1i deposited under accession number ATCC 98026. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AT205\_1i deposited under accession number ATCC 98026. In yet other preferred embodiments, such polynucleotide encodes a protein comprising the amino acid sequence  
25 of SEQ ID NO:24 from amino acid 6 to amino acid 25. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:24 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:24, or a polynucleotide encoding a protein  
30 comprising a fragment of the amino acid sequence of SEQ ID NO:24 having biological activity, the fragment comprising the amino acid sequence from amino acid 70 to amino acid 79 of SEQ ID NO:24.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected

from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:24;
- (b) the amino acid sequence of SEQ ID NO:24 from amino acid 6 to amino acid 25;
- 5 (c) fragments of the amino acid sequence of SEQ ID NO:24, each fragment comprising eight consecutive amino acids of SEQ ID NO:24; and
- (d) the amino acid sequence encoded by the cDNA insert of clone AT205\_1i deposited under accession number ATCC 98026;

the protein being substantially free from other mammalian proteins. Preferably such  
10 protein comprises the amino acid sequence of SEQ ID NO:24 or the amino acid sequence of SEQ ID NO:24 from amino acid 6 to amino acid 25. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:24 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID  
15 NO:24, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:24, the fragment comprising the amino acid sequence from amino acid 70 to amino acid 79 of SEQ ID NO:24.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group  
20 consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:25;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:25 from nucleotide 38 to nucleotide 832;
- 25 (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AT211\_1i deposited under accession number ATCC 98026;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AT211\_1i deposited under accession number ATCC  
30 98026;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AT211\_1i deposited under accession number ATCC 98026;
- (f) a polynucleotide encoding a mature protein encoded by the

cDNA insert of clone AT211\_1i deposited under accession number ATCC 98026;

(g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:26;

5 (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:26 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:26;

(i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above; and

10 (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:25 from nucleotide 38 to nucleotide 832; the nucleotide sequence of the full-length protein coding sequence of clone AT211\_1i deposited under accession number ATCC 98026; or the nucleotide sequence of a mature protein coding sequence of clone AT211\_1i  
15 deposited under accession number ATCC 98026. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AT211\_1i deposited under accession number ATCC 98026.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected  
20 from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:26;

(b) fragments of the amino acid sequence of SEQ ID NO:26, each fragment comprising eight consecutive amino acids of SEQ ID NO:26; and

25 (c) the amino acid sequence encoded by the cDNA insert of clone AT211\_1i deposited under accession number ATCC 98026;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:26. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:26 having biological activity, the fragment preferably  
30 comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:26, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:26, the fragment comprising the amino acid sequence from amino acid 127 to amino acid 136 of SEQ ID NO:26.

In one embodiment, the present invention provides a composition

comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:27;
- 5 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:27 from nucleotide 194 to nucleotide 423;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AT319\_1i deposited under accession number ATCC 98026;
- 10 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AT319\_1i deposited under accession number ATCC 98026;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AT319\_1i deposited under accession number ATCC 98026;
- 15 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AT319\_1i deposited under accession number ATCC 98026;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:28;
- 20 (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:28 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:28;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above; and
- 25 (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:27 from nucleotide 194 to nucleotide 423; the nucleotide sequence of the full-length protein coding sequence of clone AT319\_1i deposited under accession number ATCC 98026; or the nucleotide sequence of a mature protein coding sequence of clone AT319\_1i deposited under accession number ATCC 98026. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AT319\_1i deposited under accession number ATCC 98026. In yet other preferred embodiments, such polynucleotide encodes a protein comprising the amino acid sequence

of SEQ ID NO:28 from amino acid 2 to amino acid 21. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:28 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:28, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:28 having biological activity, the fragment comprising the amino acid sequence from amino acid 30 to amino acid 39 of SEQ ID NO:28.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:28;
- (b) the amino acid sequence of SEQ ID NO:28 from amino acid 2 to amino acid 21;
- (c) fragments of the amino acid sequence of SEQ ID NO:28, each fragment comprising eight consecutive amino acids of SEQ ID NO:28; and
- (d) the amino acid sequence encoded by the cDNA insert of clone AT319\_1i deposited under accession number ATCC 98026;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:28 or the amino acid sequence of SEQ ID NO:28 from amino acid 2 to amino acid 21. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:28 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:28, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:28, the fragment comprising the amino acid sequence from amino acid 30 to amino acid 39 of SEQ ID NO:28.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:30;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:30 from nucleotide 61 to nucleotide 514;

- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:30 from nucleotide 112 to nucleotide 514;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AW191\_1i deposited under accession number ATCC 98026;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AW191\_1i deposited under accession number ATCC 98026;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AW191\_1i deposited under accession number ATCC 98026;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AW191\_1i deposited under accession number ATCC 98026;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:31;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:31 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:31;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above; and
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:30 from nucleotide 61 to nucleotide 514; the nucleotide sequence of SEQ ID NO:30 from nucleotide 112 to nucleotide 514; the nucleotide sequence of the full-length protein coding sequence of clone AW191\_1i deposited under accession number ATCC 98026; or the nucleotide sequence of a mature protein coding sequence of clone AW191\_1i deposited under accession number ATCC 98026. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AW191\_1i deposited under accession number ATCC 98026. In yet other preferred embodiments, such polynucleotide encodes a protein comprising the amino acid sequence of SEQ ID NO:31 from amino acid 24 to amino acid 43. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:31 having biological activity, the

fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:31, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:31 having biological activity, the fragment comprising the amino acid sequence from amino acid 70 to amino acid 79 of SEQ ID NO:31.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:31;
  - (b) the amino acid sequence of SEQ ID NO:31 from amino acid 24 to amino acid 43;
  - (c) fragments of the amino acid sequence of SEQ ID NO:31, each fragment comprising eight consecutive amino acids of SEQ ID NO:31; and
  - (d) the amino acid sequence encoded by the cDNA insert of clone AW191\_1i deposited under accession number ATCC 98026;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:31 or the amino acid sequence of SEQ ID NO:31 from amino acid 24 to amino acid 43. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:31 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:31, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:31, the fragment comprising the amino acid sequence from amino acid 70 to amino acid 79 of SEQ ID NO:31.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:33;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:33 from nucleotide 14 to nucleotide 391;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone BB9\_1i deposited under accession number ATCC 98026;



- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone BB9\_1i deposited under accession number ATCC 98026;
- 5 (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone BB9\_1i deposited under accession number ATCC 98026;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone BB9\_1i deposited under accession number ATCC 98026;
- 10 (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:34;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:34 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:34;
- 15 (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above; and
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:33 from nucleotide 14 to nucleotide 391; the nucleotide sequence of the full-length  
20 protein coding sequence of clone BB9\_1i deposited under accession number ATCC 98026; or the nucleotide sequence of a mature protein coding sequence of clone BB9\_1i deposited under accession number ATCC 98026. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone BB9\_1i deposited under accession number ATCC 98026. In yet other preferred  
25 embodiments, such polynucleotide encodes a protein comprising the amino acid sequence of SEQ ID NO:34 from amino acid 75 to amino acid 94. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:34 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty)  
30 consecutive amino acids of SEQ ID NO:34, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:34 having biological activity, the fragment comprising the amino acid sequence from amino acid 55 to amino acid 64 of SEQ ID NO:34.

In other embodiments, the present invention provides a composition

comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:34;
  - (b) the amino acid sequence of SEQ ID NO:34 from amino acid  
5 75 to amino acid 94;
  - (c) fragments of the amino acid sequence of SEQ ID NO:34,  
each fragment comprising eight consecutive amino acids of SEQ ID NO:34; and
  - (d) the amino acid sequence encoded by the cDNA insert of  
clone BB9\_1i deposited under accession number ATCC 98026;
- 10 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:34 or the amino acid sequence of SEQ ID NO:34 from amino acid 75 to amino acid 94. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:34 having biological activity, the fragment preferably comprising  
15 eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:34, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:34, the fragment comprising the amino acid sequence from amino acid 55 to amino acid 64 of SEQ ID NO:34.

In one embodiment, the present invention provides a composition  
20 comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ  
ID NO:36;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ  
25 ID NO:36 from nucleotide 58 to nucleotide 655;
- (c) a polynucleotide comprising the nucleotide sequence of the  
full-length protein coding sequence of clone H617\_1i deposited under accession  
number ATCC 98026;
- (d) a polynucleotide encoding the full-length protein encoded  
30 by the cDNA insert of clone H617\_1i deposited under accession number ATCC  
98026;
- (e) a polynucleotide comprising the nucleotide sequence of a  
mature protein coding sequence of clone H617\_1i deposited under accession  
number ATCC 98026;

- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone H617\_1i deposited under accession number ATCC 98026;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:37;
- 5 (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:37 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:37;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above; and
- 10 (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:36 from nucleotide 58 to nucleotide 655; the nucleotide sequence of the full-length protein coding sequence of clone H617\_1i deposited under accession number ATCC 98026; or the nucleotide sequence of a mature protein coding sequence of clone H617\_1i deposited under accession number ATCC 98026. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone H617\_1i deposited under accession number ATCC 98026. In yet other preferred embodiments, such polynucleotide encodes a protein comprising the amino acid sequence of SEQ ID NO:37 from amino acid 65 to amino acid 84. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:37 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:37, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:37 having biological activity, the fragment comprising the amino acid sequence from amino acid 95 to amino acid 104 of SEQ ID NO:37.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:37;
- (b) the amino acid sequence of SEQ ID NO:37 from amino acid 65 to amino acid 84;
- (c) fragments of the amino acid sequence of SEQ ID NO:37,

each fragment comprising eight consecutive amino acids of SEQ ID NO:37; and

(d) the amino acid sequence encoded by the cDNA insert of clone H617\_1i deposited under accession number ATCC 98026;

the protein being substantially free from other mammalian proteins. Preferably such  
5 protein comprises the amino acid sequence of SEQ ID NO:37 or the amino acid sequence  
of SEQ ID NO:37 from amino acid 65 to amino acid 84. In further preferred embodiments,  
the present invention provides a protein comprising a fragment of the amino acid  
sequence of SEQ ID NO:37 having biological activity, the fragment preferably comprising  
eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID  
10 NO:37, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:37,  
the fragment comprising the amino acid sequence from amino acid 95 to amino acid 104  
of SEQ ID NO:37.

In one embodiment, the present invention provides a composition  
comprising an isolated protein encoded by a polynucleotide selected from the group  
15 consisting of:

(a) a polynucleotide comprising the nucleotide sequence of SEQ  
ID NO:39;

(b) a polynucleotide comprising the nucleotide sequence of SEQ  
ID NO:39 from nucleotide 71 to nucleotide 377;

20 (c) a polynucleotide comprising the nucleotide sequence of the  
full-length protein coding sequence of clone K39\_1i deposited under accession  
number ATCC 98026;

(d) a polynucleotide encoding the full-length protein encoded  
by the cDNA insert of clone K39\_1i deposited under accession number ATCC  
25 98026;

(e) a polynucleotide comprising the nucleotide sequence of a  
mature protein coding sequence of clone K39\_1i deposited under accession  
number ATCC 98026;

(f) a polynucleotide encoding a mature protein encoded by the  
30 cDNA insert of clone K39\_1i deposited under accession number ATCC 98026;

(g) a polynucleotide encoding a protein comprising the amino  
acid sequence of SEQ ID NO:40;

(h) a polynucleotide encoding a protein comprising a fragment  
of the amino acid sequence of SEQ ID NO:40 having biological activity, the

fragment comprising eight consecutive amino acids of SEQ ID NO:40;

(i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above; and

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above.

5 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:39 from nucleotide 71 to nucleotide 377; the nucleotide sequence of the full-length protein coding sequence of clone K39\_1i deposited under accession number ATCC 98026; or the nucleotide sequence of a mature protein coding sequence of clone K39\_1i deposited  
10 under accession number ATCC 98026. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone K39\_1i deposited under accession number ATCC 98026. In yet other preferred embodiments, such polynucleotide encodes a protein comprising the amino acid sequence of SEQ ID NO:40 from amino acid 62 to amino acid 81.

15 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:40;

(b) the amino acid sequence of SEQ ID NO:40 from amino acid  
20 62 to amino acid 81;

(c) fragments of the amino acid sequence of SEQ ID NO:40, each fragment comprising eight consecutive amino acids of SEQ ID NO:40; and

(d) the amino acid sequence encoded by the cDNA insert of clone K39\_1i deposited under accession number ATCC 98026;

25 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:40. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:40 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino  
30 acids of SEQ ID NO:40, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:40, the fragment comprising the amino acid sequence from amino acid 46 to amino acid 55 of SEQ ID NO:40.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group

consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:42;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:42 from nucleotide 1 to nucleotide 332;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone K640\_1i deposited under accession number ATCC 98026;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone K640\_1i deposited under accession number ATCC 98026;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone K640\_1i deposited under accession number ATCC 98026;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone K640\_1i deposited under accession number ATCC 98026;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:43;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:43 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:43;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above; and
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:42 from nucleotide 1 to nucleotide 332; the nucleotide sequence of the full-length protein coding sequence of clone K640\_1i deposited under accession number ATCC 98026; or the nucleotide sequence of a mature protein coding sequence of clone K640\_1i deposited under accession number ATCC 98026. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone K640\_1i deposited under accession number ATCC 98026. In yet other preferred embodiments, such polynucleotide encodes a protein comprising the amino acid sequence of SEQ ID NO:43 from amino acid 11 to amino acid 30. In further preferred embodiments,

the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:43 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:43, or a polynucleotide encoding a protein  
5 comprising a fragment of the amino acid sequence of SEQ ID NO:43 having biological activity, the fragment comprising the amino acid sequence from amino acid 50 to amino acid 59 of SEQ ID NO:43.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected  
10 from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:43;
- (b) the amino acid sequence of SEQ ID NO:43 from amino acid  
11 to amino acid 30;
- (c) fragments of the amino acid sequence of SEQ ID NO:43,  
15 each fragment comprising eight consecutive amino acids of SEQ ID NO:43; and
- (d) the amino acid sequence encoded by the cDNA insert of  
clone K640\_1i deposited under accession number ATCC 98026;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:43 or the amino acid sequence  
20 of SEQ ID NO:43 from amino acid 11 to amino acid 30. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:43 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:43, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:43,  
25 the fragment comprising the amino acid sequence from amino acid 50 to amino acid 59 of SEQ ID NO:43.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- 30 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:45;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:45 from nucleotide 12 to nucleotide 539;
- (c) a polynucleotide comprising the nucleotide sequence of the

full-length protein coding sequence of clone AE402\_1i deposited under accession number ATCC 98190;

(d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AE402\_1i deposited under accession number ATCC 98190;

(e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AE402\_1i deposited under accession number ATCC 98190;

(f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AE402\_1i deposited under accession number ATCC 98190;

(g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:46;

(h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:46 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:46;

(i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above; and

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:45 from nucleotide 12 to nucleotide 539; the nucleotide sequence of the full-length protein coding sequence of clone AE402\_1i deposited under accession number ATCC 98190; or the nucleotide sequence of a mature protein coding sequence of clone AE402\_1i deposited under accession number ATCC 98190. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AE402\_1i deposited under accession number ATCC 98190.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:46;
- (b) fragments of the amino acid sequence of SEQ ID NO:46, each fragment comprising eight consecutive amino acids of SEQ ID NO:46; and
- (c) the amino acid sequence encoded by the cDNA insert of clone AE402\_1i deposited under accession number ATCC 98190;



the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:46. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:46 having biological activity, the fragment preferably  
5 comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:46, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:46, the fragment comprising the amino acid sequence from amino acid 83 to amino acid 92 of SEQ ID NO:46.

In one embodiment, the present invention provides a composition  
10 comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:47;
- (b) a polynucleotide comprising the nucleotide sequence of  
15 SEQ ID NO:47 from nucleotide 61 to nucleotide 513;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:47 from nucleotide 322 to nucleotide 513;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AE610\_1i deposited under accession  
20 number ATCC 98190;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AE610\_1i deposited under accession number ATCC 98190;
- (f) a polynucleotide comprising the nucleotide sequence of a  
25 mature protein coding sequence of clone AE610\_1i deposited under accession number ATCC 98190;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AE610\_1i deposited under accession number ATCC 98190;
- (h) a polynucleotide encoding a protein comprising the amino  
30 acid sequence of SEQ ID NO:48;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:48 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:48;
- (j) a polynucleotide which is an allelic variant of a

polynucleotide of (a)-(g) above; and

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:47 from nucleotide 61 to nucleotide 513; the nucleotide sequence of SEQ ID NO:47 from nucleotide 322 to nucleotide 513; the nucleotide sequence of the full-length protein coding sequence of clone AE610\_1i deposited under accession number ATCC 98190; or the nucleotide sequence of a mature protein coding sequence of clone AE610\_1i deposited under accession number ATCC 98190. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AE610\_1i deposited under accession number ATCC 98190.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 15 (a) the amino acid sequence of SEQ ID NO:48;
  - (b) fragments of the amino acid sequence of SEQ ID NO:48, each fragment comprising eight consecutive amino acids of SEQ ID NO:48; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone AE610\_1i deposited under accession number ATCC 98190;
- 20 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:48. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:48 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino
- 25 acids of SEQ ID NO:48, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:48, the fragment comprising the amino acid sequence from amino acid 70 to amino acid 79 of SEQ ID NO:48.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group

30 consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:50;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:50 from nucleotide 20 to nucleotide 523;

(c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AH106\_1i deposited under accession number ATCC 98190;

5 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AH106\_1i deposited under accession number ATCC 98190;

(e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AH106\_1i deposited under accession number ATCC 98190;

10 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AH106\_1i deposited under accession number ATCC 98190;

(g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:51;

15 (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:51 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:51;

(i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above; and

20 (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:50 from nucleotide 20 to nucleotide 523; the nucleotide sequence of the full-length protein coding sequence of clone AH106\_1i deposited under accession number ATCC 98190; or the nucleotide sequence of a mature protein coding sequence of clone AH106\_1i deposited under accession number ATCC 98190. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AH106\_1i deposited under accession number ATCC 98190.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 30 (a) the amino acid sequence of SEQ ID NO:51;
- (b) fragments of the amino acid sequence of SEQ ID NO:51, each fragment comprising eight consecutive amino acids of SEQ ID NO:51; and
- (c) the amino acid sequence encoded by the cDNA insert of

clone AH106\_1i deposited under accession number ATCC 98190;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:51. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:51 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:51, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:51, the fragment comprising the amino acid sequence from amino acid 79 to amino acid 88 of SEQ ID NO:51.

10 In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:52;
- 15 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:52 from nucleotide 130 to nucleotide 309;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AH196\_1i deposited under accession number ATCC 98190;
- 20 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AH196\_1i deposited under accession number ATCC 98190;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AH196\_1i deposited under accession number ATCC 98190;
- 25 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AH196\_1i deposited under accession number ATCC 98190;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:53;
- 30 (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:53 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:53;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above; and

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:52 from nucleotide 130 to nucleotide 309; the nucleotide sequence of the full-length protein coding sequence of clone AH196\_1i deposited under accession number ATCC 98190; or the nucleotide sequence of a mature protein coding sequence of clone AH196\_1i deposited under accession number ATCC 98190. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AH196\_1i deposited under accession number ATCC 98190.

10 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:53;
- (b) fragments of the amino acid sequence of SEQ ID NO:53,
- 15 each fragment comprising eight consecutive amino acids of SEQ ID NO:53; and
- (c) the amino acid sequence encoded by the cDNA insert of clone AH196\_1i deposited under accession number ATCC 98190;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:53. In further preferred  
20 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:53 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:53, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:53, the fragment comprising the amino acid sequence from amino acid 25  
25 amino acid 34 of SEQ ID NO:53.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ  
30 ID NO:55;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:55 from nucleotide 69 to nucleotide 467;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AI6\_1i deposited under accession

number ATCC 98190;

(d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AI6\_1i deposited under accession number ATCC 98190;

5 (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AI6\_1i deposited under accession number ATCC 98190;

(f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AI6\_1i deposited under accession number ATCC 98190;

10 (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:56;

(h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:56 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:56;

15 (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above; and

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:55 from nucleotide 69 to nucleotide 467; the nucleotide sequence of the full-length protein coding sequence of clone AI6\_1i deposited under accession number ATCC 98190; or the nucleotide sequence of a mature protein coding sequence of clone AI6\_1i deposited under accession number ATCC 98190. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AI6\_1i deposited under accession number ATCC 98190. In yet other preferred  
20 embodiments, such polynucleotide encodes a protein comprising the amino acid sequence of SEQ ID NO:56 from amino acid 69 to amino acid 133. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:56 having biological  
25 activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:56, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:56 having biological activity, the fragment comprising the amino acid sequence from amino acid 61 to amino acid 70 of SEQ ID NO:56.  
30

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:56;
  - 5 (b) the amino acid sequence of SEQ ID NO:56 from amino acid 69 to amino acid 133;
  - (c) fragments of the amino acid sequence of SEQ ID NO:56, each fragment comprising eight consecutive amino acids of SEQ ID NO:56; and
  - (d) the amino acid sequence encoded by the cDNA insert of
  - 10 clone AI6\_1i deposited under accession number ATCC 98190;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:56 or the amino acid sequence of SEQ ID NO:56 from amino acid 69 to amino acid 133. In further preferred embodiments, the present invention provides a protein comprising a fragment of the
- 15 amino acid sequence of SEQ ID NO:56 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:56, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:56, the fragment comprising the amino acid sequence from amino acid 61 to amino acid 70 of SEQ ID NO:56.

20 In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:58;
- 25 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:58 from nucleotide 55 to nucleotide 363;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AJ13\_1i deposited under accession number ATCC 98190;
- 30 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AJ13\_1i deposited under accession number ATCC 98190;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AJ13\_1i deposited under accession

number ATCC 98190;

(f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AJ13\_1i deposited under accession number ATCC 98190;

(g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:59;

(h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:59 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:59;

(i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above; and

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:58 from nucleotide 55 to nucleotide 363; the nucleotide sequence of the full-length protein coding sequence of clone AJ13\_1i deposited under accession number ATCC 98190; or the nucleotide sequence of a mature protein coding sequence of clone AJ13\_1i deposited under accession number ATCC 98190. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AJ13\_1i deposited under accession number ATCC 98190.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:59;

(b) fragments of the amino acid sequence of SEQ ID NO:59, each fragment comprising eight consecutive amino acids of SEQ ID NO:59; and

(c) the amino acid sequence encoded by the cDNA insert of clone AJ13\_1i deposited under accession number ATCC 98190;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:59. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:59 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:59, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:59, the fragment comprising the amino acid sequence from amino acid 46 to



amino acid 55 of SEQ ID NO:59.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- 5 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:60;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:60 from nucleotide 33 to nucleotide 422;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ  
10 ID NO:60 from nucleotide 114 to nucleotide 422;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AJ27\_1i deposited under accession number ATCC 98190;
- (e) a polynucleotide encoding the full-length protein encoded  
15 by the cDNA insert of clone AJ27\_1i deposited under accession number ATCC 98190;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AJ27\_1i deposited under accession number ATCC 98190;
- 20 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AJ27\_1i deposited under accession number ATCC 98190;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:61;
- (i) a polynucleotide encoding a protein comprising a fragment  
25 of the amino acid sequence of SEQ ID NO:61 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:61;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above; and
- (k) a polynucleotide which encodes a species homologue of the  
30 protein of (h) or (i) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:60 from nucleotide 33 to nucleotide 422; the nucleotide sequence of SEQ ID NO:60 from nucleotide 114 to nucleotide 422; the nucleotide sequence of the full-length protein coding sequence of clone AJ27\_1i deposited under accession number ATCC 98190; or the

nucleotide sequence of a mature protein coding sequence of clone AJ27\_1i deposited under accession number ATCC 98190. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AJ27\_1i deposited under accession number ATCC 98190.

5 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:61;
- (b) fragments of the amino acid sequence of SEQ ID NO:61,
- 10 each fragment comprising eight consecutive amino acids of SEQ ID NO:61; and
- (c) the amino acid sequence encoded by the cDNA insert of clone AJ27\_1i deposited under accession number ATCC 98190;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:61. In further preferred

15 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:61 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:61, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:61, the fragment comprising the amino acid sequence from amino acid 60 to

20 amino acid 69 of SEQ ID NO:61.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ
- 25 ID NO:63;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:63 from nucleotide 47 to nucleotide 517;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:63 from nucleotide 116 to nucleotide 517;
- 30 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AJ142\_1i deposited under accession number ATCC 98190;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AJ142\_1i deposited under accession number ATCC

98190;

(f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AJ142\_1i deposited under accession number ATCC 98190;

5 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AJ142\_1i deposited under accession number ATCC 98190;

(h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:64;

10 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:64 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:64;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above; and

15 (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:63 from nucleotide 47 to nucleotide 517; the nucleotide sequence of SEQ ID NO:63 from nucleotide 116 to nucleotide 517; the nucleotide sequence of the full-length protein coding sequence of clone AJ142\_1i deposited under accession number ATCC 98190; or the  
20 nucleotide sequence of a mature protein coding sequence of clone AJ142\_1i deposited under accession number ATCC 98190. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AJ142\_1i deposited under accession number ATCC 98190.

25 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:64;

(b) fragments of the amino acid sequence of SEQ ID NO:64, each fragment comprising eight consecutive amino acids of SEQ ID NO:64; and

30 (c) the amino acid sequence encoded by the cDNA insert of clone AJ142\_1i deposited under accession number ATCC 98190;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:64. In further preferred embodiments, the present invention provides a protein comprising a fragment of the

amino acid sequence of SEQ ID NO:64 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:64, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:64, the fragment comprising the amino acid sequence from amino acid 73 to amino acid 82 of SEQ ID NO:64.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:65;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:65 from nucleotide 312 to nucleotide 417;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AK604\_1i deposited under accession number ATCC 98190;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AK604\_1i deposited under accession number ATCC 98190;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AK604\_1i deposited under accession number ATCC 98190;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AK604\_1i deposited under accession number ATCC 98190;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:66;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:66 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:66;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above; and
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:65 from nucleotide 312 to nucleotide 417; the nucleotide sequence of the full-length

protein coding sequence of clone AK604\_1i deposited under accession number ATCC 98190; or the nucleotide sequence of a mature protein coding sequence of clone AK604\_1i deposited under accession number ATCC 98190. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert  
5 of clone AK604\_1i deposited under accession number ATCC 98190.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:66;
  - 10 (b) fragments of the amino acid sequence of SEQ ID NO:66, each fragment comprising eight consecutive amino acids of SEQ ID NO:66; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone AK604\_1i deposited under accession number ATCC 98190;
- the protein being substantially free from other mammalian proteins. Preferably such  
15 protein comprises the amino acid sequence of SEQ ID NO:66. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:66 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:66, or a protein comprising a fragment of the amino acid sequence of  
20 SEQ ID NO:66, the fragment comprising the amino acid sequence from amino acid 12 to amino acid 21 of SEQ ID NO:66.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- 25 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:68;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:68 from nucleotide 57 to nucleotide 353;
- (c) a polynucleotide comprising the nucleotide sequence of the  
30 full-length protein coding sequence of clone AK620\_1i deposited under accession number ATCC 98190;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AK620\_1i deposited under accession number ATCC 98190;

- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AK620\_1i deposited under accession number ATCC 98190;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AK620\_1i deposited under accession number ATCC 98190;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:69;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:69 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:69;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above; and
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:68 from nucleotide 57 to nucleotide 353; the nucleotide sequence of the full-length protein coding sequence of clone AK620\_1i deposited under accession number ATCC 98190; or the nucleotide sequence of a mature protein coding sequence of clone AK620\_1i deposited under accession number ATCC 98190. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AK620\_1i deposited under accession number ATCC 98190.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:69;
- (b) fragments of the amino acid sequence of SEQ ID NO:69, each fragment comprising eight consecutive amino acids of SEQ ID NO:69; and
- (c) the amino acid sequence encoded by the cDNA insert of clone AK620\_1i deposited under accession number ATCC 98190;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:69. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:69 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino

acids of SEQ ID NO:69, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:69, the fragment comprising the amino acid sequence from amino acid 44 to amino acid 53 of SEQ ID NO:69.

In one embodiment, the present invention provides a composition  
5 comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:70;
- (b) a polynucleotide comprising the nucleotide sequence of  
10 SEQ ID NO:70 from nucleotide 464 to nucleotide 751;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:70 from nucleotide 542 to nucleotide 751;
- (d) a polynucleotide comprising the nucleotide sequence of the  
15 full-length protein coding sequence of clone AK650\_1i deposited under accession number ATCC 98190;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AK650\_1i deposited under accession number ATCC 98190;
- (f) a polynucleotide comprising the nucleotide sequence of a  
20 mature protein coding sequence of clone AK650\_1i deposited under accession number ATCC 98190;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AK650\_1i deposited under accession number ATCC 98190;
- (h) a polynucleotide encoding a protein comprising the amino  
25 acid sequence of SEQ ID NO:71;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:71 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:71;
- (j) a polynucleotide which is an allelic variant of a  
30 polynucleotide of (a)-(g) above; and
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:70 from nucleotide 464 to nucleotide 751; the nucleotide sequence of SEQ ID NO:70

from nucleotide 542 to nucleotide 751; the nucleotide sequence of the full-length protein coding sequence of clone AK650\_1i deposited under accession number ATCC 98190; or the nucleotide sequence of a mature protein coding sequence of clone AK650\_1i deposited under accession number ATCC 98190. In other preferred embodiments, the  
5 polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AK650\_1i deposited under accession number ATCC 98190.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 10 (a) the amino acid sequence of SEQ ID NO:71;
  - (b) fragments of the amino acid sequence of SEQ ID NO:71, each fragment comprising eight consecutive amino acids of SEQ ID NO:71; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone AK650\_1i deposited under accession number ATCC 98190;
- 15 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:71. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:71 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino  
20 acids of SEQ ID NO:71, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:71, the fragment comprising the amino acid sequence from amino acid 43 to amino acid 52 of SEQ ID NO:71.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group  
25 consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:72;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:72 from nucleotide 116 to nucleotide 310;
- 30 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:72 from nucleotide 173 to nucleotide 310;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AM226\_1i deposited under accession number ATCC 98190;



- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AM226\_1i deposited under accession number ATCC 98190;
- 5 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AM226\_1i deposited under accession number ATCC 98190;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AM226\_1i deposited under accession number ATCC 98190;
- 10 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:73;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:73 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:73;
- 15 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above; and
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:72 from nucleotide 116 to nucleotide 310; the nucleotide sequence of SEQ ID NO:72 from nucleotide 173 to nucleotide 310; the nucleotide sequence of the full-length protein coding sequence of clone AM226\_1i deposited under accession number ATCC 98190; or the nucleotide sequence of a mature protein coding sequence of clone AM226\_1i deposited under accession number ATCC 98190. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AM226\_1i deposited under accession number ATCC 98190.

25

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 30 (a) the amino acid sequence of SEQ ID NO:73;
- (b) fragments of the amino acid sequence of SEQ ID NO:73, each fragment comprising eight consecutive amino acids of SEQ ID NO:73; and
- (c) the amino acid sequence encoded by the cDNA insert of clone AM226\_1i deposited under accession number ATCC 98190;
- the protein being substantially free from other mammalian proteins. Preferably such

protein comprises the amino acid sequence of SEQ ID NO:73. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:73 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:73, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:73, the fragment comprising the amino acid sequence from amino acid 27 to amino acid 36 of SEQ ID NO:73.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:75;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:75 from nucleotide 220 to nucleotide 453;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:75 from nucleotide 352 to nucleotide 453;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AR417\_1i deposited under accession number ATCC 98190;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AR417\_1i deposited under accession number ATCC 98190;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AR417\_1i deposited under accession number ATCC 98190;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AR417\_1i deposited under accession number ATCC 98190;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:76;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:76 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:76;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above; and

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:75 from nucleotide 220 to nucleotide 453; the nucleotide sequence of SEQ ID NO:75 from nucleotide 352 to nucleotide 453; the nucleotide sequence of the full-length protein coding sequence of clone AR417\_1i deposited under accession number ATCC 98190; or the nucleotide sequence of a mature protein coding sequence of clone AR417\_1i deposited under accession number ATCC 98190. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AR417\_1i deposited under accession number ATCC 98190.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:76;
- (b) fragments of the amino acid sequence of SEQ ID NO:76, each fragment comprising eight consecutive amino acids of SEQ ID NO:76; and
- (c) the amino acid sequence encoded by the cDNA insert of clone AR417\_1i deposited under accession number ATCC 98190;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:76. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:76 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:76, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:76, the fragment comprising the amino acid sequence from amino acid 34 to amino acid 43 of SEQ ID NO:76.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:77;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:77 from nucleotide 496 to nucleotide 583;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ

ID NO:77 from nucleotide 565 to nucleotide 583;

(d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AU43\_1i deposited under accession number ATCC 98190;

5 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AU43\_1i deposited under accession number ATCC 98190;

10 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AU43\_1i deposited under accession number ATCC 98190;

(g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AU43\_1i deposited under accession number ATCC 98190;

(h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:78;

15 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:78 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:78;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above; and

20 (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:77 from nucleotide 496 to nucleotide 583; the nucleotide sequence of SEQ ID NO:77 from nucleotide 565 to nucleotide 583; the nucleotide sequence of the full-length protein coding sequence of clone AU43\_1i deposited under accession number ATCC 98190; or the  
25 nucleotide sequence of a mature protein coding sequence of clone AU43\_1i deposited under accession number ATCC 98190. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AU43\_1i deposited under accession number ATCC 98190.

30 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:78;

(b) fragments of the amino acid sequence of SEQ ID NO:78,

each fragment comprising eight consecutive amino acids of SEQ ID NO:78; and

(c) the amino acid sequence encoded by the cDNA insert of clone AU43\_1i deposited under accession number ATCC 98190;

the protein being substantially free from other mammalian proteins. Preferably such

5 protein comprises the amino acid sequence of SEQ ID NO:78. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:78 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:78, or a protein comprising a fragment of the amino acid sequence of  
10 SEQ ID NO:78, the fragment comprising the amino acid sequence from amino acid 9 to amino acid 18 of SEQ ID NO:78.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

15 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:80;

(b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:80 from nucleotide 55 to nucleotide 405;

20 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:80 from nucleotide 148 to nucleotide 405;

(d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AW60\_1i deposited under accession number ATCC 98190;

25 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AW60\_1i deposited under accession number ATCC 98190;

(f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AW60\_1i deposited under accession number ATCC 98190;

30 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AW60\_1i deposited under accession number ATCC 98190;

(h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:81;

(i) a polynucleotide encoding a protein comprising a fragment

of the amino acid sequence of SEQ ID NO:81 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:81;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above; and

5 (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:80 from nucleotide 55 to nucleotide 405; the nucleotide sequence of SEQ ID NO:80 from nucleotide 148 to nucleotide 405; the nucleotide sequence of the full-length protein  
10 coding sequence of clone AW60\_1i deposited under accession number ATCC 98190; or the nucleotide sequence of a mature protein coding sequence of clone AW60\_1i deposited under accession number ATCC 98190. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AW60\_1i deposited under accession number ATCC 98190.

15 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:81;
- (b) fragments of the amino acid sequence of SEQ ID NO:81,  
20 each fragment comprising eight consecutive amino acids of SEQ ID NO:81; and
- (c) the amino acid sequence encoded by the cDNA insert of clone AW60\_1i deposited under accession number ATCC 98190;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:81. In further preferred  
25 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:81 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:81, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:81, the fragment comprising the amino acid sequence from amino acid 53 to  
30 amino acid 62 of SEQ ID NO:81.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ

ID NO:83;

(b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:83 from nucleotide 256 to nucleotide 1338;

5 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:83 from nucleotide 1120 to nucleotide 1338;

(d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone BA176\_1i deposited under accession number ATCC 98190;

10 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone BA176\_1i deposited under accession number ATCC 98190;

(f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone BA176\_1i deposited under accession number ATCC 98190;

15 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone BA176\_1i deposited under accession number ATCC 98190;

(h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:84;

20 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:84 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:84;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above; and

25 (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:83 from nucleotide 256 to nucleotide 1338; the nucleotide sequence of SEQ ID NO:83 from nucleotide 1120 to nucleotide 1338; the nucleotide sequence of the full-length protein coding sequence of clone BA176\_1i deposited under accession number ATCC 98190; or the nucleotide sequence of a mature protein coding sequence of clone BA176\_1i deposited under accession number ATCC 98190. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone BA176\_1i deposited under accession number ATCC 98190.

In other embodiments, the present invention provides a composition

comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:84;
- (b) fragments of the amino acid sequence of SEQ ID NO:84,  
5 each fragment comprising eight consecutive amino acids of SEQ ID NO:84; and
- (c) the amino acid sequence encoded by the cDNA insert of  
clone BA176\_1i deposited under accession number ATCC 98190;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:84. In further preferred  
10 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:84 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:84, or a protein comprising a fragment of the amino acid sequence of  
15 amino acid 184 of SEQ ID NO:84.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ  
20 ID NO:85;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ  
ID NO:85 from nucleotide 199 to nucleotide 396;
- (c) a polynucleotide comprising the nucleotide sequence of the  
full-length protein coding sequence of clone BD140\_1i deposited under accession  
25 number ATCC 98190;
- (d) a polynucleotide encoding the full-length protein encoded  
by the cDNA insert of clone BD140\_1i deposited under accession number ATCC  
98190;
- (e) a polynucleotide comprising the nucleotide sequence of a  
30 mature protein coding sequence of clone BD140\_1i deposited under accession  
number ATCC 98190;
- (f) a polynucleotide encoding a mature protein encoded by the  
cDNA insert of clone BD140\_1i deposited under accession number ATCC 98190;
- (g) a polynucleotide encoding a protein comprising the amino



acid sequence of SEQ ID NO:86;

(h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:86 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:86;

5 (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above; and

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ  
10 ID NO:85 from nucleotide 199 to nucleotide 396; the nucleotide sequence of the full-length protein coding sequence of clone BD140\_1i deposited under accession number ATCC 98190; or the nucleotide sequence of a mature protein coding sequence of clone BD140\_1i deposited under accession number ATCC 98190. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert  
15 of clone BD140\_1i deposited under accession number ATCC 98190.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:86;
- 20 (b) fragments of the amino acid sequence of SEQ ID NO:86, each fragment comprising eight consecutive amino acids of SEQ ID NO:86; and
- (c) the amino acid sequence encoded by the cDNA insert of clone BD140\_1i deposited under accession number ATCC 98190;

the protein being substantially free from other mammalian proteins. Preferably such  
25 protein comprises the amino acid sequence of SEQ ID NO:86. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:86 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:86, or a protein comprising a fragment of the amino acid sequence of  
30 SEQ ID NO:86, the fragment comprising the amino acid sequence from amino acid 28 to amino acid 37 of SEQ ID NO:86.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:87;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:87 from nucleotide 303 to nucleotide 617;
- 5 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:87 from nucleotide 345 to nucleotide 617;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone BD407\_1i deposited under accession number ATCC 98190;
- 10 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone BD407\_1i deposited under accession number ATCC 98190;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone BD407\_1i deposited under accession number ATCC 98190;
- 15 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone BD407\_1i deposited under accession number ATCC 98190;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:88;
- 20 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:88 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:88;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above; and
- 25 (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:87 from nucleotide 303 to nucleotide 617; the nucleotide sequence of SEQ ID NO:87 from nucleotide 345 to nucleotide 617; the nucleotide sequence of the full-length protein coding sequence of clone BD407\_1i deposited under accession number ATCC 98190; or the nucleotide sequence of a mature protein coding sequence of clone BD407\_1i deposited under accession number ATCC 98190. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone BD407\_1i deposited under accession number ATCC 98190. In yet other preferred

30

embodiments, such polynucleotide encodes a protein comprising the amino acid sequence of SEQ ID NO:88 from amino acid 1 to amino acid 32. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:88 having biological activity, the  
5 fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:88, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:88 having biological activity, the fragment comprising the amino acid sequence from amino acid 47 to amino acid 56 of SEQ ID NO:88.

10 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:88;
  - (b) the amino acid sequence of SEQ ID NO:88 from amino acid  
15 1 to amino acid 32;
  - (c) fragments of the amino acid sequence of SEQ ID NO:88, each fragment comprising eight consecutive amino acids of SEQ ID NO:88; and
  - (d) the amino acid sequence encoded by the cDNA insert of clone BD407\_1i deposited under accession number ATCC 98190;
- 20 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:88 or the amino acid sequence of SEQ ID NO:88 from amino acid 1 to amino acid 32. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:88 having biological activity, the fragment preferably comprising  
25 eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:88, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:88, the fragment comprising the amino acid sequence from amino acid 47 to amino acid 56 of SEQ ID NO:88.

In one embodiment, the present invention provides a composition  
30 comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ  
ID NO:89;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ

ID NO:89 from nucleotide 152 to nucleotide 535;

(c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone BF290\_1i deposited under accession number ATCC 98190;

5 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone BF290\_1i deposited under accession number ATCC 98190;

(e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone BF290\_1i deposited under accession  
10 number ATCC 98190;

(f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone BF290\_1i deposited under accession number ATCC 98190;

(g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:90;

15 (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:90 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:90;

(i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above; and

20 (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:89 from nucleotide 152 to nucleotide 535; the nucleotide sequence of the full-length protein coding sequence of clone BF290\_1i deposited under accession number ATCC  
25 98190; or the nucleotide sequence of a mature protein coding sequence of clone BF290\_1i deposited under accession number ATCC 98190. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone BF290\_1i deposited under accession number ATCC 98190.

In other embodiments, the present invention provides a composition  
30 comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:90;

(b) fragments of the amino acid sequence of SEQ ID NO:90, each fragment comprising eight consecutive amino acids of SEQ ID NO:90; and

(c) the amino acid sequence encoded by the cDNA insert of clone BF290\_1i deposited under accession number ATCC 98190; the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:90. In further preferred  
5 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:90 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:90, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:90, the fragment comprising the amino acid sequence from amino acid 59 to  
10 amino acid 68 of SEQ ID NO:90.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- 15 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:91;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:91 from nucleotide 160 to nucleotide 474;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:91 from nucleotide 331 to nucleotide 474;
- 20 (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone BG236\_1i deposited under accession number ATCC 98191;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone BG236\_1i deposited under accession number ATCC  
25 98191;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone BG236\_1i deposited under accession number ATCC 98191;
- (g) a polynucleotide encoding a mature protein encoded by the  
30 cDNA insert of clone BG236\_1i deposited under accession number ATCC 98191;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:92;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:92 having biological activity, the

fragment comprising eight consecutive amino acids of SEQ ID NO:92;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above; and

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:91 from nucleotide 160 to nucleotide 474; the nucleotide sequence of SEQ ID NO:91 from nucleotide 331 to nucleotide 474; the nucleotide sequence of the full-length protein coding sequence of clone BG236\_1i deposited under accession number ATCC 98191; or the nucleotide sequence of a mature protein coding sequence of clone BG236\_1i deposited under accession number ATCC 98191. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone BG236\_1i deposited under accession number ATCC 98191.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:92;
- (b) fragments of the amino acid sequence of SEQ ID NO:92, each fragment comprising eight consecutive amino acids of SEQ ID NO:92; and
- (c) the amino acid sequence encoded by the cDNA insert of clone BG236\_1i deposited under accession number ATCC 98191;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:92. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:92 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:92, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:92, the fragment comprising the amino acid sequence from amino acid 47 to amino acid 56 of SEQ ID NO:92.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:93;

- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:93 from nucleotide 139 to nucleotide 419;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone BG237\_1i deposited under accession number ATCC 98191;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone BG237\_1i deposited under accession number ATCC 98191;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone BG237\_1i deposited under accession number ATCC 98191;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone BG237\_1i deposited under accession number ATCC 98191;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:94;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:94 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:94;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above; and
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:93 from nucleotide 139 to nucleotide 419; the nucleotide sequence of the full-length protein coding sequence of clone BG237\_1i deposited under accession number ATCC 98191; or the nucleotide sequence of a mature protein coding sequence of clone BG237\_1i deposited under accession number ATCC 98191. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone BG237\_1i deposited under accession number ATCC 98191. In yet other preferred embodiments, such polynucleotide encodes a protein comprising the amino acid sequence of SEQ ID NO:94 from amino acid 9 to amino acid 93. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:94 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty)

consecutive amino acids of SEQ ID NO:94, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:94 having biological activity, the fragment comprising the amino acid sequence from amino acid 41 to amino acid 50 of SEQ ID NO:94.

5 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:94;
  - (b) the amino acid sequence of SEQ ID NO:94 from amino acid  
10 9 to amino acid 93;
  - (c) fragments of the amino acid sequence of SEQ ID NO:94, each fragment comprising eight consecutive amino acids of SEQ ID NO:94; and
  - (d) the amino acid sequence encoded by the cDNA insert of clone BG237\_1i deposited under accession number ATCC 98191;
- 15 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:94 or the amino acid sequence of SEQ ID NO:94 from amino acid 9 to amino acid 93. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:94 having biological activity, the fragment preferably comprising  
20 eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:94, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:94, the fragment comprising the amino acid sequence from amino acid 41 to amino acid 50 of SEQ ID NO:94.

In one embodiment, the present invention provides a composition  
25 comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ  
ID NO:96;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ  
30 ID NO:96 from nucleotide 294 to nucleotide 431;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone BG255\_1i deposited under accession number ATCC 98191;
- (d) a polynucleotide encoding the full-length protein encoded



by the cDNA insert of clone BG255\_1i deposited under accession number ATCC 98191;

5 (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone BG255\_1i deposited under accession number ATCC 98191;

(f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone BG255\_1i deposited under accession number ATCC 98191;

(g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:97;

10 (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:97 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:97;

(i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above; and

15 (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:96 from nucleotide 294 to nucleotide 431; the nucleotide sequence of the full-length protein coding sequence of clone BG255\_1i deposited under accession number ATCC 98191; or the nucleotide sequence of a mature protein coding sequence of clone BG255\_1i deposited under accession number ATCC 98191. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone BG255\_1i deposited under accession number ATCC 98191.

25 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:97;  
(b) fragments of the amino acid sequence of SEQ ID NO:97, each fragment comprising eight consecutive amino acids of SEQ ID NO:97; and  
30 (c) the amino acid sequence encoded by the cDNA insert of clone BG255\_1i deposited under accession number ATCC 98191;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:97. In further preferred embodiments, the present invention provides a protein comprising a fragment of the

amino acid sequence of SEQ ID NO:97 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:97, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:97, the fragment comprising the amino acid sequence from amino acid 18 to amino acid 27 of SEQ ID NO:97.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:99;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:99 from nucleotide 57 to nucleotide 968;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:99 from nucleotide 105 to nucleotide 968;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone H541\_3i deposited under accession number ATCC 98191;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone H541\_3i deposited under accession number ATCC 98191;
- (f) a polynucleotide comprising the nucleotide sequence of a *mature protein coding sequence* of clone H541\_3i deposited under accession number ATCC 98191;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone H541\_3i deposited under accession number ATCC 98191;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:100;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:100 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:100;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above; and
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:99 from nucleotide 57 to nucleotide 968; the nucleotide sequence of SEQ ID NO:99 from nucleotide 105 to nucleotide 968; the nucleotide sequence of the full-length protein coding sequence of clone H541\_3i deposited under accession number ATCC 98191; or the nucleotide sequence of a mature protein coding sequence of clone H541\_3i deposited under accession number ATCC 98191. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone H541\_3i deposited under accession number ATCC 98191.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:100;
- (b) fragments of the amino acid sequence of SEQ ID NO:100, each fragment comprising eight consecutive amino acids of SEQ ID NO:100; and
- (c) the amino acid sequence encoded by the cDNA insert of clone H541\_3i deposited under accession number ATCC 98191;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:100. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:100 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:100, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:100, the fragment comprising the amino acid sequence from amino acid 147 to amino acid 156 of SEQ ID NO:100.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:101;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:101 from nucleotide 37 to nucleotide 220;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone H978\_1i deposited under accession number ATCC 98191;

- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone H978\_1i deposited under accession number ATCC 98191;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone H978\_1i deposited under accession number ATCC 98191;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone H978\_1i deposited under accession number ATCC 98191;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:102;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:102 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:102;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above; and
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:101 from nucleotide 37 to nucleotide 220; the nucleotide sequence of the full-length protein coding sequence of clone H978\_1i deposited under accession number ATCC 98191; or the nucleotide sequence of a mature protein coding sequence of clone H978\_1i deposited under accession number ATCC 98191. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone H978\_1i deposited under accession number ATCC 98191. In yet other preferred embodiments, such polynucleotide encodes a protein comprising the amino acid sequence of SEQ ID NO:102 from amino acid 1 to amino acid 31. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:102 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:102, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:102 having biological activity, the fragment comprising the amino acid sequence from amino acid 25 to amino acid 34 of SEQ ID NO:102.

In other embodiments, the present invention provides a composition

comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:102;
  - (b) the amino acid sequence of SEQ ID NO:102 from amino acid  
5 1 to amino acid 31;
  - (c) fragments of the amino acid sequence of SEQ ID NO:102,  
each fragment comprising eight consecutive amino acids of SEQ ID NO:102; and
  - (d) the amino acid sequence encoded by the cDNA insert of  
clone H978\_1i deposited under accession number ATCC 98191;
- 10 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:102 or the amino acid sequence of SEQ ID NO:102 from amino acid 1 to amino acid 31. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:102 having biological activity, the fragment preferably comprising  
15 eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:102, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:102, the fragment comprising the amino acid sequence from amino acid 25 to amino acid 34 of SEQ ID NO:102.

In one embodiment, the present invention provides a composition  
20 comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ  
ID NO:104;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ  
25 ID NO:104 from nucleotide 2 to nucleotide 422;
- (c) a polynucleotide comprising the nucleotide sequence of the  
full-length protein coding sequence of clone L161\_1i deposited under accession  
number ATCC 98191;
- (d) a polynucleotide encoding the full-length protein encoded  
30 by the cDNA insert of clone L161\_1i deposited under accession number ATCC  
98191;
- (e) a polynucleotide comprising the nucleotide sequence of a  
mature protein coding sequence of clone L161\_1i deposited under accession  
number ATCC 98191;

- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone L161\_1i deposited under accession number ATCC 98191;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:105;
- 5 (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:105 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:105;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above; and
- 10 (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:104 from nucleotide 2 to nucleotide 422; the nucleotide sequence of the full-length protein coding sequence of clone L161\_1i deposited under accession number ATCC 98191;

15 or the nucleotide sequence of a mature protein coding sequence of clone L161\_1i deposited under accession number ATCC 98191. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone L161\_1i deposited under accession number ATCC 98191. In yet other preferred embodiments, such polynucleotide encodes a protein comprising the amino acid sequence

20 of SEQ ID NO:105 from amino acid 72 to amino acid 91. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:105 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:105, or a polynucleotide

25 encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:105 having biological activity, the fragment comprising the amino acid sequence from amino acid 65 to amino acid 74 of SEQ ID NO:105.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected

30 from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:105;
- (b) the amino acid sequence of SEQ ID NO:105 from amino acid 72 to amino acid 91;
- (c) fragments of the amino acid sequence of SEQ ID NO:105,

each fragment comprising eight consecutive amino acids of SEQ ID NO:105; and

(d) the amino acid sequence encoded by the cDNA insert of clone L161\_1i deposited under accession number ATCC 98191;

the protein being substantially free from other mammalian proteins. Preferably such

5 protein comprises the amino acid sequence of SEQ ID NO:105 or the amino acid sequence of SEQ ID NO:105 from amino acid 72 to amino acid 91. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:105 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino  
10 acids of SEQ ID NO:105, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:105, the fragment comprising the amino acid sequence from amino acid 65 to amino acid 74 of SEQ ID NO:105.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group  
15 consisting of:

(a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:107;

(b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:107 from nucleotide 73 to nucleotide 702;

20 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:107 from nucleotide 118 to nucleotide 702;

(d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AE648\_1i deposited under accession number ATCC 98237;

25 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AE648\_1i deposited under accession number ATCC 98237;

(f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AE648\_1i deposited under accession  
30 number ATCC 98237;

(g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AE648\_1i deposited under accession number ATCC 98237;

(h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:108;

(i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:108 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:108;

5 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above; and

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:107 from nucleotide 73 to nucleotide 702; the nucleotide sequence of SEQ ID NO:107 from nucleotide 118 to nucleotide 702; the nucleotide sequence of the full-length protein coding sequence of clone AE648\_1i deposited under accession number ATCC 98237; or the nucleotide sequence of a mature protein coding sequence of clone AE648\_1i deposited under accession number ATCC 98237. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AE648\_1i deposited under accession number ATCC 98237. In yet other preferred  
10 embodiments, such polynucleotide encodes a protein comprising the amino acid sequence of SEQ ID NO:108 from amino acid 1 to amino acid 34. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:108 having biological activity, the  
15 fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:108, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:108 having biological activity, the fragment comprising the amino acid sequence from amino acid 100 to amino acid 109 of SEQ ID NO:108.

25 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:108;
- (b) the amino acid sequence of SEQ ID NO:108 from amino acid  
30 1 to amino acid 34;
- (c) fragments of the amino acid sequence of SEQ ID NO:108, each fragment comprising eight consecutive amino acids of SEQ ID NO:108; and
- (d) the amino acid sequence encoded by the cDNA insert of clone AE648\_1i deposited under accession number ATCC 98237;



the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:108 or the amino acid sequence of SEQ ID NO:108 from amino acid 1 to amino acid 34. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid  
5 sequence of SEQ ID NO:108 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:108, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:108, the fragment comprising the amino acid sequence from amino acid 100 to amino acid 109 of SEQ ID NO:108.

10 In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:109;
- 15 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:109 from nucleotide 92 to nucleotide 268;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AE693\_1i deposited under accession number ATCC 98237;
- 20 (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AE693\_1i deposited under accession number ATCC 98237;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AE693\_1i deposited under accession  
25 number ATCC 98237;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AE693\_1i deposited under accession number ATCC 98237;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:110;
- 30 (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:110 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:110;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above; and

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:109 from nucleotide 92 to nucleotide 268; the nucleotide sequence of the full-length protein coding sequence of clone AE693\_1i deposited under accession number ATCC 98237; or the nucleotide sequence of a mature protein coding sequence of clone AE693\_1i deposited under accession number ATCC 98237. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AE693\_1i deposited under accession number ATCC 98237.

10 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:110;
- (b) fragments of the amino acid sequence of SEQ ID NO:110,
- 15 each fragment comprising eight consecutive amino acids of SEQ ID NO:110; and
- (c) the amino acid sequence encoded by the cDNA insert of clone AE693\_1i deposited under accession number ATCC 98237;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:110. In further preferred  
20 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:110 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:110, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:110, the fragment comprising the amino acid sequence from amino acid 24  
25 to amino acid 33 of SEQ ID NO:110.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ  
30 ID NO:112;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:112 from nucleotide 137 to nucleotide 412;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AK438\_1i deposited under accession

number ATCC 98237;

(d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AK438\_1i deposited under accession number ATCC 98237;

5 (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AK438\_1i deposited under accession number ATCC 98237;

(f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AK438\_1i deposited under accession number ATCC 98237;

10 (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:113;

(h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:113 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:113;

15 (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above; and

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:112 from nucleotide 137 to nucleotide 412; the nucleotide sequence of the full-length protein coding sequence of clone AK438\_1i deposited under accession number ATCC 98237; or the nucleotide sequence of a mature protein coding sequence of clone AK438\_1i deposited under accession number ATCC 98237. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AK438\_1i deposited under accession number ATCC 98237.

25 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:113;
- 30 (b) fragments of the amino acid sequence of SEQ ID NO:113, each fragment comprising eight consecutive amino acids of SEQ ID NO:113; and
- (c) the amino acid sequence encoded by the cDNA insert of clone AK438\_1i deposited under accession number ATCC 98237;

the protein being substantially free from other mammalian proteins. Preferably such

protein comprises the amino acid sequence of SEQ ID NO:113. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:113 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:113, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:113, the fragment comprising the amino acid sequence from amino acid 41 to amino acid 50 of SEQ ID NO:113.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:115;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:115 from nucleotide 1 to nucleotide 285;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AK609\_1i deposited under accession number ATCC 98237;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AK609\_1i deposited under accession number ATCC 98237;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AK609\_1i deposited under accession number ATCC 98237;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AK609\_1i deposited under accession number ATCC 98237;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:116;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:116 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:116;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above; and
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:115 from nucleotide 1 to nucleotide 285; the nucleotide sequence of the full-length protein coding sequence of clone AK609\_1i deposited under accession number ATCC 98237; or the nucleotide sequence of a mature protein coding sequence of clone AK609\_1i deposited under accession number ATCC 98237. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AK609\_1i deposited under accession number ATCC 98237.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:116;
  - (b) fragments of the amino acid sequence of SEQ ID NO:116, each fragment comprising eight consecutive amino acids of SEQ ID NO:116; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone AK609\_1i deposited under accession number ATCC 98237;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:116. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:116 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:116, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:116, the fragment comprising the amino acid sequence from amino acid 42 to amino acid 51 of SEQ ID NO:116.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:118;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:118 from nucleotide 43 to nucleotide 282;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:118 from nucleotide 118 to nucleotide 282;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AM1060\_1i deposited under

accession number ATCC 98237;

(e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AM1060\_1i deposited under accession number ATCC 98237;

5 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AM1060\_1i deposited under accession number ATCC 98237;

(g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AM1060\_1i deposited under accession number ATCC 98237;

10 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:119;

(i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:119 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:119;

15 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above; and

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:118 from nucleotide 43 to nucleotide 282; the nucleotide sequence of SEQ ID NO:118 from nucleotide 118 to nucleotide 282; the nucleotide sequence of the full-length protein coding sequence of clone AM1060\_1i deposited under accession number ATCC 98237; or the nucleotide sequence of a mature protein coding sequence of clone AM1060\_1i deposited under accession number ATCC 98237. In other preferred  
25 embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AM1060\_1i deposited under accession number ATCC 98237.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

30 (a) the amino acid sequence of SEQ ID NO:119;  
(b) fragments of the amino acid sequence of SEQ ID NO:119, each fragment comprising eight consecutive amino acids of SEQ ID NO:119; and  
(c) the amino acid sequence encoded by the cDNA insert of clone AM1060\_1i deposited under accession number ATCC 98237;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:119. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:119 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:119, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:119, the fragment comprising the amino acid sequence from amino acid 35 to amino acid 44 of SEQ ID NO:119.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:121;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:121 from nucleotide 316 to nucleotide 438;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AQ2\_1i deposited under accession number ATCC 98237;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AQ2\_1i deposited under accession number ATCC 98237;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AQ2\_1i deposited under accession number ATCC 98237;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AQ2\_1i deposited under accession number ATCC 98237;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:122;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:122 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:122;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above; and
- (j) a polynucleotide which encodes a species homologue of the

protein of (g) or (h) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:121 from nucleotide 316 to nucleotide 438; the nucleotide sequence of the full-length protein coding sequence of clone AQ2\_1i deposited under accession number  
5 ATCC 98237; or the nucleotide sequence of a mature protein coding sequence of clone AQ2\_1i deposited under accession number ATCC 98237. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AQ2\_1i deposited under accession number ATCC 98237. In yet other preferred embodiments, such polynucleotide encodes a protein comprising the  
10 amino acid sequence of SEQ ID NO:122 from amino acid 1 to amino acid 25. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:122 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:122, or a polynucleotide  
15 encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:122 having biological activity, the fragment comprising the amino acid sequence from amino acid 15 to amino acid 24 of SEQ ID NO:122.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected  
20 from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:122;
- (b) the amino acid sequence of SEQ ID NO:122 from amino acid  
1 to amino acid 25;
- (c) fragments of the amino acid sequence of SEQ ID NO:122,  
25 each fragment comprising eight consecutive amino acids of SEQ ID NO:122; and
- (d) the amino acid sequence encoded by the cDNA insert of clone AQ2\_1i deposited under accession number ATCC 98237;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:122 or the amino acid sequence  
30 of SEQ ID NO:122 from amino acid 1 to amino acid 25. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:122 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:122, or a protein comprising a fragment of the amino acid sequence of SEQ ID



NO:122, the fragment comprising the amino acid sequence from amino acid 15 to amino acid 24 of SEQ ID NO:122.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group  
5 consisting of:

(a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:124;

(b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:124 from nucleotide 142 to nucleotide 285;

10 (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone K433\_1i deposited under accession number ATCC 98237;

(d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone K433\_1i deposited under accession number ATCC  
15 98237;

(e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone K433\_1i deposited under accession number ATCC 98237;

20 (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone K433\_1i deposited under accession number ATCC 98237;

(g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:125;

(h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:125 having biological activity, the  
25 fragment comprising eight consecutive amino acids of SEQ ID NO:125;

(i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above; and

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above.

30 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:124 from nucleotide 142 to nucleotide 285; the nucleotide sequence of the full-length protein coding sequence of clone K433\_1i deposited under accession number ATCC 98237; or the nucleotide sequence of a mature protein coding sequence of clone K433\_1i deposited under accession number ATCC 98237. In other preferred

embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone K433\_1i deposited under accession number ATCC 98237. In yet other preferred embodiments, such polynucleotide encodes a protein comprising the amino acid sequence of SEQ ID NO:125 from amino acid 1 to amino acid 30. In further  
5 preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:125 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:125, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:125  
10 having biological activity, the fragment comprising the amino acid sequence from amino acid 19 to amino acid 28 of SEQ ID NO:125.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 15 (a) the amino acid sequence of SEQ ID NO:125;
- (b) the amino acid sequence of SEQ ID NO:125 from amino acid 1 to amino acid 30;
- (c) fragments of the amino acid sequence of SEQ ID NO:125, each fragment comprising eight consecutive amino acids of SEQ ID NO:125; and
- 20 (d) the amino acid sequence encoded by the cDNA insert of clone K433\_1i deposited under accession number ATCC 98237;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:125 or the amino acid sequence of SEQ ID NO:125 from amino acid 1 to amino acid 30. In further preferred embodiments,  
25 the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:125 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:125, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:125, the fragment comprising the amino acid sequence from amino acid 19 to amino  
30 acid 28 of SEQ ID NO:125.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ

ID NO:127;

(b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:127 from nucleotide 47 to nucleotide 517;

5 (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone L256\_1i deposited under accession number ATCC 98237;

(d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone L256\_1i deposited under accession number ATCC 98237;

10 (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone L256\_1i deposited under accession number ATCC 98237;

(f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone L256\_1i deposited under accession number ATCC 98237;

15 (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:128;

(h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:128 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:128;

20 (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above; and

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:127 from nucleotide 47 to nucleotide 517; the nucleotide sequence of the full-length protein coding sequence of clone L256\_1i deposited under accession number ATCC 98237; or the nucleotide sequence of a mature protein coding sequence of clone L256\_1i deposited under accession number ATCC 98237. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone L256\_1i deposited under accession number ATCC 98237. In yet other preferred  
30 embodiments, such polynucleotide encodes a protein comprising the amino acid sequence of SEQ ID NO:128 from amino acid 8 to amino acid 157. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:128 having biological

activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:128, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:128 having biological activity, the fragment comprising the amino acid sequence from amino acid 73 to amino acid 82 of SEQ ID NO:128.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:128;
- 10 (b) the amino acid sequence of SEQ ID NO:128 from amino acid 8 to amino acid 157;
- (c) fragments of the amino acid sequence of SEQ ID NO:128, each fragment comprising eight consecutive amino acids of SEQ ID NO:128; and
- (d) the amino acid sequence encoded by the cDNA insert of
- 15 clone L256\_1i deposited under accession number ATCC 98237;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:128 or the amino acid sequence of SEQ ID NO:128 from amino acid 8 to amino acid 157. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:128 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:128, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:128, the fragment comprising the amino acid sequence from amino acid 73 to amino acid 82 of SEQ ID NO:128.

25 In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:130;
- 30 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:130 from nucleotide 389 to nucleotide 694;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AM207\_1i deposited under accession number ATCC 98510;

(d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AM207\_1i deposited under accession number ATCC 98510;

5 (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AM207\_1i deposited under accession number ATCC 98510;

(f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AM207\_1i deposited under accession number ATCC 98510;

10 (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:131;

(h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:131 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:131;

15 (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above; and

(j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:130 from nucleotide 389 to nucleotide 694; the nucleotide sequence of the full-length protein coding sequence of clone AM207\_1i deposited under accession number ATCC 98510; or the nucleotide sequence of a mature protein coding sequence of clone AM207\_1i deposited under accession number ATCC 98510. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AM207\_1i deposited under accession number ATCC 98510.

25 In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 30 (a) the amino acid sequence of SEQ ID NO:131;  
(b) fragments of the amino acid sequence of SEQ ID NO:131, each fragment comprising eight consecutive amino acids of SEQ ID NO:131; and  
(c) the amino acid sequence encoded by the cDNA insert of clone AM207\_1i deposited under accession number ATCC 98510;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:131. In further preferred

embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:131 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:131, or a protein comprising a fragment of the amino acid sequence  
5 of SEQ ID NO:131, the fragment comprising the amino acid sequence from amino acid 46 to amino acid 55 of SEQ ID NO:131.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- 10 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:133;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:133 from nucleotide 122 to nucleotide 685;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ  
15 ID NO:133 from nucleotide 179 to nucleotide 685;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AM910\_1i deposited under accession number ATCC 98510;
- (e) a polynucleotide encoding the full-length protein encoded  
20 by the cDNA insert of clone AM910\_1i deposited under accession number ATCC 98510;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AM910\_1i deposited under accession number ATCC 98510;
- 25 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AM910\_1i deposited under accession number ATCC 98510;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:134;
- (i) a polynucleotide encoding a protein comprising a fragment  
30 of the amino acid sequence of SEQ ID NO:134 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:134;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above; and
- (k) a polynucleotide which encodes a species homologue of the

protein of (h) or (i) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:133 from nucleotide 122 to nucleotide 685; the nucleotide sequence of SEQ ID NO:133 from nucleotide 179 to nucleotide 685; the nucleotide sequence of the full-length protein coding sequence of clone AM910\_1i deposited under accession number ATCC 98510; or the nucleotide sequence of a mature protein coding sequence of clone AM910\_1i deposited under accession number ATCC 98510. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AM910\_1i deposited under accession number ATCC 98510. In yet other preferred  
10 embodiments, such polynucleotide encodes a protein comprising the amino acid sequence of SEQ ID NO:134 from amino acid 85 to amino acid 139. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:134 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:134, or a polynucleotide  
15 encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:134 having biological activity, the fragment comprising the amino acid sequence from amino acid 89 to amino acid 98 of SEQ ID NO:134.

In other embodiments, the present invention provides a composition  
20 comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:134;
- (b) the amino acid sequence of SEQ ID NO:134 from amino acid 85 to amino acid 139;
- 25 (c) fragments of the amino acid sequence of SEQ ID NO:134, each fragment comprising eight consecutive amino acids of SEQ ID NO:134; and
- (d) the amino acid sequence encoded by the cDNA insert of clone AM910\_1i deposited under accession number ATCC 98510;

the protein being substantially free from other mammalian proteins. Preferably such  
30 protein comprises the amino acid sequence of SEQ ID NO:134 or the amino acid sequence of SEQ ID NO:134 from amino acid 85 to amino acid 139. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:134 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino

acids of SEQ ID NO:134, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:134, the fragment comprising the amino acid sequence from amino acid 89 to amino acid 98 of SEQ ID NO:134.

In one embodiment, the present invention provides a composition  
5 comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:135;
- (b) a polynucleotide comprising the nucleotide sequence of  
10 SEQ ID NO:135 from nucleotide 84 to nucleotide 269;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:135 from nucleotide 144 to nucleotide 269;
- (d) a polynucleotide comprising the nucleotide sequence of the  
15 full-length protein coding sequence of clone AR54\_1i deposited under accession number ATCC 98510;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AR54\_1i deposited under accession number ATCC 98510;
- (f) a polynucleotide comprising the nucleotide sequence of a  
20 mature protein coding sequence of clone AR54\_1i deposited under accession number ATCC 98510;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AR54\_1i deposited under accession number ATCC 98510;
- (h) a polynucleotide encoding a protein comprising the amino  
25 acid sequence of SEQ ID NO:136;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:136 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:136;
- (j) a polynucleotide which is an allelic variant of a  
30 polynucleotide of (a)-(g) above; and
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:135 from nucleotide 84 to nucleotide 269; the nucleotide sequence of SEQ ID



NO:135 from nucleotide 144 to nucleotide 269; the nucleotide sequence of the full-length protein coding sequence of clone AR54\_1i deposited under accession number ATCC 98510; or the nucleotide sequence of a mature protein coding sequence of clone AR54\_1i deposited under accession number ATCC 98510. In other preferred embodiments, the  
5 polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AR54\_1i deposited under accession number ATCC 98510.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 10 (a) the amino acid sequence of SEQ ID NO:136;
  - (b) fragments of the amino acid sequence of SEQ ID NO:136, each fragment comprising eight consecutive amino acids of SEQ ID NO:136; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone AR54\_1i deposited under accession number ATCC 98510;
- 15 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:136. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:136 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino  
20 acids of SEQ ID NO:136, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:136, the fragment comprising the amino acid sequence from amino acid 26 to amino acid 35 of SEQ ID NO:136.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group  
25 consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:137;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:137 from nucleotide 32 to nucleotide 1300;
- 30 (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:137 from nucleotide 884 to nucleotide 1300;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone L200\_1i deposited under accession number ATCC 98510;

(e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone L200\_1i deposited under accession number ATCC 98510;

5 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone L200\_1i deposited under accession number ATCC 98510;

(g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone L200\_1i deposited under accession number ATCC 98510;

10 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:138;

(i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:138 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:138;

15 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above; and

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:137 from nucleotide 32 to nucleotide 1300; the nucleotide sequence of SEQ ID NO:137 from nucleotide 884 to nucleotide 1300; the nucleotide sequence of the full-length protein coding sequence of clone L200\_1i deposited under accession number ATCC 98510; or the nucleotide sequence of a mature protein coding sequence of clone L200\_1i deposited under accession number ATCC 98510. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone L200\_1i deposited under accession number ATCC 98510. In yet other preferred  
25 embodiments, such polynucleotide encodes a protein comprising the amino acid sequence of SEQ ID NO:138 from amino acid 1 to amino acid 144. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:138 having biological  
30 activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:138, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:138 having biological activity, the fragment comprising the amino acid sequence from amino acid 206 to amino acid 215 of SEQ ID NO:138.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:138;
  - 5 (b) the amino acid sequence of SEQ ID NO:138 from amino acid 1 to amino acid 144;
  - (c) fragments of the amino acid sequence of SEQ ID NO:138, each fragment comprising eight consecutive amino acids of SEQ ID NO:138; and
  - (d) the amino acid sequence encoded by the cDNA insert of  
10 clone L200\_1i deposited under accession number ATCC 98510;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:138 or the amino acid sequence of SEQ ID NO:138 from amino acid 1 to amino acid 144. In further preferred  
15 amino acid sequence of SEQ ID NO:138 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:138, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:138, the fragment comprising the amino acid sequence from amino acid 206 to amino acid 215 of SEQ ID NO:138.

20 In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:139;
- 25 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:139 from nucleotide 85 to nucleotide 1059;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:139 from nucleotide 151 to nucleotide 1059;
- (d) a polynucleotide comprising the nucleotide sequence of the  
30 full-length protein coding sequence of clone WA129\_2i deposited under accession number ATCC 98510;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone WA129\_2i deposited under accession number ATCC 98510;

(f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone WA129\_2i deposited under accession number ATCC 98510;

5 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone WA129\_2i deposited under accession number ATCC 98510;

(h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:140;

10 (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:140 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:140;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above; and

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above.

15 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:139 from nucleotide 85 to nucleotide 1059; the nucleotide sequence of SEQ ID NO:139 from nucleotide 151 to nucleotide 1059; the nucleotide sequence of the full-length protein coding sequence of clone WA129\_2i deposited under accession number ATCC 98510; or the nucleotide sequence of a mature protein coding sequence of clone WA129\_2i  
20 deposited under accession number ATCC 98510. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone WA129\_2i deposited under accession number ATCC 98510.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected  
25 from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:140;

(b) fragments of the amino acid sequence of SEQ ID NO:140, each fragment comprising eight consecutive amino acids of SEQ ID NO:140; and

30 (c) the amino acid sequence encoded by the cDNA insert of clone WA129\_2i deposited under accession number ATCC 98510;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:140. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:140 having biological activity, the fragment preferably

comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:140, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:140, the fragment comprising the amino acid sequence from amino acid 157 to amino acid 166 of SEQ ID NO:140.

5 In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:141;
- 10 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:141 from nucleotide 128 to nucleotide 643;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:141 from nucleotide 197 to nucleotide 643;
- (d) a polynucleotide comprising the nucleotide sequence of the  
15 full-length protein coding sequence of clone WA154\_3i deposited under accession number ATCC 98510;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone WA154\_3i deposited under accession number ATCC 98510;
- 20 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone WA154\_3i deposited under accession number ATCC 98510;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone WA154\_3i deposited under accession number ATCC 98510;
- 25 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:142;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:142 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:142;
- 30 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above; and
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ

ID NO:141 from nucleotide 128 to nucleotide 643; the nucleotide sequence of SEQ ID NO:141 from nucleotide 197 to nucleotide 643; the nucleotide sequence of the full-length protein coding sequence of clone WA154\_3i deposited under accession number ATCC 98510; or the nucleotide sequence of a mature protein coding sequence of clone WA154\_3i deposited under accession number ATCC 98510. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone WA154\_3i deposited under accession number ATCC 98510. In yet other preferred embodiments, such polynucleotide encodes a protein comprising the amino acid sequence of SEQ ID NO:142 from amino acid 37 to amino acid 77. In further preferred  
10 embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:142 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:142, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:142  
15 having biological activity, the fragment comprising the amino acid sequence from amino acid 81 to amino acid 90 of SEQ ID NO:142.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- 20 (a) the amino acid sequence of SEQ ID NO:142;
- (b) the amino acid sequence of SEQ ID NO:142 from amino acid 37 to amino acid 77;
- (c) fragments of the amino acid sequence of SEQ ID NO:142, each fragment comprising eight consecutive amino acids of SEQ ID NO:142; and
- 25 (d) the amino acid sequence encoded by the cDNA insert of clone WA154\_3i deposited under accession number ATCC 98510;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:142 or the amino acid sequence of SEQ ID NO:142 from amino acid 37 to amino acid 77. In further preferred  
30 embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:142 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:142, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:142, the fragment comprising the amino acid sequence from amino acid 81

to amino acid 90 of SEQ ID NO:142.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- 5 (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:143;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:143 from nucleotide 51 to nucleotide 815;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ  
10 ID NO:143 from nucleotide 96 to nucleotide 815;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AA36\_1i deposited under accession number ATCC XXXXX;
- (e) a polynucleotide encoding the full-length protein encoded  
15 by the cDNA insert of clone AA36\_1i deposited under accession number ATCC XXXXX;
- (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AA36\_1i deposited under accession number ATCC XXXXX;
- 20 (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AA36\_1i deposited under accession number ATCC XXXXX;
- (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:144;
- (i) a polynucleotide encoding a protein comprising a fragment  
25 of the amino acid sequence of SEQ ID NO:144 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:144;
- (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above; and
- (k) a polynucleotide which encodes a species homologue of the  
30 protein of (h) or (i) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:143 from nucleotide 51 to nucleotide 815; the nucleotide sequence of SEQ ID NO:143 from nucleotide 96 to nucleotide 815; the nucleotide sequence of the full-length protein coding sequence of clone AA36\_1i deposited under accession number ATCC

XXXXX; or the nucleotide sequence of a mature protein coding sequence of clone AA36\_1i deposited under accession number ATCC XXXXX. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AA36\_1i deposited under accession number ATCC XXXXX. In yet other preferred  
5 embodiments, such polynucleotide encodes a protein comprising the amino acid sequence of SEQ ID NO:144 from amino acid 1 to amino acid 136. In further preferred embodiments, the present invention provides a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:144 having biological activity, the fragment preferably comprising eight (more preferably twenty, most  
10 preferably thirty) consecutive amino acids of SEQ ID NO:144, or a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:144 having biological activity, the fragment comprising the amino acid sequence from amino acid 122 to amino acid 131 of SEQ ID NO:144.

In other embodiments, the present invention provides a composition  
15 comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:144;
- (b) the amino acid sequence of SEQ ID NO:144 from amino acid  
20 1 to amino acid 136;
- (c) fragments of the amino acid sequence of SEQ ID NO:144, each fragment comprising eight consecutive amino acids of SEQ ID NO:144; and
- (d) the amino acid sequence encoded by the cDNA insert of  
clone AA36\_1i deposited under accession number ATCC XXXXX;

the protein being substantially free from other mammalian proteins. Preferably such  
25 protein comprises the amino acid sequence of SEQ ID NO:144 or the amino acid sequence of SEQ ID NO:144 from amino acid 1 to amino acid 136. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:144 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino  
30 acids of SEQ ID NO:144, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:144, the fragment comprising the amino acid sequence from amino acid 122 to amino acid 131 of SEQ ID NO:144.

In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group



consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:145;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:145 from nucleotide 76 to nucleotide 594;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AC175\_2i deposited under accession number ATCC XXXXX;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AC175\_2i deposited under accession number ATCC XXXXX;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AC175\_2i deposited under accession number ATCC XXXXX;
- (f) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AC175\_2i deposited under accession number ATCC XXXXX;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:146;
- (h) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:146 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:146;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above; and
- (j) a polynucleotide which encodes a species homologue of the protein of (g) or (h) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:145 from nucleotide 76 to nucleotide 594; the nucleotide sequence of the full-length protein coding sequence of clone AC175\_2i deposited under accession number ATCC XXXXX; or the nucleotide sequence of a mature protein coding sequence of clone AC175\_2i deposited under accession number ATCC XXXXX. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AC175\_2i deposited under accession number ATCC XXXXX.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected

from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:146;
- (b) fragments of the amino acid sequence of SEQ ID NO:146, each fragment comprising eight consecutive amino acids of SEQ ID NO:146; and
- 5 (c) the amino acid sequence encoded by the cDNA insert of clone AC175\_2i deposited under accession number ATCC XXXXX;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:146. In further preferred

10 amino acid sequence of SEQ ID NO:146 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:146, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:146, the fragment comprising the amino acid sequence from amino acid 81 to amino acid 90 of SEQ ID NO:146.

15 In one embodiment, the present invention provides a composition comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:147;
- 20 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:147 from nucleotide 387 to nucleotide 734;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:147 from nucleotide 639 to nucleotide 734;
- (d) a polynucleotide comprising the nucleotide sequence of the
- 25 full-length protein coding sequence of clone AV189\_1i deposited under accession number ATCC XXXXX;
- (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AV189\_1i deposited under accession number ATCC XXXXX;
- 30 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AV189\_1i deposited under accession number ATCC XXXXX;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AV189\_1i deposited under accession number ATCC XXXXX;

(h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:148;

(i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:148 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:148;

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above; and

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above.

10 Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:147 from nucleotide 387 to nucleotide 734; the nucleotide sequence of SEQ ID NO:147 from nucleotide 639 to nucleotide 734; the nucleotide sequence of the full-length protein coding sequence of clone AV189\_1i deposited under accession number ATCC XXXXX; or the nucleotide sequence of a mature protein coding sequence of clone  
15 AV189\_1i deposited under accession number ATCC XXXXX. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone AV189\_1i deposited under accession number ATCC XXXXX.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected  
20 from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:148;

(b) fragments of the amino acid sequence of SEQ ID NO:148, each fragment comprising eight consecutive amino acids of SEQ ID NO:148; and

(c) the amino acid sequence encoded by the cDNA insert of  
25 clone AV189\_1i deposited under accession number ATCC XXXXX;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:148. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:148 having biological activity, the fragment preferably  
30 comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:148, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:148, the fragment comprising the amino acid sequence from amino acid 53 to amino acid 62 of SEQ ID NO:148.

In one embodiment, the present invention provides a composition

comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:149;
- 5 (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:149 from nucleotide 66 to nucleotide 827;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:149 from nucleotide 366 to nucleotide 827;
- (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone K368\_1i deposited under accession number ATCC XXXXX;
- 10 (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone K368\_1i deposited under accession number ATCC XXXXX;
- 15 (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone K368\_1i deposited under accession number ATCC XXXXX;
- (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone K368\_1i deposited under accession number ATCC XXXXX;
- 20 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:150;
- (i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:150 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:150;
- 25 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above; and
- (k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:149 from nucleotide 66 to nucleotide 827; the nucleotide sequence of SEQ ID NO:149 from nucleotide 366 to nucleotide 827; the nucleotide sequence of the full-length protein coding sequence of clone K368\_1i deposited under accession number ATCC XXXXX; or the nucleotide sequence of a mature protein coding sequence of clone K368\_1i deposited under accession number ATCC XXXXX. In other preferred embodiments, the

polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone K368\_1i deposited under accession number ATCC XXXXX.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected  
5 from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:150;
  - (b) *fragments of the amino acid sequence of SEQ ID NO:150*,  
each fragment comprising eight consecutive amino acids of SEQ ID NO:150; and
  - (c) the amino acid sequence encoded by the cDNA insert of  
10 clone K368\_1i deposited under accession number ATCC XXXXX;
- the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:150. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:150 having biological activity, the fragment preferably  
15 comprising eight (more preferably twenty, most preferably thirty) consecutive amino acids of SEQ ID NO:150, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:150, the fragment comprising the amino acid sequence from amino acid 122 to amino acid 131 of SEQ ID NO:150.

In one embodiment, the present invention provides a composition  
20 comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ  
ID NO:151;
- (b) a polynucleotide comprising the nucleotide sequence of  
25 SEQ ID NO:151 from nucleotide 219 to nucleotide 668;
- (c) a polynucleotide comprising the nucleotide sequence of SEQ  
ID NO:151 from nucleotide 426 to nucleotide 668;
- (d) a polynucleotide comprising the nucleotide sequence of the  
full-length protein coding sequence of clone K568\_1i deposited under accession  
30 number ATCC XXXXX;
- (e) a polynucleotide encoding the full-length protein encoded  
by the cDNA insert of clone K568\_1i deposited under accession number ATCC  
XXXXX;
- (f) a polynucleotide comprising the nucleotide sequence of a

mature protein coding sequence of clone K568\_1i deposited under accession number ATCC XXXXX;

(g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone K568\_1i deposited under accession number ATCC XXXXX;

5 (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:152;

(i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:152 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:152;

10 (j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above; and

(k) a polynucleotide which encodes a species homologue of the protein of (h) or (i) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:151 from nucleotide 219 to nucleotide 668; the nucleotide sequence of SEQ ID NO:151 from nucleotide 426 to nucleotide 668; the nucleotide sequence of the full-length protein coding sequence of clone K568\_1i deposited under accession number ATCC XXXXX; or the nucleotide sequence of a mature protein coding sequence of clone K568\_1i deposited under accession number ATCC XXXXX. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone K568\_1i deposited under accession number ATCC XXXXX.

In other embodiments, the present invention provides a composition comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

25 (a) the amino acid sequence of SEQ ID NO:152;  
(b) fragments of the amino acid sequence of SEQ ID NO:152, each fragment comprising eight consecutive amino acids of SEQ ID NO:152; and  
(c) the amino acid sequence encoded by the cDNA insert of clone K568\_1i deposited under accession number ATCC XXXXX;

30 the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:152. In further preferred embodiments, the present invention provides a protein comprising a fragment of the amino acid sequence of SEQ ID NO:152 having biological activity, the fragment preferably comprising eight (more preferably twenty, most preferably thirty) consecutive amino

acids of SEQ ID NO:152, or a protein comprising a fragment of the amino acid sequence of SEQ ID NO:152, the fragment comprising the amino acid sequence from amino acid 70 to amino acid 79 of SEQ ID NO:152.

In one embodiment, the present invention provides a composition  
5 comprising an isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:153;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ  
10 ID NO:153 from nucleotide 14 to nucleotide 1438;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone T85\_1i deposited under accession number ATCC XXXXX;
- (d) a polynucleotide encoding the full-length protein encoded  
15 by the cDNA insert of clone T85\_1i deposited under accession number ATCC XXXXX;
- (e) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone T85\_1i deposited under accession number ATCC XXXXX;
- (f) a polynucleotide encoding a mature protein encoded by the  
20 cDNA insert of clone T85\_1i deposited under accession number ATCC XXXXX;
- (g) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:154;
- (h) a polynucleotide encoding a protein comprising a fragment  
25 of the amino acid sequence of SEQ ID NO:154 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:154;
- (i) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above; and
- (j) a polynucleotide which encodes a species homologue of the  
30 protein of (g) or (h) above.

Preferably, such polynucleotide comprises the nucleotide sequence of SEQ ID NO:153 from nucleotide 14 to nucleotide 1438; the nucleotide sequence of the full-length protein coding sequence of clone T85\_1i deposited under accession number ATCC XXXXX; or the nucleotide sequence of a mature protein coding sequence of clone

T85\_li deposited under accession number ATCC XXXXX. In other preferred embodiments, the polynucleotide encodes the full-length or a mature protein encoded by the cDNA insert of clone T85\_li deposited under accession number ATCC XXXXX.

In other embodiments, the present invention provides a composition  
5 comprising a protein, wherein said protein comprises an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:154;
- (b) fragments of the amino acid sequence of SEQ ID NO:154,  
each fragment comprising eight consecutive amino acids of SEQ ID NO:154; and
- 10 (c) the amino acid sequence encoded by the cDNA insert of  
clone T85\_li deposited under accession number ATCC XXXXX;

the protein being substantially free from other mammalian proteins. Preferably such protein comprises the amino acid sequence of SEQ ID NO:154. In further preferred  
embodiments, the present invention provides a protein comprising a fragment of the  
15 amino acid sequence of SEQ ID NO:154 having biological activity, the fragment preferably  
comprising eight (more preferably twenty, most preferably thirty) consecutive amino  
acids of SEQ ID NO:154, or a protein comprising a fragment of the amino acid sequence  
of SEQ ID NO:154, the fragment comprising the amino acid sequence from amino acid 232  
to amino acid 241 of SEQ ID NO:154.

20 Protein compositions of the present invention may further comprise a  
pharmaceutically acceptable carrier. Compositions comprising an antibody which  
specifically reacts with such protein are also provided by the present invention.

Methods are also provided for preventing, treating or ameliorating a  
medical condition which comprises administering to a mammalian subject a  
25 therapeutically effective amount of a composition comprising a protein of the present  
invention and a pharmaceutically acceptable carrier.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1A and 1B are schematic representations of the pED6 and pNOTs  
30 vectors, respectively, used for deposit of clones disclosed herein.

#### DETAILED DESCRIPTION

##### ISOLATED PROTEINS

Nucleotide and amino acid sequences, as presently determined, are



reported below for each clone and protein disclosed in the present application. The nucleotide sequence of each clone can readily be determined by sequencing of the deposited clone in accordance with known methods. The predicted amino acid sequence (both full-length and mature forms) can then be determined from such nucleotide  
5 sequence. The amino acid sequence of the protein encoded by a particular clone can also be determined by expression of the clone in a suitable host cell, collecting the protein and determining its sequence. For each disclosed protein applicants have identified what they have determined to be the reading frame best identifiable with sequence information available at the time of filing.

10 As used herein a "secreted" protein is one which, when expressed in a suitable host cell, is transported across or through a membrane, including transport as a result of signal sequences in its amino acid sequence. "Secreted" proteins include without limitation proteins secreted wholly (e.g., soluble proteins) or partially (e.g., receptors) from the cell in which they are expressed. "Secreted" proteins also include, without  
15 limitation proteins which are transported across the membrane of the endoplasmic reticulum.

#### Protein "AK296\_1i"

One protein of the present invention has been identified as protein  
20 "AK296\_1i". A partial cDNA clone encoding AK296\_1i was first isolated from a human fetal kidney cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of  
25 this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "AK296\_1i".

30 Applicants' methods identified clone AK296\_1i as encoding a secreted protein.

The nucleotide sequence of AK296\_1i as presently determined is reported in SEQ ID NO:1, and includes the poly(A) tail. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AK296\_1i protein

corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:2.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AK296\_1i should be approximately 1264 bp.

AK296\_1i protein was expressed in a Baculovirus expression system, and  
5 an expressed protein band of approximately 20 kDa detected in a membrane fraction using SDS polyacrylamide gel electrophoresis.

Protein "AK533\_1i"

One protein of the present invention has been identified as protein  
10 "AK533\_1i". A partial cDNA clone encoding AK533\_1i was first isolated from a human fetal kidney cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of  
15 this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "AK533\_1i".

20 Applicants' methods identified clone AK533\_1i as encoding a secreted protein.

The nucleotide sequence of AK533\_1i as presently determined is reported in SEQ ID NO:3, and includes the poly(A) tail. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AK533\_1i protein  
25 corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:4.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AK533\_1i should be approximately 1751 bp.

AK533\_1i protein was expressed in a COS cell expression system, and an expressed protein band of approximately 47 kDa detected in conditioned medium and  
30 membrane fractions using SDS polyacrylamide gel electrophoresis.

Protein "AK583\_1i"

One protein of the present invention has been identified as protein "AK583\_1i". A partial cDNA clone encoding AK583\_1i was first isolated from a human

fetal kidney cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "AK583\_1i".

Applicants' methods identified clone AK583\_1i as encoding a secreted protein.

The nucleotide sequence of AK583\_1i as presently determined is reported in SEQ ID NO:5, and includes the poly(A) tail. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AK583\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:6. Amino acids 12 to 24 of SEQ ID NO:6 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 25. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the AK583\_1i protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AK583\_1i should be approximately 870 bp.

#### Protein "AM282\_1i"

One protein of the present invention has been identified as protein "AM282\_1i". A partial cDNA clone encoding AM282\_1i was first isolated from a human fetal kidney cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR,

including a poly(A) tail. This full-length clone is also referred to herein as "AM282\_1i".

Applicants' methods identified clone AM282\_1i as encoding a secreted protein.

The nucleotide sequence of AM282\_1i as presently determined is reported in SEQ ID NO:7, and includes the poly(A) tail. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AM282\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:8. Amino acids 12 to 24 of SEQ ID NO:8 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 25. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the AM282\_1i protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AM282\_1i should be approximately 1750 bp.

AM282\_1i protein was expressed in a Baculovirus expression system, and an expressed protein band of approximately 54 kDa detected in a conditioned medium fraction using SDS polyacrylamide gel electrophoresis.

#### Protein "AM340\_1i"

One protein of the present invention has been identified as protein "AM340\_1i". A partial cDNA clone encoding AM340\_1i was first isolated from a human fetal kidney cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "AM340\_1i".

Applicants' methods identified clone AM340\_1i as encoding a secreted protein.

The nucleotide sequence of AM340\_1i as presently determined is reported in SEQ ID NO:9, and includes the poly(A) tail. What applicants believe is the proper

reading frame and the predicted amino acid sequence of the AM340\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:10. Amino acids 85 to 97 of SEQ ID NO:10 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 98. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the AM340\_1i protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AM340\_1i should be approximately 650 bp.

AM340\_1i protein was expressed in a Baculovirus expression system, and an expressed protein band of approximately 27 kDa detected in a membrane fraction using SDS polyacrylamide gel electrophoresis.

#### Protein "AM610\_1i"

One protein of the present invention has been identified as protein "AM610\_1i". A partial cDNA clone encoding AM610\_1i was first isolated from a human fetal kidney cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "AM610\_1i".

Applicants' methods identified clone AM610\_1i as encoding a secreted protein.

The nucleotide sequence of AM610\_1i as presently determined is reported in SEQ ID NO:11, and includes the poly(A) tail. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AM610\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:12. Amino acids 11 to 23 of SEQ ID NO:12 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 24. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain

should the predicted leader/signal sequence not be separated from the remainder of the AM610\_1i protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AM610\_1i should be approximately 1900 bp.

5 AM610\_1i protein was expressed in a COS cell expression system, and an expressed protein band of approximately 23 kDa detected in conditioned medium and membrane fractions using SDS polyacrylamide gel electrophoresis.

Protein "AP162\_1i"

10 One protein of the present invention has been identified as protein "AP162\_1i". A partial cDNA clone encoding AP162\_1i was first isolated from a human fetal placenta cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of  
15 the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR,  
20 including a poly(A) tail. This full-length clone is also referred to herein as "AP162\_1i".

Applicants' methods identified clone AP162\_1i as encoding a secreted protein.

The nucleotide sequence of the 5' portion of AP162\_1i as presently determined is reported in SEQ ID NO:13. What applicants believe is the proper reading  
25 frame and the predicted amino acid sequence of the AP162\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:14. Additional nucleotide sequence from the 3' portion of AP162\_1i, including a poly(A) tail, is reported in SEQ ID NO:15.

The EcoRI/NotI restriction fragment obtainable from the deposit  
30 containing clone AP162\_1i should be approximately 1200 bp.

AP162\_1i protein was expressed in a Baculovirus expression system, and an expressed protein band of approximately 20 kDa detected in a membrane fraction using SDS polyacrylamide gel electrophoresis.

Protein "AR260\_1i"

One protein of the present invention has been identified as protein "AR260\_1i". A partial cDNA clone encoding AR260\_1i was first isolated from a human adult retina cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "AR260\_1i".

Applicants' methods identified clone AR260\_1i as encoding a secreted protein.

The nucleotide sequence of AR260\_1i as presently determined is reported in SEQ ID NO:16, and includes the poly(A) tail. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AR260\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:17.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AR260\_1i should be approximately 1900 bp.

AR260\_1i protein was expressed in a COS cell expression system, and an expressed protein band of approximately 27 kDa detected in conditioned medium and membrane fractions using SDS polyacrylamide gel electrophoresis.

Protein "AS32\_1i"

One protein of the present invention has been identified as protein "AS32\_1i". A partial cDNA clone encoding AS32\_1i was first isolated from a human fetal brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor

was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "AS32\_1i".

Applicants' methods identified clone AS32\_1i as encoding a secreted protein.

5           The nucleotide sequence of the 5' portion of AS32\_1i as presently determined is reported in SEQ ID NO:18. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AS32\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:19. Additional nucleotide  
10   NO:20.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AS32\_1i should be approximately 1100 bp.

Protein "AS34\_1i"

15           One protein of the present invention has been identified as protein "AS34\_1i". A partial cDNA clone encoding AS34\_1i was first isolated from a human fetal brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or  
20   transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR,  
25   including a poly(A) tail. This full-length clone is also referred to herein as "AS34\_1i".

Applicants' methods identified clone AS34\_1i as encoding a secreted protein.

30           The nucleotide sequence of AS34\_1i as presently determined is reported in SEQ ID NO:21, and includes the poly(A) tail. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AS34\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:22. Amino acids 12 to 24 of SEQ ID NO:22 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 25. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain



should the predicted leader/signal sequence not be separated from the remainder of the AS34\_1i protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AS34\_1i should be approximately 550 bp.

- 5 AS34\_1i protein was expressed in a COS cell expression system, and an expressed protein band of approximately 8 kDa detected in conditioned medium and membrane fractions using SDS polyacrylamide gel electrophoresis.

Protein "AT205\_1i"

- 10 One protein of the present invention has been identified as protein "AT205\_1i". A partial cDNA clone encoding AT205\_1i was first isolated from a human adult blood (lymphocytes and dendritic cells, treated with mixed lymphocyte reaction) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane  
15 protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and  
20 determined to be a full-length clone, including a 5' end and 3' UTR. This full-length clone is also referred to herein as "AT205\_1i".

Applicants' methods identified clone AT205\_1i as encoding a secreted protein.

- The nucleotide sequence of AT205\_1i as presently determined is reported  
25 in SEQ ID NO:23. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AT205\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:24. Amino acids 42 to 54 of SEQ ID NO:24 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 55. Due to the hydrophobic nature of the predicted leader/signal  
30 sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the AT205\_1i protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AT205\_1i should be approximately 825 bp.

Protein "AT211\_1i"

One protein of the present invention has been identified as protein "AT211\_1i". A partial cDNA clone encoding AT211\_1i was first isolated from a human adult blood (lymphocytes and dendritic cells, treated with mixed lymphocyte reaction) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "AT211\_1i".

Applicants' methods identified clone AT211\_1i as encoding a secreted protein.

The nucleotide sequence of AT211\_1i as presently determined is reported in SEQ ID NO:25, and includes the poly(A) tail. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AT211\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:26.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AT211\_1i should be approximately 1100 bp.

Protein "AT319\_1i"

One protein of the present invention has been identified as protein "AT319\_1i". A partial cDNA clone encoding AT319\_1i was first isolated from a human adult blood (lymphocytes and dendritic cells, treated with mixed lymphocyte reaction) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR. This full-length clone

is also referred to herein as "AT319\_1i".

Applicants' methods identified clone AT319\_1i as encoding a secreted protein.

The nucleotide sequence of the 5' portion of AT319\_1i as presently  
5 determined is reported in SEQ ID NO:27. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AT319\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:28. Additional nucleotide sequence from the 3' portion of AT319\_1i is reported in SEQ ID NO:29.

The EcoRI/NotI restriction fragment obtainable from the deposit  
10 containing clone AT319\_1i should be approximately 1680 bp.

#### Protein "AW191\_1i"

One protein of the present invention has been identified as protein "AW191\_1i". A partial cDNA clone encoding AW191\_1i was first isolated from a human  
15 adult ovary (PA-1 teratocarcinoma line, pool of retinoic-acid-treated, activin-treated, and untreated tissue) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of  
20 this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "AW191\_1i".

25 Applicants' methods identified clone AW191\_1i as encoding a secreted protein.

The nucleotide sequence of the 5' portion of AW191\_1i as presently determined is reported in SEQ ID NO:30. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AW191\_1i protein corresponding to  
30 the foregoing nucleotide sequence is reported in SEQ ID NO:31. Amino acids 5 to 17 of SEQ ID NO:31 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 18. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the AW191\_1i protein.

Additional nucleotide sequence from the 3' portion of AW191\_1i, including a poly(A) tail, is reported in SEQ ID NO:32.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AW191\_1i should be approximately 1300 bp.

5

#### Protein "BB9\_1i"

One protein of the present invention has been identified as protein "BB9\_1i". A partial cDNA clone encoding BB9\_1i was first isolated from a human adult blood (peripheral blood mononuclear cells, TH1- or TH2-driven response) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "BB9\_1i".

10

15

20

Applicants' methods identified clone BB9\_1i as encoding a secreted protein.

The nucleotide sequence of the 5' portion of BB9\_1i as presently determined is reported in SEQ ID NO:33. What applicants believe is the proper reading frame and the predicted amino acid sequence of the BB9\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:34. Additional nucleotide sequence from the 3' portion of BB9\_1i, including a poly(A) tail, is reported in SEQ ID NO:35.

25

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone BB9\_1i should be approximately 1080 bp.

#### Protein "H617\_1i"

One protein of the present invention has been identified as protein "H617\_1i". A partial cDNA clone encoding H617\_1i was first isolated from a human adult blood (peripheral blood mononuclear cells, treated with phytohemagglutinin, phorbol myristate acetate, and mixed lymphocyte cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as

30

encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc.,  
5 St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "H617\_1i".

Applicants' methods identified clone H617\_1i as encoding a secreted  
10 protein.

The nucleotide sequence of the 5' portion of H617\_1i as presently determined is reported in SEQ ID NO:36. What applicants believe is the proper reading frame and the predicted amino acid sequence of the H617\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:37. Additional nucleotide  
15 sequence from the 3' portion of H617\_1i, including a poly(A) tail, is reported in SEQ ID NO:38.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone H617\_1i should be approximately 1600 bp.

20 Protein "K39\_1i"

One protein of the present invention has been identified as protein "K39\_1i". A partial cDNA clone encoding K39\_1i was first isolated from a mouse adult bone marrow (stromal line FCM-4) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as  
25 encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from  
30 the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "K39\_1i".

Applicants' methods identified clone K39\_1i as encoding a secreted protein.

The nucleotide sequence of the 5' portion of K39\_1i as presently determined is reported in SEQ ID NO:39. What applicants believe is the proper reading frame and the predicted amino acid sequence of the K39\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:40. Additional nucleotide sequence from the 3' portion of K39\_1i, including a poly(A) tail, is reported in SEQ ID NO:41.

#### Protein "K640\_1i"

One protein of the present invention has been identified as protein "K640\_1i". A partial cDNA clone encoding K640\_1i was first isolated from a mouse adult bone marrow (stromal line FCM-4) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "K640\_1i".

Applicants' methods identified clone K640\_1i as encoding a secreted protein.

The nucleotide sequence of the 5' portion of K640\_1i as presently determined is reported in SEQ ID NO:42. What applicants believe is the proper reading frame and the predicted amino acid sequence of the K640\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:43. Additional nucleotide sequence from the 3' portion of K640\_1i, including a poly(A) tail, is reported in SEQ ID NO:44.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone K640\_1i should be approximately 2400 bp.

#### Protein "AE402\_1i"

One protein of the present invention has been identified as protein "AE402\_1i". A partial cDNA clone encoding AE402\_1i was first isolated from a mouse adult spleen (stimulated with concanavalin A and mixed with dendritic cells) cDNA

library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified  
5 by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "AE402\_1i".

10 Applicants' methods identified clone AE402\_1i as encoding a secreted protein.

The nucleotide sequence of AE402\_1i as presently determined is reported in SEQ ID NO:45, and includes the poly(A) tail. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AE402\_1i protein  
15 corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:46.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AE402\_1i should be approximately 1200 bp.

#### Protein "AE610\_1i"

20 One protein of the present invention has been identified as protein "AE610\_1i". A partial cDNA clone encoding AE610\_1i was first isolated from a mouse adult spleen (stimulated with concanavalin A and mixed with dendritic cells) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified  
25 by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and  
30 determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "AE610\_1i".

Applicants' methods identified clone AE610\_1i as encoding a secreted protein.

The nucleotide sequence of the 5' portion of AE610\_1i as presently

determined is reported in SEQ ID NO:47. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AE610\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:48. Amino acids 75 to 87 of SEQ ID NO:48 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 88. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the AE610\_1i protein. Additional nucleotide sequence from the 3' portion of AE610\_1i, including a poly(A) tail, is reported in SEQ ID NO:49.

10           The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AE610\_1i should be approximately 950 bp.

#### Protein "AH106\_1i"

One protein of the present invention has been identified as protein "AH106\_1i". A partial cDNA clone encoding AH106\_1i was first isolated from a mouse fetal thymus cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "AH106\_1i".

25           Applicants' methods identified clone AH106\_1i as encoding a secreted protein.

The nucleotide sequence of AH106\_1i as presently determined is reported in SEQ ID NO:50, and includes the poly(A) tail. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AH106\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:51.

30           The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AH106\_1i should be approximately 500 bp.

#### Protein "AH196\_1i"



One protein of the present invention has been identified as protein "AH196\_1i". A partial cDNA clone encoding AH196\_1i was first isolated from a mouse fetal thymus cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "AH196\_1i".

Applicants' methods identified clone AH196\_1i as encoding a secreted protein.

The nucleotide sequence of the 5' portion of AH196\_1i as presently determined is reported in SEQ ID NO:52. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AH196\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:53. Additional nucleotide sequence from the 3' portion of AH196\_1i, including a poly(A) tail, is reported in SEQ ID NO:54.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AH196\_1i should be approximately 870 bp.

#### Protein "AI6\_1i"

One protein of the present invention has been identified as protein "AI6\_1i". A partial cDNA clone encoding AI6\_1i was first isolated from a human adult blood (peripheral blood mononuclear cells, TH1- or TH2-driven response) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A)

tail. This full-length clone is also referred to herein as "AI6\_1i".

Applicants' methods identified clone AI6\_1i as encoding a secreted protein.

The nucleotide sequence of the 5' portion of AI6\_1i as presently determined  
5 is reported in SEQ ID NO:55. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AI6\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:56. Additional nucleotide sequence from the 3' portion of AI6\_1i, including a poly(A) tail, is reported in SEQ ID NO:57.

The EcoRI/NotI restriction fragment obtainable from the deposit  
10 containing clone AI6\_1i should be approximately 900 bp.

AI6\_1i protein was expressed in a Baculovirus expression system, and an expressed protein band of approximately 6 kDa detected in a membrane fraction using SDS polyacrylamide gel electrophoresis.

15                   Protein "AJ13\_1i"

One protein of the present invention has been identified as protein  
"AJ13\_1i". A partial cDNA clone encoding AJ13\_1i was first isolated from a human adult  
testes cDNA library using methods which are selective for cDNAs encoding secreted  
proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or  
20 transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor  
25 was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "AJ13\_1i".

Applicants' methods identified clone AJ13\_1i as encoding a secreted protein.

The nucleotide sequence of AJ13\_1i as presently determined is reported in  
30 SEQ ID NO:58, and includes the poly(A) tail. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AJ13\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:59.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AJ13\_1i should be approximately 1200 bp.

Protein "AJ27\_1i"

One protein of the present invention has been identified as protein "AJ27\_1i". A partial cDNA clone encoding AJ27\_1i was first isolated from a human adult testes cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "AJ27\_1i".

Applicants' methods identified clone AJ27\_1i as encoding a secreted protein.

The nucleotide sequence of the 5' portion of AJ27\_1i as presently determined is reported in SEQ ID NO:60. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AJ27\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:61. Amino acids 15 to 27 of SEQ ID NO:61 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 28. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the AJ27\_1i protein. Additional nucleotide sequence from the 3' portion of AJ27\_1i, including a poly(A) tail, is reported in SEQ ID NO:62.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AJ27\_1i should be approximately 1500 bp.

Protein "AJ142\_1i"

One protein of the present invention has been identified as protein "AJ142\_1i". A partial cDNA clone encoding AJ142\_1i was first isolated from a human adult testes cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of

this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR ,  
5 including a poly(A) tail. This full-length clone is also referred to herein as "AJ142\_1i".

Applicants' methods identified clone AJ142\_1i as encoding a secreted protein.

The nucleotide sequence of AJ142\_1i as presently determined is reported in SEQ ID NO:63, and includes the poly(A) tail. What applicants believe is the proper  
10 reading frame and the predicted amino acid sequence of the AJ142\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:64. Amino acids 11 to 23 of SEQ ID NO:64 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 24. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain  
15 should the predicted leader/signal sequence not be separated from the remainder of the AJ142\_1i protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AJ142\_1i should be approximately 650 bp.

20                   Protein "AK604\_1i"

One protein of the present invention has been identified as protein "AK604\_1i". A partial cDNA clone encoding AK604\_1i was first isolated from a human fetal kidney cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or  
25 transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor  
30 was examined and determined to be a full-length clone, including a 5' end and 3' UTR , including a poly(A) tail. This full-length clone is also referred to herein as "AK604\_1i".

Applicants' methods identified clone AK604\_1i as encoding a secreted protein.

The nucleotide sequence of the 5' portion of AK604\_1i as presently

determined is reported in SEQ ID NO:65. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AK604\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:66. Additional nucleotide sequence from the 3' portion of AK604\_1i, including a poly(A) tail, is reported in SEQ ID NO:67.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AK604\_1i should be approximately 1350 bp.

Protein "AK620\_1i"

One protein of the present invention has been identified as protein "AK620\_1i". A partial cDNA clone encoding AK620\_1i was first isolated from a human fetal kidney cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "AK620\_1i".

Applicants' methods identified clone AK620\_1i as encoding a secreted protein.

The nucleotide sequence of AK620\_1i as presently determined is reported in SEQ ID NO:68, and includes the poly(A) tail. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AK620\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:69.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AK620\_1i should be approximately 700 bp.

Protein "AK650\_1i"

One protein of the present invention has been identified as protein "AK650\_1i". A partial cDNA clone encoding AK650\_1i was first isolated from a human fetal kidney cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or

transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "AK650\_1i".

Applicants' methods identified clone AK650\_1i as encoding a secreted protein.

The nucleotide sequence of AK650\_1i as presently determined is reported in SEQ ID NO:70, and includes the poly(A) tail. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AK650\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:71. Amino acids 14 to 26 of SEQ ID NO:71 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 27. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the AK650\_1i protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AK650\_1i should be approximately 1000 bp.

#### Protein "AM226\_1i"

One protein of the present invention has been identified as protein "AM226\_1i". A partial cDNA clone encoding AM226\_1i was first isolated from a human fetal kidney cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "AM226\_1i".

Applicants' methods identified clone AM226\_1i as encoding a secreted

protein.

The nucleotide sequence of the 5' portion of AM226\_1i as presently determined is reported in SEQ ID NO:72. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AM226\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:73. Amino acids 7 to 19 of SEQ ID NO:73 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 20. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the AM226\_1i protein. Additional nucleotide sequence from the 3' portion of AM226\_1i, including a poly(A) tail, is reported in SEQ ID NO:74.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AM226\_1i should be approximately 1500 bp.

AM226\_1i protein was expressed in a Baculovirus expression system, and an expressed protein band of approximately 50 kDa detected in a conditioned medium fraction using SDS polyacrylamide gel electrophoresis.

#### Protein "AR417\_1i"

One protein of the present invention has been identified as protein "AR417\_1i". A partial cDNA clone encoding AR417\_1i was first isolated from a human adult retina cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "AR417\_1i".

Applicants' methods identified clone AR417\_1i as encoding a secreted protein.

The nucleotide sequence of AR417\_1i as presently determined is reported in SEQ ID NO:75, and includes the poly(A) tail. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AR417\_1i protein

corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:76. Amino acids 32 to 44 of SEQ ID NO:76 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 45. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the AR417\_1i protein.

The *EcoRI/NotI* restriction fragment obtainable from the deposit containing clone AR417\_1i should be approximately 1500 bp.

10                   Protein "AU43\_1i"

One protein of the present invention has been identified as protein "AU43\_1i". A partial cDNA clone encoding AU43\_1i was first isolated from a human adult testes cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "AU43\_1i".

Applicants' methods identified clone AU43\_1i as encoding a secreted protein.

The nucleotide sequence of the 5' portion of AU43\_1i as presently determined is reported in SEQ ID NO:77. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AU43\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:78. Amino acids 11 to 23 of SEQ ID NO:78 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 24. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the AU43\_1i protein. Additional nucleotide sequence from the 3' portion of AU43\_1i, including a poly(A) tail, is reported in SEQ ID NO:79.

The *EcoRI/NotI* restriction fragment obtainable from the deposit



containing clone AU43\_1i should be approximately 950 bp.

Protein "AW60\_1i"

One protein of the present invention has been identified as protein  
5 "AW60\_1i". A partial cDNA clone encoding AW60\_1i was first isolated from a human  
adult ovary (PA-1 teratocarcinoma line, pool of retinoic-acid-treated, activin-treated, and  
untreated tissue) cDNA library using methods which are selective for cDNAs encoding  
secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or  
transmembrane protein on the basis of computer analysis of the amino acid sequence of  
10 the encoded protein. A human EST matching at least part of the nucleotide sequence of  
this clone was identified by database searches. The human cDNA clone corresponding  
to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a  
distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor  
was examined and determined to be a full-length clone, including a 5' end and 3' UTR ,  
15 including a poly(A) tail. This full-length clone is also referred to herein as "AW60\_1i".

Applicants' methods identified clone AW60\_1i as encoding a secreted  
protein.

The nucleotide sequence of the 5' portion of AW60\_1i as presently  
determined is reported in SEQ ID NO:80. What applicants believe is the proper reading  
20 frame and the predicted amino acid sequence of the AW60\_1i protein corresponding to  
the foregoing nucleotide sequence is reported in SEQ ID NO:81. Amino acids 19 to 31 of  
SEQ ID NO:81 are a predicted leader/signal sequence, with the predicted mature amino  
acid sequence beginning at amino acid 32. Due to the hydrophobic nature of the predicted  
leader/signal sequence, it is likely to act as a transmembrane domain should the predicted  
25 leader/signal sequence not be separated from the remainder of the AW60\_1i protein.  
Additional nucleotide sequence from the 3' portion of AW60\_1i, including a poly(A) tail,  
is reported in SEQ ID NO:82.

The EcoRI/NotI restriction fragment obtainable from the deposit  
containing clone AW60\_1i should be approximately 1800 bp.

30

Protein "BA176\_1i"

One protein of the present invention has been identified as protein  
"BA176\_1i". A partial cDNA clone encoding BA176\_1i was first isolated from a human  
fetal placenta cDNA library using methods which are selective for cDNAs encoding

secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "BA176\_1i".

Applicants' methods identified clone BA176\_1i as encoding a secreted protein.

The nucleotide sequence of BA176\_1i as presently determined is reported in SEQ ID NO:83, and includes the poly(A) tail. What applicants believe is the proper reading frame and the predicted amino acid sequence of the BA176\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:84. Amino acids 276 to 288 of SEQ ID NO:84 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 289. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the BA176\_1i protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone BA176\_1i should be approximately 2500 bp.

#### Protein "BD140\_1i"

One protein of the present invention has been identified as protein "BD140\_1i". A partial cDNA clone encoding BD140\_1i was first isolated from a human fetal kidney cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "BD140\_1i".

Applicants' methods identified clone BD140\_1i as encoding a secreted protein.

The nucleotide sequence of BD140\_1i as presently determined is reported in SEQ ID NO:85, and includes the poly(A) tail. What applicants believe is the proper  
5 reading frame and the predicted amino acid sequence of the BD140\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:86.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone BD140\_1i should be approximately 2550 bp.

10           Protein "BD407\_1i"

One protein of the present invention has been identified as protein "BD407\_1i". A partial cDNA clone encoding BD407\_1i was first isolated from a human fetal kidney cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or  
15 transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor  
20 was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "BD407\_1i".

Applicants' methods identified clone BD407\_1i as encoding a secreted protein.

The nucleotide sequence of BD407\_1i as presently determined is reported  
25 in SEQ ID NO:87, and includes the poly(A) tail. What applicants believe is the proper reading frame and the predicted amino acid sequence of the BD407\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:88. Amino acids 2 to 14 of SEQ ID NO:88 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 15. Due to the hydrophobic nature  
30 of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the BD407\_1i protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone BD407\_1i should be approximately 1100 bp.

Protein "BF290\_1i"

One protein of the present invention has been identified as protein "BF290\_1i". A partial cDNA clone encoding BF290\_1i was first isolated from a human fetal brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "BF290\_1i".

Applicants' methods identified clone BF290\_1i as encoding a secreted protein.

The nucleotide sequence of BF290\_1i as presently determined is reported in SEQ ID NO:89, and includes the poly(A) tail. What applicants believe is the proper reading frame and the predicted amino acid sequence of the BF290\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:90.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone BF290\_1i should be approximately 1450 bp.

Protein "BG236\_1i"

One protein of the present invention has been identified as protein "BG236\_1i". A partial cDNA clone encoding BG236\_1i was first isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "BG236\_1i".

Applicants' methods identified clone BG236\_1i as encoding a secreted protein.

The nucleotide sequence of BG236\_1i as presently determined is reported in SEQ ID NO:91, and includes the poly(A) tail. What applicants believe is the proper reading frame and the predicted amino acid sequence of the BG236\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:92. Amino acids 45 to 57 of SEQ ID NO:92 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 58. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the BG236\_1i protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone BG236\_1i should be approximately 1350 bp.

Protein "BG237\_1i"

One protein of the present invention has been identified as protein "BG237\_1i". A partial cDNA clone encoding BG237\_1i was first isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "BG237\_1i".

Applicants' methods identified clone BG237\_1i as encoding a secreted protein.

The nucleotide sequence of the 5' portion of BG237\_1i as presently determined is reported in SEQ ID NO:93. What applicants believe is the proper reading frame and the predicted amino acid sequence of the BG237\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:94. Additional nucleotide sequence from the 3' portion of BG237\_1i, including a poly(A) tail, is reported in SEQ ID NO:95.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone BG237\_1i should be approximately 1300 bp.

Protein "BG255\_1i"

5 One protein of the present invention has been identified as protein "BG255\_1i". A partial cDNA clone encoding BG255\_1i was first isolated from a human adult brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of  
10 the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR ,  
15 including a poly(A) tail. This full-length clone is also referred to herein as "BG255\_1i".

Applicants' methods identified clone BG255\_1i as encoding a secreted protein.

The nucleotide sequence of the 5' portion of BG255\_1i as presently determined is reported in SEQ ID NO:96. What applicants believe is the proper reading  
20 frame and the predicted amino acid sequence of the BG255\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:97. Additional nucleotide sequence from the 3' portion of BG255\_1i, including a poly(A) tail, is reported in SEQ ID NO:98.

The EcoRI/NotI restriction fragment obtainable from the deposit  
25 containing clone BG255\_1i should be approximately 1450 bp.

Protein "H541\_3i"

One protein of the present invention has been identified as protein "H541\_3i". A partial cDNA clone encoding H541\_3i was first isolated from a human adult  
30 blood (peripheral blood mononuclear cells treated with phytohemagglutinin and phorbol myristate acetate and mixed l cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide

sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "H541\_3i".

*Applicants' methods identified clone H541\_3i as encoding a secreted protein.*

The nucleotide sequence of H541\_3i as presently determined is reported in SEQ ID NO:99, and includes the poly(A) tail. What applicants believe is the proper reading frame and the predicted amino acid sequence of the H541\_3i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:100. Amino acids 4 to 16 of SEQ ID NO:100 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 17. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the H541\_3i protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone H541\_3i should be approximately 1500 bp.

H541\_3i protein was expressed in a COS cell expression system, and an expressed protein band of approximately 41 kDa detected in a membrane fraction using SDS polyacrylamide gel electrophoresis.

#### Protein "H978\_1i"

One protein of the present invention has been identified as protein "H978\_1i". A partial cDNA clone encoding H978\_1i was first isolated from a human adult blood (peripheral blood mononuclear cells treated with phytohemagglutinin and phorbol myristate acetate and mixed l cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from

the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "H978\_1i".

Applicants' methods identified clone H978\_1i as encoding a secreted  
5 protein.

The nucleotide sequence of the 5' portion of H978\_1i as presently determined is reported in SEQ ID NO:101. What applicants believe is the proper reading frame and the predicted amino acid sequence of the H978\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:102. Additional nucleotide  
10 sequence from the 3' portion of H978\_1i, including a poly(A) tail, is reported in SEQ ID NO:103.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone H978\_1i should be approximately 1100 bp.

15 Protein "L161\_1i"

One protein of the present invention has been identified as protein "L161\_1i". A partial cDNA clone encoding L161\_1i was first isolated from a mouse adult thymus cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or  
20 transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor  
25 was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "L161\_1i".

Applicants' methods identified clone L161\_1i as encoding a secreted protein.

The nucleotide sequence of the 5' portion of L161\_1i as presently  
30 determined is reported in SEQ ID NO:104. What applicants believe is the proper reading frame and the predicted amino acid sequence of the L161\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:105. Additional nucleotide sequence from the 3' portion of L161\_1i, including a poly(A) tail, is reported in SEQ ID NO:106.



The EcoRI/NotI restriction fragment obtainable from the deposit containing clone L161\_1i should be approximately 1300 bp.

Protein "AE648\_1i"

5 One protein of the present invention has been identified as protein "AE648\_1i". A partial cDNA clone encoding AE648\_1i was first isolated from a mouse adult spleen (stimulated with concanavalin A and mixed with dendritic cells) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on  
10 the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and  
15 determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "AE648\_1i".

Applicants' methods identified clone AE648\_1i as encoding a secreted protein.

The nucleotide sequence of AE648\_1i as presently determined is reported  
20 in SEQ ID NO:107, and includes the poly(A) tail. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AE648\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:108. Amino acids 3 to 15 of SEQ ID NO:108 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 16. Due to the  
25 hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the AE648\_1i protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AE648\_1i should be approximately 900 bp.

30

Protein "AE693\_1i"

One protein of the present invention has been identified as protein "AE693\_1i". A partial cDNA clone encoding AE693\_1i was first isolated from a mouse adult spleen (stimulated with concanavalin A and mixed with dendritic cells) cDNA

library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified  
5 by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "AE693\_1i".

10 Applicants' methods identified clone AE693\_1i as encoding a secreted protein.

The nucleotide sequence of the 5' portion of AE693\_1i as presently determined is reported in SEQ ID NO:109. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AE693\_1i protein corresponding to  
15 the foregoing nucleotide sequence is reported in SEQ ID NO:110. Additional nucleotide sequence from the 3' portion of AE693\_1i, including a poly(A) tail, is reported in SEQ ID NO:111.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AE693\_1i should be approximately 1200 bp.

20

Protein "AK438\_1i"

One protein of the present invention has been identified as protein "AK438\_1i". A partial cDNA clone encoding AK438\_1i was first isolated from a human fetal kidney cDNA library using methods which are selective for cDNAs encoding  
25 secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a  
30 distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "AK438\_1i".

Applicants' methods identified clone AK438\_1i as encoding a secreted protein.

The nucleotide sequence of the 5' portion of AK438\_1i as presently determined is reported in SEQ ID NO:112. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AK438\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:113. Additional nucleotide  
5 sequence from the 3' portion of AK438\_1i, including a poly(A) tail, is reported in SEQ ID NO:114.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AK438\_1i should be approximately 1000 bp.

10           Protein "AK609\_1i"

One protein of the present invention has been identified as protein "AK609\_1i". A partial cDNA clone encoding AK609\_1i was first isolated from a human fetal kidney cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or  
15 transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor  
20 was examined and determined to be a full-length clone, including a 5' end and 3' UTR. This full-length clone is also referred to herein as "AK609\_1i".

Applicants' methods identified clone AK609\_1i as encoding a secreted protein.

The nucleotide sequence of the 5' portion of AK609\_1i as presently  
25 determined is reported in SEQ ID NO:115. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AK609\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:116. Additional nucleotide sequence from the 3' portion of AK609\_1i is reported in SEQ ID NO:117.

The EcoRI/NotI restriction fragment obtainable from the deposit  
30 containing clone AK609\_1i should be approximately 750 bp.

Protein "AM1060\_1i"

One protein of the present invention has been identified as protein "AM1060\_1i". A partial cDNA clone encoding AM1060\_1i was first isolated from a human

fetal kidney cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "AM1060\_1i".

10 Applicants' methods identified clone AM1060\_1i as encoding a secreted protein.

The nucleotide sequence of the 5' portion of AM1060\_1i as presently determined is reported in SEQ ID NO:118. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AM1060\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:119. Amino acids 13 to 25 of SEQ ID NO:119 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 26. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the AM1060\_1i protein. Additional nucleotide sequence from the 3' portion of AM1060\_1i, including a poly(A) tail, is reported in SEQ ID NO:120.

15  
20

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AM1060\_1i should be approximately 1700 bp.

25 Protein "AQ2\_1i"

One protein of the present invention has been identified as protein "AQ2\_1i". A partial cDNA clone encoding AQ2\_1i was first isolated from a human adult ovary (PA-1 teratocarcinoma line, pool of retinoic-acid-treated, activin-treated, and untreated tissue) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a

30

distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "AQ2\_1i".

Applicants' methods identified clone AQ2\_1i as encoding a secreted  
5 protein.

The nucleotide sequence of the 5' portion of AQ2\_1i as presently determined is reported in SEQ ID NO:121. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AQ2\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:122. Additional nucleotide  
10 sequence from the 3' portion of AQ2\_1i, including a poly(A) tail, is reported in SEQ ID NO:123.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AQ2\_1i should be approximately 1370 bp.

15 Protein "K433\_1i"

One protein of the present invention has been identified as protein "K433\_1i". A partial cDNA clone encoding K433\_1i was first isolated from a mouse adult bone marrow (stromal line FCM-4) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as  
20 encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from  
25 the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "K433\_1i".

Applicants' methods identified clone K433\_1i as encoding a secreted protein.

30 The nucleotide sequence of the 5' portion of K433\_1i as presently determined is reported in SEQ ID NO:124. What applicants believe is the proper reading frame and the predicted amino acid sequence of the K433\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:125. Additional nucleotide sequence from the 3' portion of K433\_1i, including a poly(A) tail, is reported in SEQ ID

NO:126.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone K433\_1i should be approximately 1200 bp.

5                   Protein "L256\_1i"

One protein of the present invention has been identified as protein "L256\_1i". A partial cDNA clone encoding L256\_1i was first isolated from a mouse adult thymus cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or  
10 transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor  
15 was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "L256\_1i".

Applicants' methods identified clone L256\_1i as encoding a secreted protein.

The nucleotide sequence of the 5' portion of L256\_1i as presently  
20 determined is reported in SEQ ID NO:127. What applicants believe is the proper reading frame and the predicted amino acid sequence of the L256\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:128. Additional nucleotide sequence from the 3' portion of L256\_1i, including a poly(A) tail, is reported in SEQ ID NO:129.

25                   The EcoRI/NotI restriction fragment obtainable from the deposit containing clone L256\_1i should be approximately 1400 bp.

Protein "AM207\_1i"

One protein of the present invention has been identified as protein  
30 "AM207\_1i". A partial cDNA clone encoding AM207\_1i was first isolated from a human fetal kidney cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of

this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "AM207\_1i".

Applicants' methods identified clone AM207\_1i as encoding a secreted protein.

The nucleotide sequence of the 5' portion of AM207\_1i as presently determined is reported in SEQ ID NO:130. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AM207\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:131. Additional nucleotide sequence from the 3' portion of AM207\_1i, including a poly(A) tail, is reported in SEQ ID NO:132.

Protein "AM910\_1i"

One protein of the present invention has been identified as protein "AM910\_1i". A partial cDNA clone encoding AM910\_1i was first isolated from a human fetal kidney cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "AM910\_1i".

Applicants' methods identified clone AM910\_1i as encoding a secreted protein.

The nucleotide sequence of AM910\_1i as presently determined is reported in SEQ ID NO:133, and includes the poly(A) tail. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AM910\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:134. Amino acids 7 to 19 of SEQ ID NO:134 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 20. Due to the

hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the AM910\_1i protein.

The EcoRI/NotI restriction fragment obtainable from the deposit  
5 containing clone AM910\_1i should be approximately 1200 bp.

#### Protein "AR54\_1i"

One protein of the present invention has been identified as protein "AR54\_1i". A partial cDNA clone encoding AR54\_1i was first isolated from a human adult  
10 retina cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding  
15 to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "AR54\_1i".

Applicants' methods identified clone AR54\_1i as encoding a secreted  
20 protein.

The nucleotide sequence of AR54\_1i as presently determined is reported in SEQ ID NO:135, and includes the poly(A) tail. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AR54\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:136.  
25 Amino acids 8 to 20 of SEQ ID NO:136 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 21. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the AR54\_1i protein.

The EcoRI/NotI restriction fragment obtainable from the deposit  
30 containing clone AR54\_1i should be approximately 1300 bp.

#### Protein "L200\_1i"

One protein of the present invention has been identified as protein



"L200\_1i". A partial cDNA clone encoding L200\_1i was first isolated from a mouse adult thymus cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "L200\_1i".

Applicants' methods identified clone L200\_1i as encoding a secreted protein.

The nucleotide sequence of L200\_1i as presently determined is reported in SEQ ID NO:137, and includes the poly(A) tail. What applicants believe is the proper reading frame and the predicted amino acid sequence of the L200\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:138. Amino acids 272 to 284 of SEQ ID NO:138 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 285. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the L200\_1i protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone L200\_1i should be approximately 1330 bp.

#### 25      Protein "WA129\_2i"

One protein of the present invention has been identified as protein "WA129\_2i". A partial cDNA clone encoding WA129\_2i was first isolated from a Xenopus embryo (dorsal mesoderm) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from

the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "WA129\_2i".

Applicants' methods identified clone WA129\_2i as encoding a secreted  
5 protein.

The nucleotide sequence of WA129\_2i as presently determined is reported in SEQ ID NO:139, and includes the poly(A) tail. What applicants believe is the proper reading frame and the predicted amino acid sequence of the WA129\_2i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:140.  
10 Amino acids 10 to 22 of SEQ ID NO:140 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 23. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the WA129\_2i protein.

15 The EcoRI/NotI restriction fragment obtainable from the deposit containing clone WA129\_2i should be approximately 1933 bp.

WA192\_2i protein was expressed in a Baculovirus expression system, and an expressed protein band of approximately 39 kDa detected in a conditioned medium fraction using SDS polyacrylamide gel electrophoresis.

20

#### Protein "WA154\_3i"

One protein of the present invention has been identified as protein "WA154\_3i". A partial cDNA clone encoding WA154\_3i was first isolated from a Xenopus embryo (dorsal mesoderm) cDNA library using methods which are selective for cDNAs  
25 encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St.  
30 Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "WA154\_3i".

Applicants' methods identified clone WA154\_3i as encoding a secreted

protein.

The nucleotide sequence of WA154\_3i as presently determined is reported in SEQ ID NO:141, and includes the poly(A) tail. What applicants believe is the proper reading frame and the predicted amino acid sequence of the WA154\_3i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:142. Amino acids 11 to 23 of SEQ ID NO:142 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 24. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the WA154\_3i protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone WA154\_3i should be approximately 1469 bp.

WA154\_3i protein was expressed in a COS cell expression system, and an expressed protein band of approximately 17 kDa detected in conditioned medium and membrane fractions using SDS polyacrylamide gel electrophoresis.

#### Protein "AA36\_1i"

One protein of the present invention has been identified as protein "AA36\_1i". A partial cDNA clone encoding AA36\_1i was first isolated from a human fetal kidney cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "AA36\_1i".

Applicants' methods identified clone AA36\_1i as encoding a secreted protein.

The nucleotide sequence of AA36\_1i as presently determined is reported in SEQ ID NO:143, and includes the poly(A) tail. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AA36\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:144.

Amino acids 3 to 15 of SEQ ID NO:144 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 16. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated  
5 from the remainder of the AA36\_1i protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AA36\_1i should be approximately 1450 bp.

#### Protein "AC175\_2i"

10 One protein of the present invention has been identified as protein "AC175\_2i". A partial cDNA clone encoding AC175\_2i was first isolated from a human fetal placenta cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of  
15 the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR,  
20 including a poly(A) tail. This full-length clone is also referred to herein as "AC175\_2i".

Applicants' methods identified clone AC175\_2i as encoding a secreted protein.

The nucleotide sequence of AC175\_2i as presently determined is reported in SEQ ID NO:145, and includes the poly(A) tail. What applicants believe is the proper  
25 reading frame and the predicted amino acid sequence of the AC175\_2i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:146.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone AC175\_2i should be approximately 842 bp.

#### Protein "AV189\_1i"

30 One protein of the present invention has been identified as protein "AV189\_1i". A partial cDNA clone encoding AV189\_1i was first isolated from a mouse adult spleen (concanavalin A stimulated and mixed with dendritic cells) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat.

No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was  
5 ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "AV189\_1i".

Applicants' methods identified clone AV189\_1i as encoding a secreted  
10 protein.

The nucleotide sequence of AV189\_1i as presently determined is reported in SEQ ID NO:147, and includes the poly(A) tail. What applicants believe is the proper reading frame and the predicted amino acid sequence of the AV189\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:148.  
15 Amino acids 72 to 84 of SEQ ID NO:148 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 85. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the AV189\_1i protein.

20

#### Protein "K368\_1i"

One protein of the present invention has been identified as protein "K368\_1i". A partial cDNA clone encoding K368\_1i was first isolated from a mouse adult bone marrow (stromal cell line FCM-4) cDNA library using methods which are selective  
25 for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc.,  
30 St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "K368\_1i".

Applicants' methods identified clone K368\_1i as encoding a secreted

protein.

The nucleotide sequence of K368\_1i as presently determined is reported in SEQ ID NO:149, and includes the poly(A) tail. What applicants believe is the proper reading frame and the predicted amino acid sequence of the K368\_1i protein  
5 corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:150. Amino acids 88 to 100 of SEQ ID NO:150 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 101. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated  
10 from the remainder of the K368\_1i protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone K368\_1i should be approximately 983 bp.

K368\_1i protein was expressed in a Baculovirus expression system, and an expressed protein band of approximately 28 kDa detected in conditioned medium and  
15 membrane fractions using SDS polyacrylamide gel electrophoresis.

#### Protein "K568\_1i"

One protein of the present invention has been identified as protein "K568\_1i". A partial cDNA clone encoding K568\_1i was first isolated from a mouse adult  
20 bone marrow (stromal cell line FCM-4) cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA  
25 clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "K568\_1i".

30 Applicants' methods identified clone K568\_1i as encoding a secreted protein.

The nucleotide sequence of K568\_1i as presently determined is reported in SEQ ID NO:151, and includes the poly(A) tail. What applicants believe is the proper reading frame and the predicted amino acid sequence of the K568\_1i protein

corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:152. Amino acids 57 to 69 of SEQ ID NO:152 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 70. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the K568\_1i protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone K568\_1i should be approximately 1254 bp.

#### 10           Protein "T85\_1i"

One protein of the present invention has been identified as protein "T85\_1i". A partial cDNA clone encoding T85\_1i was first isolated from a mouse fetal brain cDNA library using methods which are selective for cDNAs encoding secreted proteins (see U.S. Pat. No. 5,536,637), or was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. A human EST matching at least part of the nucleotide sequence of this clone was identified by database searches. The human cDNA clone corresponding to the EST database entry was ordered from Genome Systems, Inc., St. Louis, Mo, a distributor of the I.M.A.G.E. Consortium library. The clone received from the distributor was examined and determined to be a full-length clone, including a 5' end and 3' UTR, including a poly(A) tail. This full-length clone is also referred to herein as "T85\_1i".

Applicants' methods identified clone T85\_1i as encoding a secreted protein.

The nucleotide sequence of T85\_1i as presently determined is reported in SEQ ID NO:153, and includes the poly(A) tail. What applicants believe is the proper reading frame and the predicted amino acid sequence of the T85\_1i protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:154.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone T85\_1i should be approximately 1803 bp.

30

#### Deposit of Clones

Clones AK296\_1i, AK533\_1i, AK583\_1i, AM282\_1i, AM340\_1i, AM610\_1i, AP162\_1i, AR260\_1i, AS32\_1i, AS34\_1i, AT205\_1i, AT211\_1i, AT319\_1i, AW191\_1i, BB9\_1i, H617\_1i, K39\_1i and K640\_1i were deposited on April 17, 1996 with the American

Type Culture Collection (10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were given the accession number ATCC 98026, from which each clone comprising a particular polynucleotide is obtainable.

5 Clones AE402\_1i, AE610\_1i, AH106\_1i, AH196\_1i, AI6\_1i, AJ13\_1i, AJ27\_1i, AJ142\_1i, AK604\_1i, AK620\_1i, AK650\_1i, AM226\_1i, AR417\_1i, AU43\_1i, AW60\_1i, BA176\_1i, BD140\_1i, BD407\_1i and BF290\_1i were deposited on October 2, 1996 with the American Type Culture Collection (10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under the Budapest Treaty and were  
10 given the accession number ATCC 98190, from which each clone comprising a particular polynucleotide is obtainable.

Clones BG236\_1i, BG237\_1i, BG255\_1i, H541\_3i, H978\_1i and L161\_1i were deposited on October 2, 1996 with the American Type Culture Collection (10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under  
15 the Budapest Treaty and were given the accession number ATCC 98191, from which each clone comprising a particular polynucleotide is obtainable.

Clones AE648\_1i, AE693\_1i, AK438\_1i, AK609\_1i, AM1060\_1i, AQ2\_1i, K433\_1i and L256\_1i were deposited on October 31, 1996 with the American Type Culture Collection (10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an  
20 original deposit under the Budapest Treaty and were given the accession number ATCC 98237, from which each clone comprising a particular polynucleotide is obtainable.

Clones AM207\_1i, AM910\_1i, AR54\_1i, L200\_1i, WA129\_2i and WA154\_3i were deposited on August 21, 1997 with the American Type Culture Collection (10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under  
25 the Budapest Treaty and were given the accession number ATCC 98510, from which each clone comprising a particular polynucleotide is obtainable.

Clones AA36\_1i, AC175\_2i, AV189\_1i, K368\_1i, K568\_1i and T85\_1i were deposited on December 18, 1998 with the American Type Culture Collection (10801 University Boulevard, Manassas, Virginia 20110-2209 U.S.A.) as an original deposit under  
30 the Budapest Treaty and were given the accession number ATCC XXXXX, from which each clone comprising a particular polynucleotide is obtainable.

All restrictions on the availability to the public of the deposited material will be irrevocably removed upon the granting of the patent, except for the requirements specified in 37 C.F.R. § 1.808(b), and the term of the deposit will comply with 37 C.F.R.



## § 1.806.

Each clone has been transfected into separate bacterial cells (*E. coli*) in this composite deposit. Each clone can be removed from the vector in which it was deposited by performing an EcoRI/NotI digestion (5' site, EcoRI; 3' site, NotI) to produce the appropriate fragment for such clone. Each clone was deposited in either the pED6 or pNOTs vector depicted in Figures 1A and 1B, respectively. The pED6dpc2 vector ("pED6") was derived from pED6dpc1 by insertion of a new polylinker to facilitate cDNA cloning (Kaufman *et al.*, 1991, *Nucleic Acids Res.* 19: 4485-4490); the pNOTs vector was derived from pMT2 (Kaufman *et al.*, 1989, *Mol. Cell. Biol.* 9: 946-958) by deletion of the DHFR sequences, insertion of a new polylinker, and insertion of the M13 origin of replication in the ClaI site. In some instances, the deposited clone can become "flipped" (i.e., in the reverse orientation) in the deposited isolate. In such instances, the cDNA insert can still be isolated by digestion with EcoRI and NotI. However, NotI will then produce the 5' site and EcoRI will produce the 3' site for placement of the cDNA in proper orientation for expression in a suitable vector. The cDNA may also be expressed from the vectors in which they were deposited.

Bacterial cells containing a particular clone can be obtained from the composite deposit as follows:

An oligonucleotide probe or probes should be designed to the sequence that is known for that particular clone. This sequence can be derived from the sequences provided herein, or from a combination of those sequences.

In the probe sequences derived from the sequences provided, position 2 is occupied in preferred probes/primers by a biotinylated phosphoramidite residue rather than a nucleotide (such as, for example, that produced by use of biotin phosphoramidite (1-dimethoxytrityloxy-2-(N-biotinyl-4-aminobutyl)-propyl-3-O-(2-cyanoethyl)-(N,N-diisopropyl)-phosphoramidite) (Glen Research, cat. no. 10-1953)).

The design of the oligonucleotide probe should preferably follow these parameters:

- (a) It should be designed to an area of the sequence which has the fewest ambiguous bases ("N's"), if any;
- (b) It should be designed to have a  $T_m$  of approx. 80 ° C (assuming 2° for each A or T and 4 degrees for each G or C).

The oligonucleotide should preferably be labeled with  $^{-32}\text{P}$  ATP (specific activity 6000 Ci/mmol) and T4 polynucleotide kinase using commonly employed techniques for labeling oligonucleotides. Other labeling techniques can also be used. Unincorporated label should preferably be removed by gel filtration chromatography or other established methods. The amount of radioactivity incorporated into the probe should be quantitated by measurement in a scintillation counter. Preferably, specific activity of the resulting probe should be approximately  $4 \times 10^6$  dpm/pmol.

The bacterial culture containing the pool of full-length clones should preferably be thawed and 100 l of the stock used to inoculate a sterile culture flask containing 25 ml of sterile L-broth containing ampicillin at 100 g/ml. The culture should preferably be grown to saturation at  $37^\circ\text{C}$ , and the saturated culture should preferably be diluted in fresh L-broth. Aliquots of these dilutions should preferably be plated to determine the dilution and volume which will yield approximately 5000 distinct and well-separated colonies on solid bacteriological media containing L-broth containing ampicillin at 100 g/ml and agar at 1.5% in a 150 mm petri dish when grown overnight at  $37^\circ\text{C}$ . Other known methods of obtaining distinct, well-separated colonies can also be employed.

Standard colony hybridization procedures should then be used to transfer the colonies to nitrocellulose filters and lyse, denature and bake them.

The filter is then preferably incubated at  $65^\circ\text{C}$  for 1 hour with gentle agitation in 6X SSC (20X stock is 175.3 g NaCl/liter, 88.2 g Na citrate/liter, adjusted to pH 7.0 with NaOH) containing 0.5% SDS, 100 g/ml of yeast RNA, and 10 mM EDTA (approximately 10 mL per 150 mm filter). Preferably, the probe is then added to the hybridization mix at a concentration greater than or equal to  $1 \times 10^6$  dpm/mL. The filter is then preferably incubated at  $65^\circ\text{C}$  with gentle agitation overnight. The filter is then preferably washed in 500 mL of 2X SSC/0.5% SDS at room temperature without agitation, preferably followed by 500 mL of 2X SSC/0.1% SDS at room temperature with gentle shaking for 15 minutes. A third wash with 0.1X SSC/0.5% SDS at  $65^\circ\text{C}$  for 30 minutes to 1 hour is optional. The filter is then preferably dried and subjected to autoradiography for sufficient time to visualize the positives on the X-ray film. Other known hybridization methods can also be employed.

The positive colonies are picked, grown in culture, and plasmid DNA isolated using standard procedures. The clones can then be verified by restriction analysis, hybridization analysis, or DNA sequencing.

Fragments of the proteins of the present invention which are capable of exhibiting biological activity are also encompassed by the present invention. Fragments of the protein may be in linear form or they may be cyclized using known methods, for example, as described in H.U. Saragovi, *et al.*, *Bio/Technology* 10, 773-778 (1992) and in  
5 R.S. McDowell, *et al.*, *J. Amer. Chem. Soc.* 114, 9245-9253 (1992), both of which are incorporated herein by reference. Such fragments may be fused to carrier molecules such as immunoglobulins for many purposes, including increasing the valency of protein binding sites. For example, fragments of the protein may be fused through "linker" sequences to the Fc portion of an immunoglobulin. For a bivalent form of the protein,  
10 such a fusion could be to the Fc portion of an IgG molecule. Other immunoglobulin isotypes may also be used to generate such fusions. For example, a protein - IgM fusion would generate a decavalent form of the protein of the invention.

The present invention also provides both full-length and mature forms of the disclosed proteins. The full-length form of the such proteins is identified in the  
15 sequence listing by translation of the nucleotide sequence of each disclosed clone. The mature form(s) of such protein may be obtained by expression of the disclosed full-length polynucleotide (preferably those deposited with ATCC) in a suitable mammalian cell or other host cell. The sequence(s) of the mature form(s) of the protein may also be determinable from the amino acid sequence of the full-length form.

20 Where the protein of the present invention is membrane-bound (e.g., is a receptor), the present invention also provides for soluble forms of such protein. In such forms, part or all of the intracellular and transmembrane domains of the protein are deleted such that the protein is fully secreted from the cell in which it is expressed. The intracellular and transmembrane domains of proteins of the invention can be identified  
25 in accordance with known techniques for determination of such domains from sequence information. For example, the TopPredII computer program can be used to predict the location of transmembrane domains in an amino acid sequence, domains which are described by the location of the center of the transmembrane domain, with at least ten transmembrane amino acids on each side of the reported central residue(s).

30 Proteins and protein fragments of the present invention include proteins with amino acid sequence lengths that are at least 25% (more preferably at least 50%, and most preferably at least 75%) of the length of a disclosed protein and have at least 60% sequence identity (more preferably, at least 75% identity; most preferably at least 90% or 95% identity) with that disclosed protein, where sequence identity is determined by

comparing the amino acid sequences of the proteins when aligned so as to maximize overlap and identity while minimizing sequence gaps. Also included in the present invention are proteins and protein fragments that contain a segment preferably comprising 8 or more (more preferably 20 or more, most preferably 30 or more) contiguous amino acids that shares at least 75% sequence identity (more preferably, at least 85% identity; most preferably at least 95% identity) with any such segment of any of the disclosed proteins.

In particular, sequence identity may be determined using WU-BLAST (Washington University BLAST) version 2.0 software, which builds upon WU-BLAST version 1.4, which in turn is based on the public domain NCBI-BLAST version 1.4 (Altschul and Gish, 1996, Local alignment statistics, Doolittle *ed.*, *Methods in Enzymology* 266: 460-480; Altschul *et al.*, 1990, Basic local alignment search tool, *Journal of Molecular Biology* 215: 403-410; Gish and States, 1993, Identification of protein coding regions by database similarity search, *Nature Genetics* 3: 266-272; Karlin and Altschul, 1993, Applications and statistics for multiple high-scoring segments in molecular sequences, *Proc. Natl. Acad. Sci. USA* 90: 5873-5877; all of which are incorporated by reference herein). WU-BLAST version 2.0 executable programs for several UNIX platforms can be downloaded from <ftp://blast.wustl.edu/blast/executables>. The complete suite of search programs (BLASTP, BLASTN, BLASTX, TBLASTN, and TBLASTX) is provided at that site, in addition to several support programs. WU-BLAST 2.0 is copyrighted and may not be sold or redistributed in any form or manner without the express written consent of the author; but the posted executables may otherwise be freely used for commercial, nonprofit, or academic purposes. In all search programs in the suite -- BLASTP, BLASTN, BLASTX, TBLASTN and TBLASTX -- the gapped alignment routines are integral to the database search itself, and thus yield much better sensitivity and selectivity while producing the more easily interpreted output. Gapping can optionally be turned off in all of these programs, if desired. The default penalty (Q) for a gap of length one is Q=9 for proteins and BLASTP, and Q=10 for BLASTN, but may be changed to any integer value including zero, one through eight, nine, ten, eleven, twelve through twenty, twenty-one through fifty, fifty-one through one hundred, etc. The default per-residue penalty for extending a gap (R) is R=2 for proteins and BLASTP, and R=10 for BLASTN, but may be changed to any integer value including zero, one, two, three, four, five, six,

seven, eight, nine, ten, eleven, twelve through twenty, twenty-one through fifty, fifty-one through one hundred, etc. Any combination of values for Q and R can be used in order to align sequences so as to maximize overlap and identity while minimizing sequence gaps. The default amino acid comparison matrix is BLOSUM62, but other amino acid  
5 comparison matrices such as PAM can be utilized.

Species homologues of the disclosed proteins are also provided by the present invention. As used herein, a "species homologue" is a protein or polynucleotide with a different species of origin from that of a given protein or polynucleotide, but with significant sequence similarity to the  
10 given protein or polynucleotide. Preferably, polynucleotide species homologues have at least 60% sequence identity (more preferably, at least 75% identity; most preferably at least 90% identity) with the given polynucleotide, and protein species homologues have at least 30% sequence identity (more preferably, at least 45% identity; most preferably at least 60% identity) with the given protein, where sequence  
15 identity is determined by comparing the nucleotide sequences of the polynucleotides or the amino acid sequences of the proteins when aligned so as to maximize overlap and identity while minimizing sequence gaps. Species homologues may be isolated and identified by making suitable probes or primers from the sequences provided herein and screening a suitable nucleic acid source from the desired species. Preferably, species  
20 homologues are those isolated from mammalian species. Most preferably, species homologues are those isolated from certain mammalian species such as, for example, *Pan troglodytes*, *Gorilla gorilla*, *Pongo pygmaeus*, *Hylobates concolor*, *Macaca mulatta*, *Papio papio*, *Papio hamadryas*, *Cercopithecus aethiops*, *Cebus capucinus*, *Aotus trivirgatus*, *Sanguinus oedipus*, *Microcebus murinus*, *Mus musculus*, *Rattus norvegicus*, *Cricetulus griseus*, *Felis catus*, *Mustela*  
25 *vison*, *Canis familiaris*, *Oryctolagus cuniculus*, *Bos taurus*, *Ovis aries*, *Sus scrofa*, and *Equus caballus*, for which genetic maps have been created allowing the identification of syntenic relationships between the genomic organization of genes in one species and the genomic organization of the related genes in another species (O'Brien and Seuánez, 1988, *Ann. Rev. Genet.* 22: 323-351; O'Brien *et al.*, 1993, *Nature Genetics* 3:103-112; Johansson *et al.*, 1995, *Genomics* 25: 682-690; Lyons *et al.*, 1997, *Nature Genetics* 15: 47-56; O'Brien *et al.*, 1997, *Trends in Genetics* 13(10): 393-399; Carver and Stubbs, 1997, *Genome Research* 7:1123-1137; all of which are incorporated by reference herein).

The invention also encompasses allelic variants of the disclosed proteins;

that is, naturally-occurring alternative forms of the isolated proteins which are identical or have significantly similar sequences to those encoded by the disclosed polynucleotides. Preferably, allelic variants have at least 60% sequence identity (more preferably, at least 75% identity; most preferably at least 90% identity) with the given polynucleotide.

5 where sequence identity is determined by comparing the nucleotide sequences of the polynucleotides when aligned so as to maximize overlap and identity while minimizing sequence gaps. Allelic variants may be isolated and identified by making suitable probes or primers from the sequences provided herein and screening a suitable nucleic acid source from individuals of the appropriate species.

10 The invention also includes polynucleotides with sequences complementary to those of the polynucleotides disclosed herein.

The isolated polynucleotide encoding the protein of the invention may be operably linked to an expression control sequence such as the pMT2 or pED expression vectors disclosed in Kaufman *et al.*, Nucleic Acids Res. 19, 4485-4490 (1991), in order to  
15 produce the protein recombinantly. Many suitable expression control sequences are known in the art. General methods of expressing recombinant proteins are also known and are exemplified in R. Kaufman, Methods in Enzymology 185, 537-566 (1990). As defined herein "operably linked" means that the isolated polynucleotide of the invention and an expression control sequence are situated within a vector or cell in such a way that  
20 the protein is expressed by a host cell which has been transformed (transfected) with the ligated polynucleotide/expression control sequence.

A number of types of cells may act as suitable host cells for expression of the protein. Mammalian host cells include, for example, monkey COS cells, Chinese Hamster Ovary (CHO) cells, human kidney 293 cells, human epidermal A431 cells, human  
25 Colo205 cells, 3T3 cells, CV-1 cells, other transformed primate cell lines, normal diploid cells, cell strains derived from in vitro culture of primary tissue, primary explants, HeLa cells, mouse L cells, BHK, HL-60, U937, HaK or Jurkat cells.

Alternatively, it may be possible to produce the protein in lower eukaryotes such as yeast or in prokaryotes such as bacteria. Potentially suitable yeast strains include  
30 *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Kluyveromyces* strains, *Candida*, or any yeast strain capable of expressing heterologous proteins. Potentially suitable bacterial strains include *Escherichia coli*, *Bacillus subtilis*, *Salmonella typhimurium*, or any bacterial strain capable of expressing heterologous proteins. If the protein is made in yeast or

bacteria, it may be necessary to modify the protein produced therein, for example by phosphorylation or glycosylation of the appropriate sites, in order to obtain the functional protein. Such covalent attachments may be accomplished using known chemical or enzymatic methods.

5           The protein may also be produced by operably linking the isolated polynucleotide of the invention to suitable control sequences in one or more insect expression vectors, and employing an insect expression system. Materials and methods for baculovirus/insect cell expression systems are commercially available in kit form from, *e.g.*, Invitrogen, San Diego, California, U.S.A. (the MaxBac® kit), and such methods are  
10 well known in the art, as described in Summers and Smith, Texas Agricultural Experiment Station Bulletin No. 1555 (1987), incorporated herein by reference. As used herein, an insect cell capable of expressing a polynucleotide of the present invention is "transformed."

          The protein of the invention may be prepared by culturing transformed  
15 host cells under culture conditions suitable to express the recombinant protein. The resulting expressed protein may then be purified from such culture (*i.e.*, from culture medium or cell extracts) using known purification processes, such as gel filtration and ion exchange chromatography. The purification of the protein may also include an affinity column containing agents which will bind to the protein; one or more column steps over  
20 such affinity resins as concanavalin A-agarose, heparin-toyopearl® or Cibacrom blue 3GA Sepharose®; one or more steps involving hydrophobic interaction chromatography using such resins as phenyl ether, butyl ether, or propyl ether; or immunoaffinity chromatography.

          Alternatively, the protein of the invention may also be expressed in a form  
25 which will facilitate purification. For example, it may be expressed as a fusion protein, such as those of maltose binding protein (MBP), glutathione-S-transferase (GST) or thioredoxin (TRX). Kits for expression and purification of such fusion proteins are commercially available from New England BioLabs (Beverly, MA), Pharmacia (Piscataway, NJ) and Invitrogen Corporation (Carlsbad, CA), respectively. The protein  
30 can also be tagged with an epitope and subsequently purified by using a specific antibody directed to such epitope. One such epitope ("Flag") is commercially available from the Eastman Kodak Company (New Haven, CT).

          Finally, one or more reverse-phase high performance liquid chromatography (RP-HPLC) steps employing hydrophobic RP-HPLC media, *e.g.*, silica

gel having pendant methyl or other aliphatic groups, can be employed to further purify the protein. Some or all of the foregoing purification steps, in various combinations, can also be employed to provide a substantially homogeneous isolated recombinant protein. The protein thus purified is substantially free of other mammalian proteins and is defined  
5 in accordance with the present invention as an "isolated protein."

The protein of the invention may also be expressed as a product of transgenic animals, e.g., as a component of the milk of transgenic cows, goats, pigs, or sheep which are characterized by somatic or germ cells containing a nucleotide sequence encoding the protein.

10 The protein may also be produced by known conventional chemical synthesis. Methods for constructing the proteins of the present invention by synthetic means are known to those skilled in the art. The synthetically-constructed protein sequences, by virtue of sharing primary, secondary or tertiary structural and/or conformational characteristics with proteins may possess biological properties in common  
15 therewith, including protein activity. Thus, they may be employed as biologically active or immunological substitutes for natural, purified proteins in screening of therapeutic compounds and in immunological processes for the development of antibodies.

The proteins provided herein also include proteins characterized by amino acid sequences similar to those of purified proteins but into which modification are  
20 naturally provided or deliberately engineered. For example, modifications in the peptide or DNA sequences can be made by those skilled in the art using known techniques. Modifications of interest in the protein sequences may include the alteration, substitution, replacement, insertion or deletion of a selected amino acid residue in the coding sequence. For example, one or more of the cysteine residues may be deleted or replaced with  
25 another amino acid to alter the conformation of the molecule. Techniques for such alteration, substitution, replacement, insertion or deletion are well known to those skilled in the art (see, e.g., U.S. Patent No. 4,518,584). Preferably, such alteration, substitution, replacement, insertion or deletion retains the desired activity of the protein.

Other fragments and derivatives of the sequences of proteins which would  
30 be expected to retain protein activity in whole or in part and may thus be useful for screening or other immunological methodologies may also be easily made by those skilled in the art given the disclosures herein. Such modifications are believed to be encompassed by the present invention.



### USES AND BIOLOGICAL ACTIVITY

The proteins of the present invention are expected to exhibit one or more of the uses or biological activities (including those associated with assays cited herein) identified below. Uses or activities described for proteins of the present invention may be provided by administration or use of such proteins or by administration or use of polynucleotides encoding such proteins (such as, for example, in gene therapies or vectors suitable for introduction of DNA).

#### Research Uses and Utilities

The proteins provided by the present invention can similarly be used in assay to determine biological activity, including in a panel of multiple proteins for high-throughput screening; to raise antibodies or to elicit another immune response; as a reagent (including the labeled reagent) in assays designed to quantitatively determine levels of the protein (or its receptor) in biological fluids; as markers for tissues in which the corresponding protein is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in a disease state); and, of course, to isolate correlative receptors or ligands. Where the protein binds or potentially binds to another protein (such as, for example, in a receptor-ligand interaction), the protein can be used to identify the other protein with which binding occurs or to identify inhibitors of the binding interaction. Proteins involved in these binding interactions can also be used to screen for peptide or small molecule inhibitors or agonists of the binding interaction.

Any or all of these research utilities are capable of being developed into reagent grade or kit format for commercialization as research products.

Methods for performing the uses listed above are well known to those skilled in the art. References disclosing such methods include without limitation "Molecular Cloning: A Laboratory Manual", 2d ed., Cold Spring Harbor Laboratory Press, Sambrook, J., E.F. Fritsch and T. Maniatis eds., 1989, and "Methods in Enzymology: Guide to Molecular Cloning Techniques", Academic Press, Berger, S.L. and A.R. Kimmel eds., 1987.

#### Nutritional Uses

Proteins of the present invention can also be used as nutritional sources or supplements. Such uses include without limitation use as a protein or amino acid

supplement, use as a carbon source, use as a nitrogen source and use as a source of carbohydrate. In such cases the protein of the invention can be added to the feed of a particular organism or can be administered as a separate solid or liquid preparation, such as in the form of powder, pills, solutions, suspensions or capsules. In the case of  
5 microorganisms, the protein of the invention can be added to the medium in or on which the microorganism is cultured.

Cytokine and Cell Proliferation/Differentiation Activity

A protein of the present invention may exhibit cytokine, cell proliferation (either inducing or inhibiting) or cell differentiation (either inducing or inhibiting) activity  
10 or may induce production of other cytokines in certain cell populations. Many protein factors discovered to date, including all known cytokines, have exhibited activity in one or more factor-dependent cell proliferation assays, and hence the assays serve as a convenient confirmation of cytokine activity. The activity of a protein of the present invention is evidenced by any one of a number of routine factor dependent cell  
15 proliferation assays for cell lines including, without limitation, 32D, DA2, DA1G, T10, B9, B9/11, BaF3, MC9/G, M+ (preB M+), 2E8, RB5, DA1, 123, T1165, HT2, CTLL2, TF-1, Mo7e and CMK.

The activity of a protein of the invention may, among other means, be measured by the following methods:

20 Assays for T-cell or thymocyte proliferation include without limitation those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Takai et al., J. Immunol. 137:3494-3500, 1986;  
25 Bertagnolli et al., J. Immunol. 145:1706-1712, 1990; Bertagnolli et al., Cellular Immunology 133:327-341, 1991; Bertagnolli, et al., J. Immunol. 149:3778-3783, 1992; Bowman et al., J. Immunol. 152: 1756-1761, 1994.

Assays for cytokine production and/or proliferation of spleen cells, lymph node cells or thymocytes include, without limitation, those described in: Polyclonal T cell  
30 stimulation, Kruisbeek, A.M. and Shevach, E.M. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 3.12.1-3.12.14, John Wiley and Sons, Toronto. 1994; and Measurement of mouse and human Interferon, Schreiber, R.D. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.8.1-6.8.8, John Wiley and Sons, Toronto. 1994.

Assays for proliferation and differentiation of hematopoietic and

- lymphopoietic cells include, without limitation, those described in: Measurement of Human and Murine Interleukin 2 and Interleukin 4, Bottomly, K., Davis, L.S. and Lipsky, P.E. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.3.1-6.3.12, John Wiley and Sons, Toronto. 1991; deVries et al., J. Exp. Med. 173:1205-1211, 1991; Moreau et al., Nature 336:690-692, 1988; Greenberger et al., Proc. Natl. Acad. Sci. U.S.A. 80:2931-2938, 1983; Measurement of mouse and human interleukin 6 - Nordan, R. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.6.1-6.6.5, John Wiley and Sons, Toronto. 1991; Smith et al., Proc. Natl. Acad. Sci. U.S.A. 83:1857-1861, 1986; Measurement of human Interleukin 11 - Bennett, F., Giannotti, J., Clark, S.C. and Turner, K. J. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.15.1 John Wiley and Sons, Toronto. 1991; Measurement of mouse and human Interleukin 9 - Ciarletta, A., Giannotti, J., Clark, S.C. and Turner, K.J. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.13.1, John Wiley and Sons. Toronto. 1991.

- Assays for T-cell clone responses to antigens (which will identify, among others, proteins that affect APC-T cell interactions as well as direct T-cell effects by measuring proliferation and cytokine production) include, without limitation, those described in: *Current Protocols in Immunology*, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function; Chapter 6, Cytokines and their cellular receptors; Chapter 7, Immunologic studies in Humans); Weinberger et al., Proc. Natl. Acad. Sci. USA 77:6091-6095, 1980; Weinberger et al., Eur. J. Immun. 11:405-411, 1981; Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988.

#### 25 Immune Stimulating or Suppressing Activity

- A protein of the present invention may also exhibit immune stimulating or immune suppressing activity, including without limitation the activities for which assays are described herein. A protein may be useful in the treatment of various immune deficiencies and disorders (including severe combined immunodeficiency (SCID)), e.g., in regulating (up or down) growth and proliferation of T and/or B lymphocytes, as well as effecting the cytolytic activity of NK cells and other cell populations. These immune deficiencies may be genetic or be caused by viral (e.g., HIV) as well as bacterial or fungal infections, or may result from autoimmune disorders. More specifically, infectious diseases caused by viral, bacterial, fungal or other infection may be treatable using a

protein of the present invention, including infections by HIV, hepatitis viruses, herpesviruses, mycobacteria, Leishmania spp., malaria spp. and various fungal infections such as candidiasis. Of course, in this regard, a protein of the present invention may also be useful where a boost to the immune system generally may be desirable, *i.e.*, in the  
5 treatment of cancer.

Autoimmune disorders which may be treated using a protein of the present invention include, for example, connective tissue disease, multiple sclerosis, systemic lupus erythematosus, rheumatoid arthritis, autoimmune pulmonary inflammation, Guillain-Barre syndrome, autoimmune thyroiditis, insulin dependent diabetes mellitis,  
10 myasthenia gravis, graft-versus-host disease and autoimmune inflammatory eye disease. Such a protein of the present invention may also to be useful in the treatment of allergic reactions and conditions, such as asthma (particularly allergic asthma) or other respiratory problems. Other conditions, in which immune suppression is desired (including, for example, organ transplantation), may also be treatable using a protein of the present  
15 invention.

Using the proteins of the invention it may also be possible to regulate immune responses in a number of ways. Down regulation may be in the form of inhibiting or blocking an immune response already in progress or may involve preventing the induction of an immune response. The functions of activated T cells may be inhibited  
20 by suppressing T cell responses or by inducing specific tolerance in T cells, or both. Immunosuppression of T cell responses is generally an active, non-antigen-specific, process which requires continuous exposure of the T cells to the suppressive agent. Tolerance, which involves inducing non-responsiveness or anergy in T cells, is distinguishable from immunosuppression in that it is generally antigen-specific and  
25 persists after exposure to the tolerizing agent has ceased. Operationally, tolerance can be demonstrated by the lack of a T cell response upon reexposure to specific antigen in the absence of the tolerizing agent.

Down regulating or preventing one or more antigen functions (including without limitation B lymphocyte antigen functions (such as , for example, B7)), *e.g.*,  
30 preventing high level lymphokine synthesis by activated T cells, will be useful in situations of tissue, skin and organ transplantation and in graft-versus-host disease (GVHD). For example, blockage of T cell function should result in reduced tissue destruction in tissue transplantation. Typically, in tissue transplants, rejection of the transplant is initiated through its recognition as foreign by T cells, followed by an immune

reaction that destroys the transplant. The administration of a molecule which inhibits or blocks interaction of a B7 lymphocyte antigen with its natural ligand(s) on immune cells (such as a soluble, monomeric form of a peptide having B7-2 activity alone or in conjunction with a monomeric form of a peptide having an activity of another B lymphocyte antigen (*e.g.*, B7-1, B7-3) or blocking antibody), prior to transplantation can lead to the binding of the molecule to the natural ligand(s) on the immune cells without transmitting the corresponding costimulatory signal. Blocking B lymphocyte antigen function in this matter prevents cytokine synthesis by immune cells, such as T cells, and thus acts as an immunosuppressant. Moreover, the lack of costimulation may also be sufficient to anergize the T cells, thereby inducing tolerance in a subject. Induction of long-term tolerance by B lymphocyte antigen-blocking reagents may avoid the necessity of repeated administration of these blocking reagents. To achieve sufficient immunosuppression or tolerance in a subject, it may also be necessary to block the function of a combination of B lymphocyte antigens.

The efficacy of particular blocking reagents in preventing organ transplant rejection or GVHD can be assessed using animal models that are predictive of efficacy in humans. Examples of appropriate systems which can be used include allogeneic cardiac grafts in rats and xenogeneic pancreatic islet cell grafts in mice, both of which have been used to examine the immunosuppressive effects of CTLA4Ig fusion proteins *in vivo* as described in Lenschow *et al.*, *Science* 257:789-792 (1992) and Turka *et al.*, *Proc. Natl. Acad. Sci USA*, 89:11102-11105 (1992). In addition, murine models of GVHD (see Paul ed., *Fundamental Immunology*, Raven Press, New York, 1989, pp. 846-847) can be used to determine the effect of blocking B lymphocyte antigen function *in vivo* on the development of that disease.

Blocking antigen function may also be therapeutically useful for treating autoimmune diseases. Many autoimmune disorders are the result of inappropriate activation of T cells that are reactive against self tissue and which promote the production of cytokines and autoantibodies involved in the pathology of the diseases. Preventing the activation of autoreactive T cells may reduce or eliminate disease symptoms. Administration of reagents which block costimulation of T cells by disrupting receptor:ligand interactions of B lymphocyte antigens can be used to inhibit T cell activation and prevent production of autoantibodies or T cell-derived cytokines which may be involved in the disease process. Additionally, blocking reagents may induce antigen-specific tolerance of autoreactive T cells which could lead to long-term relief from

the disease. The efficacy of blocking reagents in preventing or alleviating autoimmune disorders can be determined using a number of well-characterized animal models of human autoimmune diseases. Examples include murine experimental autoimmune encephalitis, systemic lupus erythmatosis in MRL/*lpr/lpr* mice or NZB hybrid mice, 5 murine autoimmune collagen arthritis, diabetes mellitus in NOD mice and BB rats, and murine experimental myasthenia gravis (see Paul ed., Fundamental Immunology, Raven Press, New York, 1989, pp. 840-856).

Upregulation of an antigen function (preferably a B lymphocyte antigen function), as a means of up regulating immune responses, may also be useful in therapy.

- 10 Upregulation of immune responses may be in the form of enhancing an existing immune response or eliciting an initial immune response. For example, enhancing an immune response through stimulating B lymphocyte antigen function may be useful in cases of viral infection. In addition, systemic viral diseases such as influenza, the common cold, and encephalitis might be alleviated by the administration of stimulatory forms of B 15 lymphocyte antigens systemically.

- Alternatively, anti-viral immune responses may be enhanced in an infected patient by removing T cells from the patient, costimulating the T cells *in vitro* with viral antigen- pulsed APCs either expressing a peptide of the present invention or together with a stimulatory form of a soluble peptide of the present invention and reintroducing 20 the *in vitro* activated T cells into the patient. Another method of enhancing anti-viral immune responses would be to isolate infected cells from a patient, transfect them with a nucleic acid encoding a protein of the present invention as described herein such that the cells express all or a portion of the protein on their surface, and reintroduce the transfected cells into the patient. The infected cells would now be capable of delivering 25 a costimulatory signal to, and thereby activate, T cells *in vivo*.

- In another application, up regulation or enhancement of antigen function (preferably B lymphocyte antigen function) may be useful in the induction of tumor immunity. Tumor cells (*e.g.*, sarcoma, melanoma, lymphoma, leukemia, neuroblastoma, carcinoma) transfected with a nucleic acid encoding at least one peptide of the present 30 invention can be administered to a subject to overcome tumor-specific tolerance in the subject. If desired, the tumor cell can be transfected to express a combination of peptides. For example, tumor cells obtained from a patient can be transfected *ex vivo* with an expression vector directing the expression of a peptide having B7-2-like activity alone, or in conjunction with a peptide having B7-1-like activity and/or B7-3-like activity. The

transfected tumor cells are returned to the patient to result in expression of the peptides on the surface of the transfected cell. Alternatively, gene therapy techniques can be used to target a tumor cell for transfection *in vivo*.

The presence of the peptide of the present invention having the activity of a B lymphocyte antigen(s) on the surface of the tumor cell provides the necessary costimulation signal to T cells to induce a T cell mediated immune response against the transfected tumor cells. In addition, tumor cells which lack MHC class I or MHC class II molecules, or which fail to reexpress sufficient amounts of MHC class I or MHC class II molecules, can be transfected with nucleic acid encoding all or a portion of (e.g., a cytoplasmic-domain truncated portion) of an MHC class I chain protein and a microglobulin protein or an MHC class II chain protein and an MHC class II chain protein to thereby express MHC class I or MHC class II proteins on the cell surface. Expression of the appropriate class I or class II MHC in conjunction with a peptide having the activity of a B lymphocyte antigen (e.g., B7-1, B7-2, B7-3) induces a T cell mediated immune response against the transfected tumor cell. Optionally, a gene encoding an antisense construct which blocks expression of an MHC class II associated protein, such as the invariant chain, can also be cotransfected with a DNA encoding a peptide having the activity of a B lymphocyte antigen to promote presentation of tumor associated antigens and induce tumor specific immunity. Thus, the induction of a T cell mediated immune response in a human subject may be sufficient to overcome tumor-specific tolerance in the subject.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for thymocyte or splenocyte cytotoxicity include, without limitation, those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Herrmann et al., Proc. Natl. Acad. Sci. USA 78:2488-2492, 1981; Herrmann et al., J. Immunol. 128:1968-1974, 1982; Handa et al., J. Immunol. 135:1564-1572, 1985; Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988; Herrmann et al., Proc. Natl. Acad. Sci. USA 78:2488-2492, 1981; Herrmann et al., J. Immunol. 128:1968-1974, 1982; Handa et al., J. Immunol. 135:1564-1572, 1985; Takai et al., J. Immunol. 137:3494-3500, 1986; Bowman et al., J. Virology 61:1992-1998; Takai et al., J. Immunol. 140:508-512, 1988; Bertagnolli et al.,

Cellular Immunology 133:327-341, 1991; Brown et al., J. Immunol. 153:3079-3092, 1994.

Assays for T-cell-dependent immunoglobulin responses and isotype switching (which will identify, among others, proteins that modulate T-cell dependent antibody responses and that affect Th1/Th2 profiles) include, without limitation, those  
5 described in: Maliszewski, J. Immunol. 144:3028-3033, 1990; and Assays for B cell function: *In vitro* antibody production, Mond, J.J. and Brunswick, M. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 3.8.1-3.8.16, John Wiley and Sons, Toronto. 1994.

Mixed lymphocyte reaction (MLR) assays (which will identify, among  
10 others, proteins that generate predominantly Th1 and CTL responses) include, without limitation, those described in: *Current Protocols in Immunology*, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, *In Vitro* assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, *Immunologic studies in Humans*); Takai et al., J. Immunol.  
15 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988; Bertagnoli et al., J. Immunol. 149:3778-3783, 1992.

Dendritic cell-dependent assays (which will identify, among others, proteins expressed by dendritic cells that activate naive T-cells) include, without limitation, those described in: Guery et al., J. Immunol. 134:536-544, 1995; Inaba et al.,  
20 *Journal of Experimental Medicine* 173:549-559, 1991; Macatonia et al., *Journal of Immunology* 154:5071-5079, 1995; Porgador et al., *Journal of Experimental Medicine* 182:255-260, 1995; Nair et al., *Journal of Virology* 67:4062-4069, 1993; Huang et al., *Science* 264:961-965, 1994; Macatonia et al., *Journal of Experimental Medicine* 169:1255-1264, 1989; Bhardwaj et al., *Journal of Clinical Investigation* 94:797-807, 1994; and Inaba et al., *Journal*  
25 *of Experimental Medicine* 172:631-640, 1990.

Assays for lymphocyte survival/apoptosis (which will identify, among others, proteins that prevent apoptosis after superantigen induction and proteins that regulate lymphocyte homeostasis) include, without limitation, those described in: Darzynkiewicz et al., *Cytometry* 13:795-808, 1992; Gorczyca et al., *Leukemia* 7:659-670,  
30 1993; Gorczyca et al., *Cancer Research* 53:1945-1951, 1993; Itoh et al., *Cell* 66:233-243, 1991; Zacharchuk, *Journal of Immunology* 145:4037-4045, 1990; Zamai et al., *Cytometry* 14:891-897, 1993; Gorczyca et al., *International Journal of Oncology* 1:639-648, 1992.

Assays for proteins that influence early steps of T-cell commitment and development include, without limitation, those described in: Antica et al., *Blood*



84:111-117, 1994; Fine et al., Cellular Immunology 155:111-122, 1994; Galy et al., Blood 85:2770-2778, 1995; Toki et al., Proc. Nat. Acad. Sci. USA 88:7548-7551, 1991.

#### Hematopoiesis Regulating Activity

5 A protein of the present invention may be useful in regulation of hematopoiesis and, consequently, in the treatment of myeloid or lymphoid cell deficiencies. Even marginal biological activity in support of colony forming cells or of factor-dependent cell lines indicates involvement in regulating hematopoiesis, e.g. in supporting the growth and proliferation of erythroid progenitor cells alone or in  
10 combination with other cytokines, thereby indicating utility, for example, in treating various anemias or for use in conjunction with irradiation/chemotherapy to stimulate the production of erythroid precursors and/or erythroid cells; in supporting the growth and proliferation of myeloid cells such as granulocytes and monocytes/macrophages (i.e., traditional CSF activity) useful, for example, in conjunction with chemotherapy to prevent  
15 or treat consequent myelo-suppression; in supporting the growth and proliferation of megakaryocytes and consequently of platelets thereby allowing prevention or treatment of various platelet disorders such as thrombocytopenia, and generally for use in place of or complimentary to platelet transfusions; and/or in supporting the growth and proliferation of hematopoietic stem cells which are capable of maturing to any and all of  
20 the above-mentioned hematopoietic cells and therefore find therapeutic utility in various stem cell disorders (such as those usually treated with transplantation, including, without limitation, aplastic anemia and paroxysmal nocturnal hemoglobinuria), as well as in repopulating the stem cell compartment post irradiation/chemotherapy, either *in-vivo* or *ex-vivo* (i.e., in conjunction with bone marrow transplantation or with peripheral  
25 progenitor cell transplantation (homologous or heterologous)) as normal cells or genetically manipulated for gene therapy.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for proliferation and differentiation of various  
30 hematopoietic lines are cited above.

Assays for embryonic stem cell differentiation (which will identify, among others, proteins that influence embryonic differentiation hematopoiesis) include, without limitation, those described in: Johansson et al. Cellular Biology 15:141-151, 1995; Keller et al., Molecular and Cellular Biology 13:473-486, 1993; McClanahan et al., Blood

81:2903-2915, 1993.

- Assays for stem cell survival and differentiation (which will identify, among others, proteins that regulate lympho-hematopoiesis) include, without limitation, those described in: Methylcellulose colony forming assays, Freshney, M.G. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 265-268, Wiley-Liss, Inc., New York, NY. 1994; Hirayama et al., Proc. Natl. Acad. Sci. USA 89:5907-5911, 1992; Primitive hematopoietic colony forming cells with high proliferative potential, McNiece, I.K. and Briddell, R.A. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 23-39, Wiley-Liss, Inc., New York, NY. 1994; Neben et al., Experimental Hematology 22:353-359, 1994; Cobblestone area forming cell assay, Ploemacher, R.E. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 1-21, Wiley-Liss, Inc., New York, NY. 1994; Long term bone marrow cultures in the presence of stromal cells, Spooncer, E., Dexter, M. and Allen, T. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 163-179, Wiley-Liss, Inc., New York, NY. 1994; Long term culture initiating cell assay, Sutherland, H.J. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 139-162, Wiley-Liss, Inc., New York, NY. 1994.

#### Tissue Growth Activity

- A protein of the present invention also may have utility in compositions used for bone, cartilage, tendon, ligament and/or nerve tissue growth or regeneration, as well as for wound healing and tissue repair and replacement, and in the treatment of burns, incisions and ulcers.

- A protein of the present invention, which induces cartilage and/or bone growth in circumstances where bone is not normally formed, has application in the healing of bone fractures and cartilage damage or defects in humans and other animals. Such a preparation employing a protein of the invention may have prophylactic use in closed as well as open fracture reduction and also in the improved fixation of artificial joints. *De novo* bone formation induced by an osteogenic agent contributes to the repair of congenital, trauma induced, or oncologic resection induced craniofacial defects, and also is useful in cosmetic plastic surgery.

A protein of this invention may also be used in the treatment of periodontal disease, and in other tooth repair processes. Such agents may provide an environment to attract bone-forming cells, stimulate growth of bone-forming cells or induce differentiation of progenitors of bone-forming cells. A protein of the invention may also

be useful in the treatment of osteoporosis or osteoarthritis, such as through stimulation of bone and/or cartilage repair or by blocking inflammation or processes of tissue destruction (collagenase activity, osteoclast activity, etc.) mediated by inflammatory processes.

5 Another category of tissue regeneration activity that may be attributable to the protein of the present invention is tendon/ligament formation. A protein of the present invention, which induces tendon/ligament-like tissue or other tissue formation in circumstances where such tissue is not normally formed, has application in the healing of tendon or ligament tears, deformities and other tendon or ligament defects in humans  
10 and other animals. Such a preparation employing a tendon/ligament-like tissue inducing protein may have prophylactic use in preventing damage to tendon or ligament tissue, as well as use in the improved fixation of tendon or ligament to bone or other tissues, and in repairing defects to tendon or ligament tissue. De novo tendon/ligament-like tissue formation induced by a composition of the present invention contributes to the repair of  
15 congenital, trauma induced, or other tendon or ligament defects of other origin, and is also useful in cosmetic plastic surgery for attachment or repair of tendons or ligaments. The compositions of the present invention may provide an environment to attract tendon- or ligament-forming cells, stimulate growth of tendon- or ligament-forming cells, induce differentiation of progenitors of tendon- or ligament-forming cells, or induce growth of  
20 tendon/ligament cells or progenitors *ex vivo* for return *in vivo* to effect tissue repair. The compositions of the invention may also be useful in the treatment of tendinitis, carpal tunnel syndrome and other tendon or ligament defects. The compositions may also include an appropriate matrix and/or sequestering agent as a carrier as is well known in the art.

25 The protein of the present invention may also be useful for proliferation of neural cells and for regeneration of nerve and brain tissue, *i.e.* for the treatment of central and peripheral nervous system diseases and neuropathies, as well as mechanical and traumatic disorders, which involve degeneration, death or trauma to neural cells or nerve tissue. More specifically, a protein may be used in the treatment of diseases of the  
30 peripheral nervous system, such as peripheral nerve injuries, peripheral neuropathy and localized neuropathies, and central nervous system diseases, such as Alzheimer's, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, and Shy-Drager syndrome. Further conditions which may be treated in accordance with the present invention include mechanical and traumatic disorders, such as spinal cord disorders, head

trauma and cerebrovascular diseases such as stroke. Peripheral neuropathies resulting from chemotherapy or other medical therapies may also be treatable using a protein of the invention.

Proteins of the invention may also be useful to promote better or faster  
5 closure of non-healing wounds, including without limitation pressure ulcers, ulcers associated with vascular insufficiency, surgical and traumatic wounds, and the like.

It is expected that a protein of the present invention may also exhibit activity for generation or regeneration of other tissues, such as organs (including, for example, pancreas, liver, intestine, kidney, skin, endothelium), muscle (smooth, skeletal  
10 or cardiac) and vascular (including vascular endothelium) tissue, or for promoting the growth of cells comprising such tissues. Part of the desired effects may be by inhibition or modulation of fibrotic scarring to allow normal tissue to regenerate. A protein of the invention may also exhibit angiogenic activity.

A protein of the present invention may also be useful for gut protection or  
15 regeneration and treatment of lung or liver fibrosis, reperfusion injury in various tissues, and conditions resulting from systemic cytokine damage.

A protein of the present invention may also be useful for promoting or inhibiting differentiation of tissues described above from precursor tissues or cells; or for inhibiting the growth of tissues described above.

20 The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for tissue generation activity include, without limitation, those described in: International Patent Publication No. WO95/16035 (bone, cartilage, tendon); International Patent Publication No. WO95/05846 (nerve, neuronal); International Patent  
25 Publication No. WO91/07491 (skin, endothelium ).

Assays for wound healing activity include, without limitation, those described in: Winter, Epidermal Wound Healing, pps. 71-112 (Maibach, HI and Rovee, DT, eds.), Year Book Medical Publishers, Inc., Chicago, as modified by Eaglstein and Mertz, J. Invest. Dermatol 71:382-84 (1978).

30

#### Activin/Inhibin Activity

A protein of the present invention may also exhibit activin- or inhibin-related activities. Inhibins are characterized by their ability to inhibit the release of follicle stimulating hormone (FSH), while activins and are characterized by their ability

to stimulate the release of follicle stimulating hormone (FSH). Thus, a protein of the present invention, alone or in heterodimers with a member of the inhibin family, may be useful as a contraceptive based on the ability of inhibins to decrease fertility in female mammals and decrease spermatogenesis in male mammals. Administration of sufficient amounts of other inhibins can induce infertility in these mammals. Alternatively, the protein of the invention, as a homodimer or as a heterodimer with other protein subunits of the inhibin- group, may be useful as a fertility inducing therapeutic, based upon the ability of activin molecules in stimulating FSH release from cells of the anterior pituitary. See, for example, United States Patent 4,798,885. A protein of the invention may also be useful for advancement of the onset of fertility in sexually immature mammals, so as to increase the lifetime reproductive performance of domestic animals such as cows, sheep and pigs.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for activin/inhibin activity include, without limitation, those described in: Vale et al., *Endocrinology* 91:562-572, 1972; Ling et al., *Nature* 321:779-782, 1986; Vale et al., *Nature* 321:776-779, 1986; Mason et al., *Nature* 318:659-663, 1985; Forage et al., *Proc. Natl. Acad. Sci. USA* 83:3091-3095, 1986.

#### Chemotactic/Chemokinetic Activity

A protein of the present invention may have chemotactic or chemokinetic activity (e.g., act as a chemokine) for mammalian cells, including, for example, monocytes, fibroblasts, neutrophils, T-cells, mast cells, eosinophils, epithelial and/or endothelial cells. Chemotactic and chemokinetic proteins can be used to mobilize or attract a desired cell population to a desired site of action. Chemotactic or chemokinetic proteins provide particular advantages in treatment of wounds and other trauma to tissues, as well as in treatment of localized infections. For example, attraction of lymphocytes, monocytes or neutrophils to tumors or sites of infection may result in improved immune responses against the tumor or infecting agent.

A protein or peptide has chemotactic activity for a particular cell population if it can stimulate, directly or indirectly, the directed orientation or movement of such cell population. Preferably, the protein or peptide has the ability to directly stimulate directed movement of cells. Whether a particular protein has chemotactic activity for a population of cells can be readily determined by employing such protein or

peptide in any known assay for cell chemotaxis.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for chemotactic activity (which will identify proteins that induce  
5 or prevent chemotaxis) consist of assays that measure the ability of a protein to induce the  
migration of cells across a membrane as well as the ability of a protein to induce the  
adhesion of one cell population to another cell population. Suitable assays for movement  
and adhesion include, without limitation, those described in: Current Protocols in  
Immunology, Ed by J.E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach,  
10 W.Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 6.12,  
Measurement of alpha and beta Chemokines 6.12.1-6.12.28; Taub et al. J. Clin. Invest.  
95:1370-1376, 1995; Lind et al. APMIS 103:140-146, 1995; Muller et al Eur. J. Immunol. 25:  
1744-1748; Gruber et al. J. of Immunol. 152:5860-5867, 1994; Johnston et al. J. of Immunol.  
153: 1762-1768, 1994.

15

#### Hemostatic and Thrombolytic Activity

A protein of the invention may also exhibit hemostatic or thrombolytic  
activity. As a result, such a protein is expected to be useful in treatment of various  
coagulation disorders (including hereditary disorders, such as hemophilias) or to enhance  
20 coagulation and other hemostatic events in treating wounds resulting from trauma,  
surgery or other causes. A protein of the invention may also be useful for dissolving or  
inhibiting formation of thromboses and for treatment and prevention of conditions  
resulting therefrom (such as, for example, infarction of cardiac and central nervous system  
vessels (e.g., stroke).

25 The activity of a protein of the invention may, among other means, be  
measured by the following methods:

Assay for hemostatic and thrombolytic activity include, without limitation,  
those described in: Linet et al., J. Clin. Pharmacol. 26:131-140, 1986; Burdick et al.,  
Thrombosis Res. 45:413-419, 1987; Humphrey et al., Fibrinolysis 5:71-79 (1991); Schaub,  
30 Prostaglandins 35:467-474, 1988.

#### Receptor/Ligand Activity

A protein of the present invention may also demonstrate activity as  
receptors, receptor ligands or inhibitors or agonists of receptor/ligand interactions.

Examples of such receptors and ligands include, without limitation, cytokine receptors and their ligands, receptor kinases and their ligands, receptor phosphatases and their ligands, receptors involved in cell-cell interactions and their ligands (including without limitation, cellular adhesion molecules (such as selectins, integrins and their ligands) and  
5 receptor/ligand pairs involved in antigen presentation, antigen recognition and development of cellular and humoral immune responses). Receptors and ligands are also useful for screening of potential peptide or small molecule inhibitors of the relevant receptor/ligand interaction. A protein of the present invention (including, without limitation, fragments of receptors and ligands) may themselves be useful as inhibitors of  
10 receptor/ligand interactions.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for receptor-ligand activity include without limitation those described in: Current Protocols in Immunology, Ed by J.E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W. Strober, Pub. Greene Publishing Associates and  
15 Wiley-Interscience (Chapter 7.28, Measurement of Cellular Adhesion under static conditions 7.28.1-7.28.22), Takai et al., Proc. Natl. Acad. Sci. USA 84:6864-6868, 1987; Bierer et al., J. Exp. Med. 168:1145-1156, 1988; Rosenstein et al., J. Exp. Med. 169:149-160 1989; Stoltenberg et al., J. Immunol. Methods 175:59-68, 1994; Stitt et al., Cell 80:661-670,  
20 1995.

#### Anti-Inflammatory Activity

Proteins of the present invention may also exhibit anti-inflammatory activity. The anti-inflammatory activity may be achieved by providing a stimulus to cells  
25 involved in the inflammatory response, by inhibiting or promoting cell-cell interactions (such as, for example, cell adhesion), by inhibiting or promoting chemotaxis of cells involved in the inflammatory process, inhibiting or promoting cell extravasation, or by stimulating or suppressing production of other factors which more directly inhibit or promote an inflammatory response. Proteins exhibiting such activities can be used to treat  
30 inflammatory conditions including chronic or acute conditions), including without limitation inflammation associated with infection (such as septic shock, sepsis or systemic inflammatory response syndrome (SIRS)), ischemia-reperfusion injury, endotoxin lethality, arthritis, complement-mediated hyperacute rejection, nephritis, cytokine or chemokine-induced lung injury, inflammatory bowel disease, Crohn's disease or resulting

from over production of cytokines such as TNF or IL-1. Proteins of the invention may also be useful to treat anaphylaxis and hypersensitivity to an antigenic substance or material.

#### Cadherin/Tumor Invasion Suppressor Activity

5 Cadherins are calcium-dependent adhesion molecules that appear to play major roles during development, particularly in defining specific cell types. Loss or alteration of normal cadherin expression can lead to changes in cell adhesion properties linked to tumor growth and metastasis. Cadherin malfunction is also implicated in other human diseases, such as pemphigus vulgaris and pemphigus foliaceus (auto-immune  
10 blistering skin diseases), Crohn's disease, and some developmental abnormalities.

The cadherin superfamily includes well over forty members, each with a distinct pattern of expression. All members of the superfamily have in common conserved extracellular repeats (cadherin domains), but structural differences are found in other parts of the molecule. The cadherin domains bind calcium to form their tertiary  
15 structure and thus calcium is required to mediate their adhesion. Only a few amino acids in the first cadherin domain provide the basis for homophilic adhesion; modification of this recognition site can change the specificity of a cadherin so that instead of recognizing only itself, the mutant molecule can now also bind to a different cadherin. In addition, some cadherins engage in heterophilic adhesion with other cadherins.

20 E-cadherin, one member of the cadherin superfamily, is expressed in epithelial cell types. Pathologically, if E-cadherin expression is lost in a tumor, the malignant cells become invasive and the cancer metastasizes. Transfection of cancer cell lines with polynucleotides expressing E-cadherin has reversed cancer-associated changes by returning altered cell shapes to normal, restoring cells' adhesiveness to each other and  
25 to their substrate, decreasing the cell growth rate, and drastically reducing anchorage-independent cell growth. Thus, reintroducing E-cadherin expression reverts carcinomas to a less advanced stage. It is likely that other cadherins have the same invasion suppressor role in carcinomas derived from other tissue types. Therefore, proteins of the present invention with cadherin activity, and polynucleotides of the  
30 present invention encoding such proteins, can be used to treat cancer. Introducing such proteins or polynucleotides into cancer cells can reduce or eliminate the cancerous changes observed in these cells by providing normal cadherin expression.

Cancer cells have also been shown to express cadherins of a different tissue type than their origin, thus allowing these cells to invade and metastasize in a different



tissue in the body. Proteins of the present invention with cadherin activity, and polynucleotides of the present invention encoding such proteins, can be substituted in these cells for the inappropriately expressed cadherins, restoring normal cell adhesive properties and reducing or eliminating the tendency of the cells to metastasize.

5           Additionally, proteins of the present invention with cadherin activity, and polynucleotides of the present invention encoding such proteins, can be used to generate antibodies recognizing and binding to cadherins. Such antibodies can be used to block the adhesion of inappropriately expressed tumor-cell cadherins, preventing the cells from forming a tumor elsewhere. Such an anti-cadherin antibody can also be used as a marker  
10 for the grade, pathological type, and prognosis of a cancer, i.e. the more progressed the cancer, the less cadherin expression there will be, and this decrease in cadherin expression can be detected by the use of a cadherin-binding antibody.

          Fragments of proteins of the present invention with cadherin activity, preferably a polypeptide comprising a decapeptide of the cadherin recognition site, and  
15 poly-nucleotides of the present invention encoding such protein fragments, can also be used to block cadherin function by binding to cadherins and preventing them from binding in ways that produce undesirable effects. Additionally, fragments of proteins of the present invention with cadherin activity, preferably truncated soluble cadherin fragments which have been found to be stable in the circulation of cancer patients, and  
20 polynucleotides encoding such protein fragments, can be used to disturb proper cell-cell adhesion.

          Assays for cadherin adhesive and invasive suppressor activity include, without limitation, those described in: Hortsch et al. J Biol Chem 270 (32): 18809-18817, 1995; Miyaki et al. Oncogene 11: 2547-2552, 1995; Ozawa et al. Cell 63: 1033-1038, 1990.

25

#### Tumor Inhibition Activity

          In addition to the activities described above for immunological treatment or prevention of tumors, a protein of the invention may exhibit other anti-tumor activities. A protein may inhibit tumor growth directly or indirectly (such as, for example, via  
30 antibody-dependent cell-mediated cytotoxicity (ADCC)). A protein may exhibit its tumor inhibitory activity by acting on tumor tissue or tumor precursor tissue, by inhibiting formation of tissues necessary to support tumor growth (such as, for example, by inhibiting angiogenesis), by causing production of other factors, agents or cell types which inhibit tumor growth, or by suppressing, eliminating or inhibiting factors, agents or cell

types which promote tumor growth.

#### Other Activities

A protein of the invention may also exhibit one or more of the following additional activities or effects: inhibiting the growth, infection or function of, or killing, infectious agents, including, without limitation, bacteria, viruses, fungi and other parasites; effecting (suppressing or enhancing) bodily characteristics, including, without limitation, height, weight, hair color, eye color, skin, fat to lean ratio or other tissue pigmentation, or organ or body part size or shape (such as, for example, breast augmentation or diminution, change in bone form or shape); effecting biorhythms or circadian cycles or rhythms; effecting the fertility of male or female subjects; effecting the metabolism, catabolism, anabolism, processing, utilization, storage or elimination of dietary fat, lipid, protein, carbohydrate, vitamins, minerals, cofactors or other nutritional factors or component(s); effecting behavioral characteristics, including, without limitation, appetite, libido, stress, cognition (including cognitive disorders), depression (including depressive disorders) and violent behaviors; providing analgesic effects or other pain reducing effects; promoting differentiation and growth of embryonic stem cells in lineages other than hematopoietic lineages; hormonal or endocrine activity; in the case of enzymes, correcting deficiencies of the enzyme and treating deficiency-related diseases; treatment of hyperproliferative disorders (such as, for example, psoriasis); immunoglobulin-like activity (such as, for example, the ability to bind antigens or complement); and the ability to act as an antigen in a vaccine composition to raise an immune response against such protein or another material or entity which is cross-reactive with such protein.

25

#### ADMINISTRATION AND DOSING

A protein of the present invention (from whatever source derived, including without limitation from recombinant and non-recombinant sources) may be used in a pharmaceutical composition when combined with a pharmaceutically acceptable carrier. Such a composition may also contain (in addition to protein and a carrier) diluents, fillers, salts, buffers, stabilizers, solubilizers, and other materials well known in the art. The term "pharmaceutically acceptable" means a non-toxic material that does not interfere with the effectiveness of the biological activity of the active ingredient(s). The characteristics of the carrier will depend on the route of administration. The

pharmaceutical composition of the invention may also contain cytokines, lymphokines, or other hematopoietic factors such as M-CSF, GM-CSF, TNF, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IFN, TNF0, TNF1, TNF2, G-CSF, Meg-CSF, thrombopoietin, stem cell factor, and erythropoietin. The  
5 pharmaceutical composition may further contain other agents which either enhance the activity of the protein or compliment its activity or use in treatment. Such additional factors and/or agents may be included in the pharmaceutical composition to produce a synergistic effect with protein of the invention, or to minimize side effects. Conversely, protein of the present invention may be included in formulations of the particular  
10 cytokine, lymphokine, other hematopoietic factor, thrombolytic or anti-thrombotic factor, or anti-inflammatory agent to minimize side effects of the cytokine, lymphokine, other hematopoietic factor, thrombolytic or anti-thrombotic factor, or anti-inflammatory agent.

A protein of the present invention may be active in multimers (e.g., heterodimers or homodimers) or complexes with itself or other proteins. As a result,  
15 pharmaceutical compositions of the invention may comprise a protein of the invention in such multimeric or complexed form.

The pharmaceutical composition of the invention may be in the form of a complex of the protein(s) of present invention along with protein or peptide antigens. The protein and/or peptide antigen will deliver a stimulatory signal to both B and T  
20 lymphocytes. B lymphocytes will respond to antigen through their surface immunoglobulin receptor. T lymphocytes will respond to antigen through the T cell receptor (TCR) following presentation of the antigen by MHC proteins. MHC and structurally related proteins including those encoded by class I and class II MHC genes on host cells will serve to present the peptide antigen(s) to T lymphocytes. The antigen  
25 components could also be supplied as purified MHC-peptide complexes alone or with co-stimulatory molecules that can directly signal T cells. Alternatively antibodies able to bind surface immunoglobulin and other molecules on B cells as well as antibodies able to bind the TCR and other molecules on T cells can be combined with the pharmaceutical composition of the invention.

30 The pharmaceutical composition of the invention may be in the form of a liposome in which protein of the present invention is combined, in addition to other pharmaceutically acceptable carriers, with amphipathic agents such as lipids which exist in aggregated form as micelles, insoluble monolayers, liquid crystals, or lamellar layers in aqueous solution. Suitable lipids for liposomal formulation include, without limitation,

monoglycerides, diglycerides, sulfatides, lysolecithin, phospholipids, saponin, bile acids, and the like. Preparation of such liposomal formulations is within the level of skill in the art, as disclosed, for example, in U.S. Patent No. 4,235,871; U.S. Patent No. 4,501,728; U.S. Patent No. 4,837,028; and U.S. Patent No. 4,737,323, all of which are incorporated herein  
5 by reference.

As used herein, the term "therapeutically effective amount" means the total amount of each active component of the pharmaceutical composition or method that is sufficient to show a meaningful patient benefit, i.e., treatment, healing, prevention or amelioration of the relevant medical condition, or an increase in rate of treatment, healing,  
10 prevention or amelioration of such conditions. When applied to an individual active ingredient, administered alone, the term refers to that ingredient alone. When applied to a combination, the term refers to combined amounts of the active ingredients that result in the therapeutic effect, whether administered in combination, serially or simultaneously.

In practicing the method of treatment or use of the present invention, a  
15 therapeutically effective amount of protein of the present invention is administered to a mammal having a condition to be treated. Protein of the present invention may be administered in accordance with the method of the invention either alone or in combination with other therapies such as treatments employing cytokines, lymphokines or other hematopoietic factors. When co-administered with one or more cytokines,  
20 lymphokines or other hematopoietic factors, protein of the present invention may be administered either simultaneously with the cytokine(s), lymphokine(s), other hematopoietic factor(s), thrombolytic or anti-thrombotic factors, or sequentially. If administered sequentially, the attending physician will decide on the appropriate sequence of administering protein of the present invention in combination with  
25 cytokine(s), lymphokine(s), other hematopoietic factor(s), thrombolytic or anti-thrombotic factors.

Administration of protein of the present invention used in the pharmaceutical composition or to practice the method of the present invention can be carried out in a variety of conventional ways, such as oral ingestion, inhalation, topical  
30 application or cutaneous, subcutaneous, intraperitoneal, parenteral or intravenous injection. Intravenous administration to the patient is preferred.

When a therapeutically effective amount of protein of the present invention is administered orally, protein of the present invention will be in the form of a tablet, capsule, powder, solution or elixir. When administered in tablet form, the pharmaceutical

composition of the invention may additionally contain a solid carrier such as a gelatin or an adjuvant. The tablet, capsule, and powder contain from about 5 to 95% protein of the present invention, and preferably from about 25 to 90% protein of the present invention. When administered in liquid form, a liquid carrier such as water, petroleum, oils of animal or plant origin such as peanut oil, mineral oil, soybean oil, or sesame oil, or synthetic oils may be added. The liquid form of the pharmaceutical composition may further contain physiological saline solution, dextrose or other saccharide solution, or glycols such as ethylene glycol, propylene glycol or polyethylene glycol. When administered in liquid form, the pharmaceutical composition contains from about 0.5 to 90% by weight of protein of the present invention, and preferably from about 1 to 50% protein of the present invention.

When a therapeutically effective amount of protein of the present invention is administered by intravenous, cutaneous or subcutaneous injection, protein of the present invention will be in the form of a pyrogen-free, parenterally acceptable aqueous solution. The preparation of such parenterally acceptable protein solutions, having due regard to pH, isotonicity, stability, and the like, is within the skill in the art. A preferred pharmaceutical composition for intravenous, cutaneous, or subcutaneous injection should contain, in addition to protein of the present invention, an isotonic vehicle such as Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, Lactated Ringer's Injection, or other vehicle as known in the art. The pharmaceutical composition of the present invention may also contain stabilizers, preservatives, buffers, antioxidants, or other additives known to those of skill in the art.

The amount of protein of the present invention in the pharmaceutical composition of the present invention will depend upon the nature and severity of the condition being treated, and on the nature of prior treatments which the patient has undergone. Ultimately, the attending physician will decide the amount of protein of the present invention with which to treat each individual patient. Initially, the attending physician will administer low doses of protein of the present invention and observe the patient's response. Larger doses of protein of the present invention may be administered until the optimal therapeutic effect is obtained for the patient, and at that point the dosage is not increased further. It is contemplated that the various pharmaceutical compositions used to practice the method of the present invention should contain about 0.01 g to about 100 mg (preferably about 0.1mg to about 10 mg, more preferably about 0.1 g to about 1 mg) of protein of the present invention per kg body weight.

The duration of intravenous therapy using the pharmaceutical composition of the present invention will vary, depending on the severity of the disease being treated and the condition and potential idiosyncratic response of each individual patient. It is contemplated that the duration of each application of the protein of the present invention will be in the range of 12 to 24 hours of continuous intravenous administration. Ultimately the attending physician will decide on the appropriate duration of intravenous therapy using the pharmaceutical composition of the present invention.

Protein of the invention may also be used to immunize animals to obtain polyclonal and monoclonal antibodies which specifically react with the protein. Such antibodies may be obtained using either the entire protein or fragments thereof as an immunogen. The peptide immunogens additionally may contain a cysteine residue at the carboxyl terminus, and are conjugated to a hapten such as keyhole limpet hemocyanin (KLH). Methods for synthesizing such peptides are known in the art, for example, as in R.P. Merrifield, *J. Amer.Chem.Soc.* 85, 2149-2154 (1963); J.L. Krstenansky, *et al.*, *FEBS Lett.* 211, 10 (1987). Monoclonal antibodies binding to the protein of the invention may be useful diagnostic agents for the immunodetection of the protein. Neutralizing monoclonal antibodies binding to the protein may also be useful therapeutics for both conditions associated with the protein and also in the treatment of some forms of cancer where abnormal expression of the protein is involved. In the case of cancerous cells or leukemic cells, neutralizing monoclonal antibodies against the protein may be useful in detecting and preventing the metastatic spread of the cancerous cells, which may be mediated by the protein.

For compositions of the present invention which are useful for bone, cartilage, tendon or ligament regeneration, the therapeutic method includes administering the composition topically, systematically, or locally as an implant or device. When administered, the therapeutic composition for use in this invention is, of course, in a pyrogen-free, physiologically acceptable form. Further, the composition may desirably be encapsulated or injected in a viscous form for delivery to the site of bone, cartilage or tissue damage. Topical administration may be suitable for wound healing and tissue repair. Therapeutically useful agents other than a protein of the invention which may also optionally be included in the composition as described above, may alternatively or additionally, be administered simultaneously or sequentially with the composition in the methods of the invention. Preferably for bone and/or cartilage formation, the composition would include a matrix capable of delivering the protein-containing

composition to the site of bone and/or cartilage damage, providing a structure for the developing bone and cartilage and optimally capable of being resorbed into the body. Such matrices may be formed of materials presently in use for other implanted medical applications.

5           The choice of matrix material is based on biocompatibility, biodegradability, mechanical properties, cosmetic appearance and interface properties. The particular application of the compositions will define the appropriate formulation. Potential matrices for the compositions may be biodegradable and chemically defined calcium sulfate, tricalciumphosphate, hydroxyapatite, polylactic acid, polyglycolic acid  
10 and polyanhydrides. Other potential materials are biodegradable and biologically well-defined, such as bone or dermal collagen. Further matrices are comprised of pure proteins or extracellular matrix components. Other potential matrices are nonbiodegradable and chemically defined, such as sintered hydroxapatite, bioglass, aluminates, or other ceramics. Matrices may be comprised of combinations of any of the  
15 above mentioned types of material, such as polylactic acid and hydroxyapatite or collagen and tricalciumphosphate. The bioceramics may be altered in composition, such as in calcium-aluminate-phosphate and processing to alter pore size, particle size, particle shape, and biodegradability.

          Presently preferred is a 50:50 (mole weight) copolymer of lactic acid and  
20 glycolic acid in the form of porous particles having diameters ranging from 150 to 800 microns. In some applications, it will be useful to utilize a sequestering agent, such as carboxymethyl cellulose or autologous blood clot, to prevent the protein compositions from disassociating from the matrix.

          A preferred family of sequestering agents is cellulosic materials such as  
25 alkylcelluloses (including hydroxyalkylcelluloses), including methylcellulose, ethylcellulose, hydroxyethylcellulose, hydroxypropylcellulose, hydroxypropyl-methylcellulose, and carboxymethylcellulose, the most preferred being cationic salts of carboxymethylcellulose (CMC). Other preferred sequestering agents include hyaluronic acid, sodium alginate, poly(ethylene glycol), polyoxyethylene oxide,  
30 carboxyvinyl polymer and poly(vinyl alcohol). The amount of sequestering agent useful herein is 0.5-20 wt%, preferably 1-10 wt% based on total formulation weight, which represents the amount necessary to prevent desorption of the protein from the polymer matrix and to provide appropriate handling of the composition, yet not so much that the progenitor cells are prevented from infiltrating the matrix, thereby providing the protein

the opportunity to assist the osteogenic activity of the progenitor cells.

In further compositions, proteins of the invention may be combined with other agents beneficial to the treatment of the bone and/or cartilage defect, wound, or tissue in question. These agents include various growth factors such as epidermal growth factor (EGF), platelet derived growth factor (PDGF), transforming growth factors (TGF- and TGF-), and insulin-like growth factor (IGF).

The therapeutic compositions are also presently valuable for veterinary applications. Particularly domestic animals and thoroughbred horses, in addition to humans, are desired patients for such treatment with proteins of the present invention.

10 The dosage regimen of a protein-containing pharmaceutical composition to be used in tissue regeneration will be determined by the attending physician considering various factors which modify the action of the proteins, e.g., amount of tissue weight desired to be formed, the site of damage, the condition of the damaged tissue, the size of a wound, type of damaged tissue (e.g., bone), the patient's age, sex, and diet, the  
15 severity of any infection, time of administration and other clinical factors. The dosage may vary with the type of matrix used in the reconstitution and with inclusion of other proteins in the pharmaceutical composition. For example, the addition of other known growth factors, such as IGF I (insulin like growth factor I), to the final composition, may also effect the dosage. Progress can be monitored by periodic assessment of tissue/bone  
20 growth and/or repair, for example, X-rays, histomorphometric determinations and tetracycline labeling.

Polynucleotides of the present invention can also be used for gene therapy. Such polynucleotides can be introduced either *in vivo* or *ex vivo* into cells for expression in a mammalian subject. Polynucleotides of the invention may also be administered by  
25 other known methods for introduction of nucleic acid into a cell or organism (including, without limitation, in the form of viral vectors or naked DNA).

Cells may also be cultured *ex vivo* in the presence of proteins of the present invention in order to proliferate or to produce a desired effect on or activity in such cells. Treated cells can then be introduced *in vivo* for therapeutic purposes.

30

Patent and literature references cited herein are incorporated by reference as if fully set forth.



What is claimed is:

1. An isolated protein encoded by a polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:1;
- (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:1 from nucleotide 19 to nucleotide 561;
- (c) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AK296\_1i deposited under accession number ATCC 98026;
- (d) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AK296\_1i deposited under accession number ATCC 98026;
- (e) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:2;
- (f) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:2 having biological activity, the fragment comprising eight consecutive amino acids of SEQ ID NO:2; and
- (g) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(f) above.

2. A composition comprising the protein of claim 1 and a pharmaceutically acceptable carrier.

3. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:2;
  - (b) the amino acid sequence of SEQ ID NO:2 from amino acid 3 to amino acid 181;
  - (c) fragments of the amino acid sequence of SEQ ID NO:2, each fragment comprising eight consecutive amino acids of SEQ ID NO:2; and
  - (d) the amino acid sequence encoded by the cDNA insert of clone AK296\_1i deposited under accession number ATCC 98026;
- the protein being substantially free from other mammalian proteins.

4. The protein of claim 3, wherein said protein comprises the amino acid sequence of SEQ ID NO:2.
5. The protein of claim 3, wherein said protein comprises a fragment of the amino acid sequence of SEQ ID NO:2, the fragment comprising eight consecutive amino acids of SEQ ID NO:2.
6. The protein of claim 3, wherein said protein comprises the amino acid sequence of SEQ ID NO:2 from amino acid 3 to amino acid 181.
7. A composition comprising the protein of claim 3 and a pharmaceutically acceptable carrier.
8. An isolated protein encoded by a polynucleotide selected from the group consisting of:
  - (a) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:21;
  - (b) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:21 from nucleotide 65 to nucleotide 490;
  - (c) a polynucleotide comprising the nucleotide sequence of SEQ ID NO:21 from nucleotide 137 to nucleotide 490;
  - (d) a polynucleotide comprising the nucleotide sequence of the full-length protein coding sequence of clone AS34\_1i deposited under accession number ATCC 98026;
  - (e) a polynucleotide encoding the full-length protein encoded by the cDNA insert of clone AS34\_1i deposited under accession number ATCC 98026;
  - (f) a polynucleotide comprising the nucleotide sequence of a mature protein coding sequence of clone AS34\_1i deposited under accession number ATCC 98026;
  - (g) a polynucleotide encoding a mature protein encoded by the cDNA insert of clone AS34\_1i deposited under accession number ATCC 98026;
  - (h) a polynucleotide encoding a protein comprising the amino acid sequence of SEQ ID NO:22;


(i) a polynucleotide encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:22 having biological activity, the *fragment comprising eight consecutive amino acids of SEQ ID NO:22; and*

(j) a polynucleotide which is an allelic variant of a polynucleotide of (a)-(g) above.

9. A protein comprising an amino acid sequence selected from the group consisting of:


- (a) the amino acid sequence of SEQ ID NO:22;
  - (b) fragments of the amino acid sequence of SEQ ID NO:22, each fragment comprising eight consecutive amino acids of SEQ ID NO:22; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone AS34\_1i deposited under accession number ATCC 98026;
- the protein being substantially free from other mammalian proteins.

Fig. 1A



2 / 2

Fig. 1B



## SEQUENCE LISTING

<110> Jacobs, Kenneth  
 McCoy, John M.  
 LaVallie, Edward R.  
 Collins-Racie, Lisa A.  
 Evans, Cheryl  
 Merberg, David  
 Treacy, Maurice  
 Bowman, Michael R.  
 Genetics Institute, Inc.

<120> SECRETED PROTEINS

<130> GI 6505

<140>

<141>

<160> 154

<170> PatentIn Ver. 2.0

<210> 1

<211> 1248

<212> DNA

<213> Homo sapiens

<400> 1

```

aaaggcctac gtcgacctat gaccatgatt acgccaagct tggcacgagg cagggagggtc 60
ctgaccccaa cgagcacttc tgacaatgag accagagact cctcaattat tgatccagga 120
actgagcaag atcttccttc cctgaaaat agttctgtta aagaataccg aatggaagtt 180
ccatcttcgt ttccagaaga catgtcaaat atcaggtcac agcatgcaga agaacagtcc 240
aacaatggta gatatgacga ttgtaaaagaa tttaaagacc tccactgttc caaggattct 300
accctagccg aggaagaatc tgagttccct tctacttcta tctctgcagt tctgtctgac 360
ttagctgact tgagaagctg tgatggccaa gctttgccct cccaggaccc tgaggttgct 420
ttatctctca gttgtggcca ttccagagga ctctttagtc atatgcagca acatgacatt 480
ttagataccc tgtgtaggac cattgaatct acaatccatg tcgtcacaag gatattctggc 540
aaaggaaacc aagctgtctc ttgacattag gtgtagcatg tctactttta agtccctcac 600
ccccaaaccc catgctgttt gtataagttt tgcttatttg tttttgtgct tcagtttgtc 660
cagtgtcttc tgcttgaatg gcaagataga tttataggct caattcttgg tcaggcagaa 720
ctccagatga aaaaaacttg catcttcagt atacttcta aagggcaatc agataatgga 780
tatgttttat gtaattaaga gttcacttta gtggctttca tttaatatgg ctgtctggga 840
agaacagggt tgccatgccc tgtacaatgt aatttaaact tacagcattt ttactgtgta 900
tgatatggtg tcctctgtgc cagttttgta ccttatagag gcagattgcc tccgatcgct 960
gtggttctta ttatcaaat taagtttact tgtatacgga acaacc?caa gaaatttgat 1020
tctgtaaaga atcctcttta gctgtggcct ggcagtatat aaatgggtgt ttatttaaca 1080
gaatacctgt ggaggaaata aagcacactt gatgtaaaaa taattgtttt atttttattg 1140
acatgactga ttgattgcta ttctgtgcac ttaattaaac tgattgtgat gactttttaa 1200
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaa 1248

```

<210> 2

<211> 181

<212> PRT

<213> Homo sapiens

<400> 2

```

Met Thr Met Ile Thr Pro Ser Leu Ala Arg Gly Arg Glu Val Leu Thr
  1           5           10          15

```

Pro Thr Ser Thr Ser Asp Asn Glu Thr Arg Asp Ser Ser Ile Ile Asp  
                   20                  25                  30  
 Pro Gly Thr Glu Gln Asp Leu Pro Ser Pro Glu Asn Ser Ser Val Lys  
                   35                  40                  45  
 Glu Tyr Arg Met Glu Val Pro Ser Ser Phe Ser Glu Asp Met Ser Asn  
                   50                  55                  60  
 Ile Arg Ser Gln His Ala Glu Glu Gln Ser Asn Asn Gly Arg Tyr Asp  
                   65                  70                  75                  80  
 Asp Cys Lys Glu Phe Lys Asp Leu His Cys Ser Lys Asp Ser Thr Leu  
                   85                  90                  95  
 Ala Glu Glu Glu Ser Glu Phe Pro Ser Thr Ser Ile Ser Ala Val Leu  
                   100                  105                  110  
 Ser Asp Leu Ala Asp Leu Arg Ser Cys Asp Gly Gln Ala Leu Pro Ser  
                   115                  120                  125  
 Gln Asp Pro Glu Val Ala Leu Ser Leu Ser Cys Gly His Ser Arg Gly  
                   130                  135                  140  
 Leu Phe Ser His Met Gln Gln His Asp Ile Leu Asp Thr Leu Cys Arg  
                   145                  150                  155                  160  
 Thr Ile Glu Ser Thr Ile His Val Val Thr Arg Ile Ser Gly Lys Gly  
                   165                  170                  175  
 Asn Gln Ala Ala Ser  
                   180

&lt;210&gt; 3

&lt;211&gt; 1706

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 3

gcgcagtga ggcagtggaa ccggaatata caaagagtgg tttgaaggag aaagaagcat 60  
 tgtggcttta taccctctgg gcctgggttt cctgaagtca ccaacacatag aggagagaga 120  
 aaatggctga gttaaagtac atttctggat ttgggaatga gtgttcttca gaggatcctc 180  
 gctgcccagg ttccctgcca gaaggacaga ataatcctca ggtctgcccc tacaatctct 240  
 atgctgagca gctctcagga tcggctttca cttgtccacg gacaccaat aagagaagct 300  
 ggctgtatag gattctacct tcagtttctc acaagccctt tgaatccatt gacgaaggcc 360  
 atgtcactca caactgggat gaagttgatc ctgacccata ccagcttaga tggaaacccat 420  
 ttgagattcc aaaagcatct cagaagaaag tagactttgt gagtggcctg cataccttgt 480  
 gtggagctgg agacataaag tctaacaatg ggcttgctat ccacattttc ctctgcaata 540  
 cctccatgga gaacagatgc ttttacaatt cagatgggga cttcttgatt gttccgcaga 600  
 aagggaaacct tctcatttac accgagtttg gcaagatgct tgtacagccc aatgagatct 660  
 gcgtcattca gagaggaatg cggttcagca tagatgtctt tgaggagacc aggggctaca 720  
 tcttggaggt ctatggtgac cactttgagt tacctgacct tggaccaatt ggggccaatg 780  
 gcttgcccaa tcctcgatg ttcttgatac ccattgcctg gtatgaggat cgccaagtac 840  
 caggtgggta cacggtcatt aataaatacc agggcaagct gtttgctgcc aaacaggatg 900  
 tctcccggtt caatgttgtg gcctggcacg ggaattatac accctacaag tacaacctga 960  
 agaatttcac ggttatcaac tcagtgccct ttgacctgac agaccatcc attttcacag 1020  
 tattgactgc taagtctgtc cgcctgggag tggccattgc tgattttgtc atcttccac 1080  
 ctcgatgggg ggttgctgat aagaccttca ggctcctta ttacctagg aactgcatga 1140  
 gtgagttcat gggactcatc cgaggtcact atgaggcaaa gcaaggtggg ttcctgccag 1200

```

ggggaggag tctacacagc acaatgaccc cccatggacc tgatgctgac tgctttgaga 1260
agggcagcaa ggtcaagctg gcacctgaga ggattgccga tggcaccatg gcatttatgt 1320
ttgaatcatc ttttaagtctg gcggtcacaa agtggggact caaggcctcc aggtgttttg 1380
atgagaacta ccacaagtgc tgggagccac tcaagagcca cttcactccc aactccagga 1440
acccagcaga acctaattga gactggaaca ttgctaccat aattaagagt agatttgtga 1500
agatttcttc agaattctcat gctttctggt agtattggag gagggggttg gttaaaatga 1560
aaattcactt ttcatagtca agtaactcag aacttttatg gaaacgcatt tgcaaagtgc 1620
tatggctgtc accttaatta ctcaataaac ttgctggtgt tctgtggaaa aaaaaaaaaa 1680
aaaaaaaaaa aaaaaaaaaa aaaaaa 1706

```

<210> 4  
 <211> 445  
 <212> PRT  
 <213> Homo sapiens

<400> 4  
 Met Ala Glu Leu Lys Tyr Ile Ser Gly Phe Gly Asn Glu Cys Ser Ser  
     1                    5                    10                    15  
  
 Glu Asp Pro Arg Cys Pro Gly Ser Leu Pro Glu Gly Gln Asn Asn Pro  
             20                    25                    30  
  
 Gln Val Cys Pro Tyr Asn Leu Tyr Ala Glu Gln Leu Ser Gly Ser Ala  
             35                    40                    45  
  
 Phe Thr Cys Pro Arg Ser Thr Asn Lys Arg Ser Trp Leu Tyr Arg Ile  
             50                    55                    60  
  
 Leu Pro Ser Val Ser His Lys Pro Phe Glu Ser Ile Asp Glu Gly His  
             65                    70                    75                    80  
  
 Val Thr His Asn Trp Asp Glu Val Asp Pro Asp Pro Asn Gln Leu Arg  
             85                    90                    95  
  
 Trp Lys Pro Phe Glu Ile Pro Lys Ala Ser Gln Lys Lys Val Asp Phe  
             100                    105                    110  
  
 Val Ser Gly Leu His Thr Leu Cys Gly Ala Gly Asp Ile Lys Ser Asn  
             115                    120                    125  
  
 Asn Gly Leu Ala Ile His Ile Phe Leu Cys Asn Thr Ser Met Glu Asn  
             130                    135                    140  
  
 Arg Cys Phe Tyr Asn Ser Asp Gly Asp Phe Leu Ile Val Pro Gln Lys  
             145                    150                    155                    160  
  
 Gly Asn Leu Leu Ile Tyr Thr Glu Phe Gly Lys Met Leu Val Gln Pro  
             165                    170                    175  
  
 Asn Glu Ile Cys Val Ile Gln Arg Gly Met Arg Phe Ser Ile Asp Val  
             180                    185                    190  
  
 Phe Glu Glu Thr Arg Gly Tyr Ile Leu Glu Val Tyr Gly Val His Phe  
             195                    200                    205  
  
 Glu Leu Pro Asp Leu Gly Pro Ile Gly Ala Asn Gly Leu Ala Asn Pro  
             210                    215                    220  
  
 Arg Asp Phe Leu Ile Pro Ile Ala Trp Tyr Glu Asp Arg Gln Val Pro  
             225                    230                    235                    240



Gly Gly Tyr Thr Val Ile Asn Lys Tyr Gln Gly Lys Leu Phe Ala Ala  
 245 250 255  
 Lys Gln Asp Val Ser Pro Phe Asn Val Val Ala Trp His Gly Asn Tyr  
 260 265 270  
 Thr Pro Tyr Lys Tyr Asn Leu Lys Asn Phe Met Val Ile Asn Ser Val  
 275 280 285  
 Ala Phe Asp His Ala Asp Pro Ser Ile Phe Thr Val Leu Thr Ala Lys  
 290 295 300  
 Ser Val Arg Pro Gly Val Ala Ile Ala Asp Phe Val Ile Phe Pro Pro  
 305 310 315 320  
 Arg Trp Gly Val Ala Asp Lys Thr Phe Arg Pro Pro Tyr Tyr His Arg  
 325 330 335  
 Asn Cys Met Ser Glu Phe Met Gly Leu Ile Arg Gly His Tyr Glu Ala  
 340 345 350  
 Lys Gln Gly Gly Phe Leu Pro Gly Gly Gly Ser Leu His Ser Thr Met  
 355 360 365  
 Thr Pro His Gly Pro Asp Ala Asp Cys Phe Glu Lys Ala Ser Lys Val  
 370 375 380  
 Lys Leu Ala Pro Glu Arg Ile Ala Asp Gly Thr Met Ala Phe Met Phe  
 385 390 395 400  
 Glu Ser Ser Leu Ser Leu Ala Val Thr Lys Trp Gly Leu Lys Ala Ser  
 405 410 415  
 Arg Cys Leu Asp Glu Asn Tyr His Lys Cys Trp Glu Pro Leu Lys Ser  
 420 425 430  
 His Phe Thr Pro Asn Ser Arg Asn Pro Ala Glu Pro Asn  
 435 440 445

&lt;210&gt; 5

&lt;211&gt; 912

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 5

gtggaaattt gtggctagtg tgatttttgt ttgttttcctt ttaagtactg ttgatcagtt 60  
 gtgacactta ctggttaaac ttacgttgct aaagatttct ctataataag ccacacatta 120  
 tatttagact atattaaggg accttggttt tcttctagat agcagctgtc ccaaagaaaa 180  
 tatttcttct ttgtctgtta agatttagct attatctgcc agttgttaag aggttttggt 240  
 tccaaactca accagcaatg ttgagagctg aacttaagat agctgttgta ctttttgctt 300  
 tccatctgtt actgtccttc attcttggtt cctactatc tataaacagc tgctgtgaag 360  
 aagaaaagtt gaataagagt tggcttaaat tttaaaaaag aaaaagaaaa ttgaggtttt 420  
 aggattttca tggtaacaag ctctgggtata agctaaggct ggcaagtcca gatactaaaa 480  
 tattatttga tcatatcttg gatccttttg aaaaagttaa gactatatga aggtaaatta 540  
 gaaataagta tgaatattaa taaaatagca tttatcttat ttctctattt tatgttgkga 600  
 mttaacctaa ttttattttt ttaamatttt yttatttytt ataatatgaa tgstgatatt 660  
 taaaggtaga tytatgttgt attytttgtg ttttyttaatt gtttaamtyt taagattatt 720  
 tgtgatytgg atttatgtat ttgttagata mtaacgratt gttaaaatgg ratgcaagtt 780

tttcaaaagc ccaggtytaa atgtaatggk tggtttattg tttataacc ccagcccatc 840  
atttttctgtg taaatyataa acaataaaca gratatamtc ggtggtcatt tytaaaaaaa 900  
aaaaaaaaaa aa 912

<210> 6  
<211> 45  
<212> PRT  
<213> Homo sapiens

<400> 6  
Met Leu Arg Ala Glu Leu Lys Ile Ala Val Val Leu Phe Ala Phe His  
1 5 10 15  
Leu Leu Leu Ser Phe Ile Leu Gly Ser Leu Leu Ser Ile Asn Ser Cys  
20 25 30  
Cys Glu Glu Glu Lys Leu Asn Lys Ser Trp Leu Lys Phe  
35 40 45

<210> 7  
<211> 1767  
<212> DNA  
<213> Homo sapiens

<400> 7  
cgaagatgaa attccttata ttgcattttt tgggtgggtt tcacctttta tccctgtgct 60  
ctgggaaagc tatatgcaag aatggcatct ctaagaggac ttttgaagaa ataaaagaag 120  
aaatagccag ctgtggagat gttgctaaag caatcatcaa cctagctgtt tatggtaaag 180  
cccagaacag atcctatgag cgattggcac ttctggttga tactgttga cccagactga 240  
gtggctccaa gaacctagaa aaagccatcc aaattatgta ccaaaccctg cagcaagatg 300  
ggctggagaa agttcacctg gagccagtga gaatacccca ctgggagagg ggagaagaat 360  
cagctgtgat gctggagcca agaattcata agatagccat cctgggtctt ggcagcagca 420  
ttgggactcc tccagaaggc attacagcag aagttctggt ggtgacctct ttcgatgaac 480  
tgcagagaag ggcctcagaa gcaagaggga agattgttgt ttataaccaa ccttacatca 540  
actactcaag gacggtgcaa taccgaacgc agggggcggt ggaagctgcc aaggtggggg 600  
ctttggcatc tctcattcga tccgtggcct ccttctccat ctacagtctt cacacaggta 660  
ttcaggaata ccaggatggc gtgcccaga ttccaacagc ctgtattacg gtggaagatg 720  
cagaaatgat gtcaagaatg gcttctcatg ggatcaaaat tgcattcag ctaaagatgg 780  
gggcaaagac ctaccagat actgattcct tcaacactgt agcagagatc actgggagca 840  
aatatccaga acaggttgta ctggctcagtg gacatctgga cagctgggat gttgggcagg 900  
gtgccatgga tgatggcggt ggagccttta tatcatggga agcactctca cttattaaag 960  
atcttgggct gcgtccaaag aggactctgc ggctgggtgt ctggactgca ggagaacaag 1020  
gtggagtggg tgocctccag tattatcagt tacacaaggt aaatatttcc aactacagtc 1080  
tggtgatgga gtctgacgca ggaaccttct taccactgg gctgcaattc actggcagtg 1140  
aaaaggccag ggccatcatg gaggagggtta tgagcctgct gcagcccctc aatatcactc 1200  
aggctctgag ccatggagaa gggacagaca tcaacttttg gatccaagct ggagtgcctg 1260  
gagccagtct acttgatgac ttatacaagt atttcttctt ccactactcc cacggagaca 1320  
ccatgactgt catggatcca aagcagatga atgttgctgc tgctgtttgg gctgtgtgtt 1380  
cttatgttgt tgcagacatg gaagaaatgc tgcctaggtc ctagaaacag taagaaagaa 1440  
acgttttcat gcttctggcc aggaatcctg ggtctgcaac tttggaaaac tctcttccac 1500  
ataacaattt catccaattc atcttcaaag cacaactcta tttcatgctt tctgttatta 1560  
tctttcttga tactttccaa attctctgat tctagaaaaa ggaatcatc tcccctccct 1620  
cccaccacat agaatcaaca tatggtaggg attacagtgg gggcatttct ttataccacc 1680  
tcttaaaaac attgtttcca ctttaaaagt aaacacttaa taaatttttg gaagatctct 1740  
gaaaaaaaaa aaaaaaaaaa aaaaaaa 1767

<210> 8  
<211> 472  
<212> PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 8

Met Lys Phe Leu Ile Phe Ala Phe Phe Gly Gly Val His Leu Leu Ser  
 1 5 10 15

Leu Cys Ser Gly Lys Ala Ile Cys Lys Asn Gly Ile Ser Lys Arg Thr  
 20 25 30

Phe Glu Glu Ile Lys Glu Glu Ile Ala Ser Cys Gly Asp Val Ala Lys  
 35 40 45

Ala Ile Ile Asn Leu Ala Val Tyr Gly Lys Ala Gln Asn Arg Ser Tyr  
 50 55 60

Glu Arg Leu Ala Leu Leu Val Asp Thr Val Gly Pro Arg Leu Ser Gly  
 65 70 75 80

Ser Lys Asn Leu Glu Lys Ala Ile Gln Ile Met Tyr Gln Asn Leu Gln  
 85 90 95

Gln Asp Gly Leu Glu Lys Val His Leu Glu Pro Val Arg Ile Pro His  
 100 105 110

Trp Glu Arg Gly Glu Glu Ser Ala Val Met Leu Glu Pro Arg Ile His  
 115 120 125

Lys Ile Ala Ile Leu Gly Leu Gly Ser Ser Ile Gly Thr Pro Pro Glu  
 130 135 140

Gly Ile Thr Ala Glu Val Leu Val Val Thr Ser Phe Asp Glu Leu Gln  
 145 150 155 160

Arg Arg Ala Ser Glu Ala Arg Gly Lys Ile Val Val Tyr Asn Gln Pro  
 165 170 175

Tyr Ile Asn Tyr Ser Arg Thr Val Gln Tyr Arg Thr Gln Gly Ala Val  
 180 185 190

Glu Ala Ala Lys Val Gly Ala Leu Ala Ser Leu Ile Arg Ser Val Ala  
 195 200 205

Ser Phe Ser Ile Tyr Ser Pro His Thr Gly Ile Gln Glu Tyr Gln Asp  
 210 215 220

Gly Val Pro Lys Ile Pro Thr Ala Cys Ile Thr Val Glu Asp Ala Glu  
 225 230 235 240

Met Met Ser Arg Met Ala Ser His Gly Ile Lys Ile Val Ile Gln Leu  
 245 250 255

Lys Met Gly Ala Lys Thr Tyr Pro Asp Thr Asp Ser Phe Asn Thr Val  
 260 265 270

Ala Glu Ile Thr Gly Ser Lys Tyr Pro Glu Gln Val Val Leu Val Ser  
 275 280 285

Gly His Leu Asp Ser Trp Asp Val Gly Gln Gly Ala Met Asp Asp Gly  
 290 295 300

Gly Gly Ala Phe Ile Ser Trp Glu Ala Leu Ser Leu Ile Lys Asp Leu  
 305 310 315 320  
 Gly Leu Arg Pro Lys Arg Thr Leu Arg Leu Val Leu Trp Thr Ala Gly  
 325 330 335  
 Glu Gln Gly Gly Val Gly Ala Phe Gln Tyr Tyr Gln Leu His Lys Val  
 340 345 350  
 Asn Ile Ser Asn Tyr Ser Leu Val Met Glu Ser Asp Ala Gly Thr Phe  
 355 360 365  
 Leu Pro Thr Gly Leu Gln Phe Thr Gly Ser Glu Lys Ala Arg Ala Ile  
 370 375 380  
 Met Glu Glu Val Met Ser Leu Leu Gln Pro Leu Asn Ile Thr Gln Val  
 385 390 395 400  
 Leu Ser His Gly Glu Gly Thr Asp Ile Asn Phe Trp Ile Gln Ala Gly  
 405 410 415  
 Val Pro Gly Ala Ser Leu Leu Asp Asp Leu Tyr Lys Tyr Phe Phe Phe  
 420 425 430  
 His His Ser His Gly Asp Thr Met Thr Val Met Asp Pro Lys Gln Met  
 435 440 445  
 Asn Val Ala Ala Ala Val Trp Ala Val Val Ser Tyr Val Val Ala Asp  
 450 455 460  
 Met Glu Glu Met Leu Pro Arg Ser  
 465 470

<210> 9  
 <211> 571  
 <212> DNA  
 <213> Homo sapiens

<400> 9  
 cacgagggt tttgacaagg cctatgttgt ccttggccag tttctggtgc taaagaaaga 60  
 tgaagacctc ttccgggaat ggctgaaaga cacttgtggc gccaacgcca agcagtcctg 120  
 ggactgcttc ggatgccttc gagagtgggt cgacgccttc ttgtgatgct ctctgggaag 180  
 ctctcaatcc ccagccctca tccagagttt gcagccgagt agggactcct cccctgtcct 240  
 ctacgaagga aaagattgct attgtcgtac tcacctccga cgtactccgg ggtcttttgg 300  
 gagttttctc ccctaaccat ttcaactttt tttggattct cgctcttgca tgcctcccc 360  
 gtcctttttc ccttgccagt tccctgggtga cagttaccag ctttctgaa tggattcccg 420  
 gccccatccc tcacccccac cctcactttc aatccgtttg ataccatttg gtccttttt 480  
 tggcagaaca gtcactgtcc ttgtaaagtt ttttagatca ataaagtcag tggctttcaa 540  
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa a 571

<210> 10  
 <211> 124  
 <212> PRT  
 <213> Homo sapiens

<400> 10  
 Lys Thr Leu Val Ala Pro Thr Pro Ser Ser Pro Gly Thr Ala Ser Asp  
 1 5 10 15

Ala Phe Glu Ser Gly Ala Thr Pro Ser Cys Asp Ala Leu Trp Glu Ala  
 20 25 30

Leu Asn Pro Gln Pro Ser Ser Arg Val Cys Ser Arg Val Gly Thr Pro  
 35 40 45

Pro Leu Ser Ser Thr Lys Glu Lys Ile Ala Ile Val Val Leu Thr Ser  
 50 55 60

Asp Val Leu Arg Gly Leu Leu Gly Val Phe Ser Pro Asn His Phe Asn  
 65 70 75 80

Phe Phe Trp Ile Leu Ala Leu Ala Cys Leu Pro Arg Pro Phe Ser Leu  
 85 90 95

Ala Ser Ser Leu Val Thr Val Thr Ser Phe Pro Glu Trp Ile Pro Gly  
 100 105 110

Pro Ile Pro His Pro His Pro His Phe Gln Ser Val  
 115 120

<210> 11  
 <211> 1713  
 <212> DNA  
 <213> Homo sapiens

<400> 11  
 gggcaggata ttagaaatgg ctactcccca gtcaattttc atctttgcaa tctgcatttt 60  
 aatgataaca gaattaattc tggcctcaaa aagctactat gatattcttag gtgtgccaaa 120  
 atcggcatca gagcgccaaa tcaagaaggc ctttcacaag ttggccatga agtaccaccc 180  
 tgacaaaaat aagagcccag atgctgaagc aaaattcaga gagattgcag aagcatatga 240  
 aacactctca gatgctaata gacgaaaaga gtatgataca cttggacaca gtgcttttac 300  
 tagtggttaa ggacaaagag gtagtggaag ttcttttgag cagtcattta acttcaattt 360  
 tgatgactta tttaaagact ttggcttttt tggcctcaaa caaacactg gatccaagaa 420  
 gcgttttgaa aatcattttc agacacgcca ggatgggtgt tccagtagac aaaggcatca 480  
 ttccaagaa tttctttttg gaggtggatt atttgatgac atgtttgaag atatggagaa 540  
 aatgtttttt tttagtgtt ttgactctac caatcagcat acagtacaga ctgaaaatag 600  
 atttcaagga tctagcaagc actgcaggac tgtcactcaa cgaagaggaa atatgggtac 660  
 tacatacact gactgttcag gacagtagt cttattctat tctcactaaa tccaactggt 720  
 tgactcttcc tcattatctt tgatgctaaa caattttctg tgaactattt tgacaagtgc 780  
 atgatttcac tttaaacaat ttgatatagc tattaatat atttaagggt tttttttttt 840  
 gacaaattca acattcaacg agtagacaaa atgctaatta ttccctgat taggaaagt 900  
 tctttaaaaa acacgtaatt ttgcctagt ctttttctct acctgccctt gggctcacta 960  
 atatcaccag tattattacc aagaaaatat tgagtttacc tgattaaact ttaaaagtta 1020  
 attgtagatt taaattgtgt gaacctaatg atttttgcag tgaacacctt actaattcaa 1080  
 agttgcatgt tctatgacat ctgtgacttg cgttcagag tgtagatgaa actgtataat 1140  
 tgagtcattc agtaaaggag aacagtatct tggtaattg ctactgaaag gttgagaaag 1200  
 gaatggtttg atatttacca cagcgtctgt cctttctaca gtagaactgg ggtaaaggaa 1260  
 atggttttat tgccatagt catttaggct ggaaaaaagt tgaactta acgaaatatt 1320  
 gccaaagat tgttatgtgt ttgggtccag cctaaaaatg attttgtagt gttgaaatca 1380  
 tagctactta catagctttt tcatttttct ttcttagttg ttggcactct taggtcttag 1440  
 tatggattta tgtgtttgtg tgtgtgtagt ttatcctctc tctcatcttt atctagagat 1500  
 tgactgatac ctcatctctg ttgtaaaacc agccagtaat ttctgtgcaa ccttactatg 1560  
 tgcaatattt ttaaactctg agaaatgtgt gcttttgttt tcggatagac ttatttcttt 1620  
 agttctgcac ttttccacat tatactccat atgagtatta atcctatgga tacatatcaa 1680  
 aacaagtgtc tcataaaaaa aaaaaaaaaa aaa 1713

<210> 12  
 <211> 223

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 12

Met Ala Thr Pro Gln Ser Ile Phe Ile Phe Ala Ile Cys Ile Leu Met  
 1 5 10 15

Ile Thr Glu Leu Ile Leu Ala Ser Lys Ser Tyr Tyr Asp Ile Leu Gly  
 20 25 30

Val Pro Lys Ser Ala Ser Glu Arg Gln Ile Lys Lys Ala Phe His Lys  
 35 40 45

Leu Ala Met Lys Tyr His Pro Asp Lys Asn Lys Ser Pro Asp Ala Glu  
 50 55 60

Ala Lys Phe Arg Glu Ile Ala Glu Ala Tyr Glu Thr Leu Ser Asp Ala  
 65 70 75 80

Asn Arg Arg Lys Glu Tyr Asp Thr Leu Gly His Ser Ala Phe Thr Ser  
 85 90 95

Gly Lys Gly Gln Arg Gly Ser Gly Ser Ser Phe Glu Gln Ser Phe Asn  
 100 105 110

Phe Asn Phe Asp Asp Leu Phe Lys Asp Phe Gly Phe Phe Gly Gln Asn  
 115 120 125

Gln Asn Thr Gly Ser Lys Lys Arg Phe Glu Asn His Phe Gln Thr Arg  
 130 135 140

Gln Asp Gly Gly Ser Ser Arg Gln Arg His His Phe Gln Glu Phe Ser  
 145 150 155 160

Phe Gly Gly Gly Leu Phe Asp Asp Met Phe Glu Asp Met Glu Lys Met  
 165 170 175

Phe Ser Phe Ser Gly Phe Asp Ser Thr Asn Gln His Thr Val Gln Thr  
 180 185 190

Glu Asn Arg Phe His Gly Ser Ser Lys His Cys Arg Thr Val Thr Gln  
 195 200 205

Arg Arg Gly Asn Met Val Thr Thr Tyr Thr Asp Cys Ser Gly Gln  
 210 215 220

&lt;210&gt; 13

&lt;211&gt; 505

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (430)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (436)

<220>  
 <221> unsure  
 <222> (452)

<400> 13  
 gaattcggca cgagggcgcg gggtcggywa tggcgscggc agccgaaggc gtactggcga 60  
 cccggagtga tgagcccgcc cgagacgatg ccscctgga gacagctgag gaarcaaagg 120  
 agcctgctga aagctgacat cactgagctc tgccgggaca tgttctccaa aatggccact 180  
 tacctgactg gggaaactgac ggccaccagt gaagactata agctcctgga aaatatgaat 240  
 aaactcacca gcttgaagta tyttgaaatg aaagatatg ctataaacat tagtaggaac 300  
 ttaaaggact taaaccagaa atatgctgga ctgcagcctt atytggattc agattcaatg 360  
 ttcattggaa gaggaggtag cagctttttg agcaggcagc ttacaagttg grtgcmtwtt 420  
 tcaaaaaaan tggaanccca agtacaagaa gntggagaag cgatgagaaa attattttta 480  
 tgggacagag tttttttttt ttaat 505

<210> 14  
 <211> 144  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> UNSURE  
 <222> (65)

<220>  
 <221> UNSURE  
 <222> (115)..(117)

<220>  
 <221> UNSURE  
 <222> (121)

<220>  
 <221> UNSURE  
 <222> (123)

<220>  
 <221> UNSURE  
 <222> (128)

<400> 14  
 Met Ser Pro Pro Glu Thr Met Pro Pro Trp Arg Gln Leu Arg Lys Gln  
 1 5 10 15  
 Arg Ser Leu Leu Lys Ala Asp Ile Thr Glu Leu Cys Arg Asp Met Phe  
 20 25 30  
 Ser Lys Met Ala Thr Tyr Leu Thr Gly Glu Leu Thr Ala Thr Ser Glu  
 35 40 45  
 Asp Tyr Lys Leu Leu Glu Asn Met Asn Lys Leu Thr Ser Leu Lys Tyr  
 50 55 60  
 Xaa Glu Met Lys Asp Ile Ala Ile Asn Ile Ser Arg Asn Leu Lys Asp  
 65 70 75 80  
 Leu Asn Gln Lys Tyr Ala Gly Leu Gln Pro Tyr Leu Asp Ser Asp Ser  
 85 90 95  
 Met Phe Ile Gly Arg Ala Gly Ser Ser Phe Leu Ser Arg Gln Leu Thr

	100		105		110
Ser Trp Xaa Xaa Xaa Ser Lys Lys Xaa Glu Xaa Gln Val Gln Glu Xaa					
115		120		125	
Gly Glu Ala Met Arg Lys Leu Phe Leu Trp Asp Arg Val Phe Phe Phe					
130	135		140		

<210> 15  
 <211> 315  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> unsure  
 <222> (65)

<220>  
 <221> unsure  
 <222> (163)

<220>  
 <221> unsure  
 <222> (269)

<220>  
 <221> unsure  
 <222> (275)

<400> 15  
 gcagcttatac accttgtgaa tgcggtaac ttacttttcc ataataattgc aaataacata 60  
 aaatntttaa ataattccaa gctgagtttt ctagattgag cagaaatggt gaaaggagta 120  
 ttgataactt ggcgtatgtg atgggcccc cttgtttatt ttntatgtga gtcacattga 180  
 catgcatga gtttggggaa atgtgatgaa aacaaagact agatgggtat gtgtgtttat 240  
 gtgttgggta gggaggtgac gattgccant catanaataa aggattttat aaaataccaa 300  
 aaaaaaaaa aaaaa 315

<210> 16  
 <211> 1719  
 <212> DNA  
 <213> Homo sapiens

<400> 16  
 aatttcacag ttacagtga gaagcagaag caaaagaatt aaccagctct tcagtcaagc 60  
 aaatcctcta ctcaccatgc ttccctctgc cattcatttc tatctccttc cccttgcag 120  
 catcctaatag aaaagctgtt tggcttttaa aaatgatgcc acagaaatcc tttattcaca 180  
 tgtgggttaa cctgttccag cacaccccag cagcaacagc acgttgaatc aagccagaaa 240  
 tggaggcagg catttcagta acactggact ggatcggaac actcgggttc aagtgggttg 300  
 ccgggaactg cgttccacca aatacatctc tgaaggccag tgcaccagca tcagccctct 360  
 gaaggagctg gtgtgtgctg gcgagtgttt gcccttgcca gtgctcccta actggattgg 420  
 agggagctat ggaacaaagt actggagcag gaggagctcc caggagtggc ggtgtgtcaa 480  
 tgacaaaacc cgtaccaga gaatccagct gcagtgccaa gatggcagca cacgcacctc 540  
 caaaatcaca gtagtcactg cctgcaagtg caagaggtac acccggcagc acaacgagtc 600  
 cagtcaaac tttgagagca tgtcacctgc caagccagtc cagcatcaca gagagcggaa 660  
 aagagccagc aaatccagca agcacagcat gagttagaac tcagactccc ataactagac 720  
 ttactagtaa ccactgtgtt tacagatttg attgcttgga agactcaagc ctgccactgc 780  
 tgtttttctc cttgaaagta tatgctttct gctttgatca aaccagcaa gctgtcttaa 840  
 gtatcaggac cttctttggg aatagttttt ccttttcaag tttttcaaga ttaggtata 900  
 tccatgaatg caatttgcac ttaattcca cgtatcctgt agttttaatt cctcattgtt 960



```

cttaaaagac tgttgatact ataaacatca gtgaatcatt atattttaaa acagaaaagg 1020
gcttctcaga taccctccat ctactggccc atcccctctc ctaaacaaaa ctccttcaaa 1080
acaggttaaa aaaaatatgt tgtcatgaat cttcacagta acatttcaga aagggtgcttt 1140
tttggtactc ttcattggga cagtttagca gccatgagtg atcttccttt gaaagagaat 1200
gaaagaccct gtgacatttc acttcaaaaa taagccctgt agctctttac ggtcgcatag 1260
tatgaaatta taccctgcat gctgaccctc gcttggaatg gaatgccaga aatgcatggc 1320
agcagctaata aagtaaagct gattaactat ttatttgtca atgttattat ttaatgagct 1380
ttcacatgtg atttggttca aaactttaat tttttaatgt tttgaaactt tttcatggac 1440
ctaaatattt tcctatatga tttgtggttg attagaaata tgaaatacat gttgtagata 1500
tgtaaaatga atattttagt ctccttatta catatatgtt catggtgaac tttatcaata 1560
gtatggatct ttttaaatca ataagatgct ttgtaaagtt gaaataagta atactttctt 1620
gtttaatctg tgcaatcaga aggtgtcttg accttcaatt caattggttt cttttaacaa 1680
aaataaacac tgctaaaagt taaaaaaaaa aaaaaaaaaa 1719

```

&lt;210&gt; 17

&lt;211&gt; 206

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 17

```

Met Leu Pro Pro Ala Ile His Phe Tyr Leu Leu Pro Leu Ala Cys Ile
  1             5             10             15

```

```

Leu Met Lys Ser Cys Leu Ala Phe Lys Asn Asp Ala Thr Glu Ile Leu
      20             25             30

```

```

Tyr Ser His Val Val Lys Pro Val Pro Ala His Pro Ser Ser Asn Ser
      35             40             45

```

```

Thr Leu Asn Gln Ala Arg Asn Gly Gly Arg His Phe Ser Asn Thr Gly
      50             55             60

```

```

Leu Asp Arg Asn Thr Arg Val Gln Val Gly Cys Arg Glu Leu Arg Ser
      65             70             75             80

```

```

Thr Lys Tyr Ile Ser Asp Gly Gln Cys Thr Ser Ile Ser Pro Leu Lys
      85             90             95

```

```

Glu Leu Val Cys Ala Gly Glu Cys Leu Pro Leu Pro Val Leu Pro Asn
      100            105            110

```

```

Trp Ile Gly Gly Gly Tyr Gly Thr Lys Tyr Trp Ser Arg Arg Ser Ser
      115            120            125

```

```

Gln Glu Trp Arg Cys Val Asn Asp Lys Thr Arg Thr Gln Arg Ile Gln
      130            135            140

```

```

Leu Gln Cys Gln Asp Gly Ser Thr Arg Thr Tyr Lys Ile Thr Val Val
      145            150            155            160

```

```

Thr Ala Cys Lys Cys Lys Arg Tyr Thr Arg Gln His Asn Glu Ser Ser
      165            170            175

```

```

His Asn Phe Glu Ser Met Ser Pro Ala Lys Pro Val Gln His His Arg
      180            185            190

```

```

Glu Arg Lys Arg Ala Ser Lys Ser Ser Lys His Ser Met Ser
      195            200            205

```

<210> 18  
 <211> 676  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> unsure  
 <222> (35)

<220>  
 <221> unsure  
 <222> (151)

<220>  
 <221> unsure  
 <222> (165)

<220>  
 <221> unsure  
 <222> (229)

<220>  
 <221> unsure  
 <222> (245)

<220>  
 <221> unsure  
 <222> (612)

<220>  
 <221> unsure  
 <222> (625)

<400> 18  
 gagattttca gcaccctgcg atatgcaagc cgagntcagc gggtcaccac ccgaccacag 60  
 gcccccaagt ttcctgtggc aaagcagccc cagcgtttgg agacagagat gctgcagctc 120  
 caggaggaga accgtcgcct gcagttccag ntggaccaa tggantgcaa ggcctcaggg 180  
 ttcagtggag ccgggtggc ctgggccag cggaacctgt acgggatgnt acaggagttt 240  
 catgntagag aatgagaggc tcaggaaaga aaagagccag ctgcagaata gccgagagct 300  
 agcccagaat ggcagcgca tcctggccca gcaggtccat gcactagaga rgcgtctcct 360  
 ctctgcctgc taccatcacc agcagggtcc tggcctgacc ccaccgtgtc cctgcttgat 420  
 ggccccagct ccccttgcc atgcactgcc acccctctac tctgccccct gctgccacat 480  
 ctgcccactg tgtckagtgc ccctggccca ctgggyykgc ctgscmaggg ggcaccacc 540  
 ttgccccagc ctctctctg gggctctgar gagtcagaaa tagaccagac gtggtttctt 600  
 ggttctcagg anggttttta gtttnaggag agggacggta gaagaacctt tttgttgcaa 660  
 aaagaagggg accaag 676

<210> 19  
 <211> 218  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> UNSURE  
 <222> (5)

<220>  
 <221> UNSURE  
 <222> (43)

<220>  
 <221> UNSURE  
 <222> (48)

<220>  
 <221> UNSURE  
 <222> (69)

<220>  
 <221> UNSURE  
 <222> (75)

<220>  
 <221> UNSURE  
 <222> (110)

<220>  
 <221> UNSURE  
 <222> (158)

<220>  
 <221> UNSURE  
 <222> (165) .. (166)

<220>  
 <221> UNSURE  
 <222> (168)

<220>  
 <221> UNSURE  
 <222> (183)

<220>  
 <221> UNSURE  
 <222> (187)

<220>  
 <221> UNSURE  
 <222> (197)

<220>  
 <221> UNSURE  
 <222> (201)

<400> 19  
 Met Gln Ala Glu Xaa Ser Gly Ser Pro Pro Asp His Arg Pro Pro Ser  
     1                    5                    10                    15  
 Phe Leu Trp Gln Ser Ser Pro Ser Val Trp Arg Gln Arg Cys Cys Ser  
     20                    25                    30  
 Ser Arg Arg Arg Thr Val Ala Cys Ser Ser Xaa Trp Thr Lys Trp Xaa  
     35                    40                    45  
 Ala Arg Pro Gln Gly Ser Val Glu Pro Gly Trp Pro Gly Pro Ser Gly  
     50                    55                    60  
 Thr Cys Thr Gly Xaa Tyr Arg Ser Phe Met Xaa Glu Asn Glu Arg Leu  
     65                    70                    75                    80

Arg Lys Glu Lys Ser Gln Leu Gln Asn Ser Arg Glu Leu Ala Gln Asn  
                     85                    90                    95  
 Glu Gln Arg Ile Leu Ala Gln Gln Val His Ala Leu Glu Xaa Arg Leu  
                     100                    105                    110  
 Leu Ser Ala Cys Tyr His His Gln Gln Gly Pro Gly Leu Thr Pro Pro  
                     115                    120                    125  
 Cys Pro Cys Leu Met Ala Pro Ala Pro Pro Cys His Ala Leu Pro Pro  
                     130                    135                    140  
 Leu Tyr Ser Cys Pro Cys Cys His Ile Cys Pro Leu Cys Xaa Val Pro  
                     145                    150                    155                    160  
 Leu Ala His Trp Xaa Xaa Leu Xaa Arg Gly Ala Pro Pro Cys Pro Ser  
                     165                    170                    175  
 Leu Ser Ser Gly Ala Leu Xaa Ser Gln Lys Xaa Thr Arg Arg Gly Phe  
                     180                    185                    190  
 Leu Val Leu Arg Xaa Val Phe Ser Xaa Arg Arg Gly Thr Val Glu Glu  
                     195                    200                    205  
 Pro Phe Cys Cys Lys Lys Lys Gly Thr Lys  
                     210                    215

<210> 20  
 <211> 373  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> unsure  
 <222> (50)

<220>  
 <221> unsure  
 <222> (54)

<220>  
 <221> unsure  
 <222> (60)

<220>  
 <221> unsure  
 <222> (102)

<220>  
 <221> unsure  
 <222> (114)

<220>  
 <221> unsure  
 <222> (118)

<220>  
 <221> unsure  
 <222> (144) .. (145)

<220>  
 <221> unsure  
 <222> (147)..(148)

<220>  
 <221> unsure  
 <222> (228)

<400> 20  
 ttccaaactt ggcccagaga ctggaggcct tcagagacca gattggcagn tccntgcgan 60  
 gtggccgcag ccagccaccc tgcagtgagg gcgcacggag cncaggccaa gtcntccntc 120  
 cccattgaag gccaaagtggg aacnnannag aatgctgtgt gacctcagac tgggctccac 180  
 actcttgggc ttcagtctgc ccactctgtg aatggagaca gcagctgnta ctccacctgc 240  
 agctgggcta gggcgggga ctgggggtgc tatttagggg aacaaggga tttcaggaga 300  
 aacccaggca gcaggggatg aaatacatga ataaagagag gcacagctc caaaaaaaaa 360  
 aaaaaaaaa aaa 373

<210> 21  
 <211> 1485  
 <212> DNA  
 <213> Homo sapiens

<400> 21  
 cccgcgcttc tctaaactaa ctatttaaag gtctgcggtc gcaaatgggt tgactaaacg 60  
 taggatggga cttaagttga acggcagata tatttcaactg atcctcgcgg tgcaaatagc 120  
 gtatctggtg caggccgtga gagcagcggg caagtgcgat gcggtcttca agggcttttc 180  
 ggactgtttg ctcaagctgg gcgacagcat ggccaactac ccgcagggcc tggacgacaa 240  
 gacgaacatc aagaccgtgt gcacatactg ggaggatttc cacagctgca cggtcacagc 300  
 ccttacggat tgccagggaag gggcgaaaga tatgtgggat aaactgagaa aagaatccaa 360  
 aaccccaac atccaaggca gcttattcga actctgcggc agcggcaacg gggcgggcggg 420  
 gtccctgctc ccggcggttc cgggtgctctc ggtgtctctc tcggcagctt tagcgacctg 480  
 gctttccctc tgagcgtggg gccagctccc cccgcgcgcc caccacact cactccatgc 540  
 tcccggaat cgagagggaag atccattagt tctttgggga cgttgtgatt ctctgtgatg 600  
 ctgaaaacac tcatatagga ttgtgggaaa tcctgattct cttttttatt tcgtttgatt 660  
 tcttgtgttt tatttgccaa atgttaccac tcagttagca agcaagcaca gccaaaatcg 720  
 gacctcagct ttagtcctgc ttcacacaca aataagaaaa cggcaaaccc accccatttt 780  
 ttaattttat tattattaat tttttttgtt ggcaaaagaa tctcaggaaac ggccctgggc 840  
 caccactat attaatcatg ctagttaacat gaaaaatgat gggctcctcc taatagggaag 900  
 gcgaggagag gagaaggcca ggggaatgaa ttcaagagag atgtccacgg acgaaacata 960  
 cgggtgaataa ttcacgctca cgtcgttctt ccacagtatc ttgttttgat catttccact 1020  
 gcacatttct cctcaagaaa agcgaaagga cagactgttg gctttgtgtt tggaggatag 1080  
 gagggagaga ggggaagggc tgaggaaatc tctggggtaa gagtaaggc ttcagaaga 1140  
 catgctgcta tggtcactga ggggttagct ttatctgctg ttgttgatgc atccgtccaa 1200  
 gttcactgcc tttattttcc ctccctccctc ttgttttagc tggtacacac acagtaatac 1260  
 ctgaatatcc aacggtatag atcacaaggg ggggatgtta aatgttaac taaaatatag 1320  
 ctaaaaaaaa attttgacat aaaagagcct tgatttttaa aaaaaagag agagagatgt 1380  
 aatttaaaaa gtttattata aattaaatc agcaaaaaaa gatttgctac aaagtataga 1440  
 gaagtataaa ataaagtta ttgtttgaaa aaaaaaaaa aaaaa 1485

<210> 22  
 <211> 142  
 <212> PRT  
 <213> Homo sapiens

<400> 22  
 Met Gly Leu Lys Leu Asn Gly Arg Tyr Ile Ser Leu Ile Leu Ala Val  
 1 5 10 15

Gln Ile Ala Tyr Leu Val Gln Ala Val Arg Ala Ala Gly Lys Cys Asp

20                      25                      30  
 Ala Val Phe Lys Gly Phe Ser Asp Cys Leu Leu Lys Leu Gly Asp Ser  
                     35                      40                      45  
 Met Ala Asn Tyr Pro Gln Gly Leu Asp Asp Lys Thr Asn Ile Lys Thr  
                     50                      55                      60  
 Val Cys Thr Tyr Trp Glu Asp Phe His Ser Cys Thr Val Thr Ala Leu  
                     65                      70                      75                      80  
 Thr Asp Cys Gln Glu Gly Ala Lys Asp Met Trp Asp Lys Leu Arg Lys  
                     85                      90                      95  
 Glu Ser Lys Asn Leu Asn Ile Gln Gly Ser Leu Phe Glu Leu Cys Gly  
                     100                      105                      110  
 Ser Gly Asn Gly Ala Ala Gly Ser Leu Leu Pro Ala Phe Pro Val Leu  
                     115                      120                      125  
 Leu Val Ser Leu Ser Ala Ala Leu Ala Thr Trp Leu Ser Phe  
                     130                      135                      140

<210> 23  
 <211> 825  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> unsure  
 <222> (290)

<220>  
 <221> unsure  
 <222> (480)

<220>  
 <221> unsure  
 <222> (491)

<220>  
 <221> unsure  
 <222> (589)

<220>  
 <221> unsure  
 <222> (698)

<400> 23  
 gaattcgcca cgagtttttt tttctgcag ttgtgtgtat gtgtgtttgt gtgaagaaaa 60  
 acagactctg tccaggtaga aatggtgagg agggggaaga gaattacatt tccaggtgca 120  
 gaaacttgcc aacagttttc ctakagtgcac tcagacacac cacagtaaca actctcgctg 180  
 caattttatt ttaatttgag aaataaagat ttctccaag ccacatgagg actctggcac 240  
 ccaccacaaa agcaagacct gtatttataa gccgaggggtg cagggagctn aactgcggga 300  
 cccgctcagg ccccggtggac ccacccccgt cccaccccc cctccaccg ytggggcccc 360  
 tcagtgtgtg ttggggggga tgcttgggca gctggggggg gagggagaca acaaacctyg 420  
 gggaaytggg agccagagct gcggcctgac tgacgccttt tgatgctcac gggaaatttn 480  
 tgcccaggat ntcagcccca ggctggttgt ttctacaaat ctctctcaaa tgtattattn 540  
 tgggtgacaaa aatgaaggag ctttgtaaat ttttttaaaa ttatgaatnc atatcaagta 600

```

gttggtttaca tttcttgaaa aaataggaac tcgggcagca gaatcagatt ggcagaatct 660
ttagactaca caggcaataa tcaagtctgc tgttttgncc tttcgtagta gaagtgggtg 720
tagtggttag atatctgttt ggtcttgctt cttgtattgc atttttttca ataaacaaca 780
acaaaaagaa aaaaaaaaaa aaaaaaaaaa aagatcttta attaa 825

```

<210> 24  
 <211> 200  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> UNSURE  
 <222> (75)

<220>  
 <221> UNSURE  
 <222> (86)

<220>  
 <221> UNSURE  
 <222> (89)

<220>  
 <221> UNSURE  
 <222> (118)

<220>  
 <221> UNSURE  
 <222> (122)

<220>  
 <221> UNSURE  
 <222> (134)

<220>  
 <221> UNSURE  
 <222> (152)

<220>  
 <221> UNSURE  
 <222> (158)

<220>  
 <221> UNSURE  
 <222> (199)

<400> 24  
 Met Arg Thr Leu Ala Pro Thr His Lys Ala Arg Pro Val Phe Ile Ser  
 1 5 10 15  
 Arg Gly Cys Arg Glu Leu Asn Cys Gly Thr Arg Gln Gly Pro Val Asp  
 20 25 30  
 Pro Ser Pro Ser Pro Pro Pro Pro Pro Leu Gly Pro Ile Ser Val  
 35 40 45  
 Cys Trp Gly Gly Cys Leu Gly Ser Trp Gly Val Arg Glu Thr Thr Asn  
 50 55 60  
 Leu Gly Glu Leu Gly Ala Arg Ala Ala Ala Xaa Leu Thr Pro Phe Asp

<220>  
<221> UNSURE



&lt;222&gt; (150)

&lt;400&gt; 26

Met Thr Glu Lys Ser Pro Phe Thr Thr Pro Ile Gly Arg Lys Asp Glu  
 1 5 10 15

Ala Asp Leu Ala Lys Ser Ala Leu Ala Met Ala Asp Ser Asp His Leu  
 20 25 30

Thr Ile Tyr Asn Ala Tyr Leu Gly Trp Lys Lys Ala Arg Gln Glu Gly  
 35 40 45

Gly Tyr Arg Ser Glu Ile Thr Tyr Cys Arg Arg Asn Phe Leu Asn Arg  
 50 55 60

Thr Ser Leu Leu Thr Leu Glu Asp Val Lys Gln Glu Leu Ile Lys Leu  
 65 70 75 80

Val Lys Ala Ala Gly Phe Ser Ser Ser Thr Thr Ser Thr Ser Trp Glu  
 85 90 95

Gly Asn Arg Ala Ser Gln Thr Leu Ser Phe Gln Glu Ile Ala Leu Leu  
 100 105 110

Lys Ala Val Leu Val Ala Gly Leu Tyr Asp Asn Val Gly Lys Ile Ile  
 115 120 125

Tyr Thr Lys Ser Val Asp Val Thr Glu Lys Leu Ala Cys Ile Val Glu  
 130 135 140

Thr Ala Gln Gly Lys Xaa Gln Val His Pro Ser Ser Val Asn Arg Asp  
 145 150 155 160

Leu Gln Thr His Gly Trp Leu Leu Tyr Gln Glu Lys Ile Arg Tyr Ala  
 165 170 175

Arg Val Tyr Leu Arg Glu Thr Thr Leu Ile Thr Pro Phe Pro Phe Leu  
 180 185 190

Leu Phe Gly Gly Asp Ile Glu Val Gln His Arg Glu Arg Leu Leu Ser  
 195 200 205

Ile Asp Gly Trp Ile Tyr Phe Gln Ala Pro Val Lys Ile Ala Val Ile  
 210 215 220

Phe Lys Gln Leu Arg Val Leu Ile Asp Ser Val Leu Arg Lys Lys Leu  
 225 230 235 240

Glu Asn Pro Lys Met Ser Leu Glu Asn Asp Lys Ile Leu Gln Ile Ile  
 245 250 255

Thr Glu Leu Ile Lys Thr Glu Asn Asn  
 260 265

&lt;210&gt; 27

&lt;211&gt; 423

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

<220>  
 <221> unsure  
 <222> (13)..(14)

<220>  
 <221> unsure  
 <222> (18)

<220>  
 <221> unsure  
 <222> (412)

<400> 27  
 gaattcggca cgnngtgnaa tataaaaatt tatttttaag tcaaagtatg caacaaataa 60  
 acctacagaa aacattttcc catcacaaac tgggtgctta ccaaataata ttttgaaaac 120  
 acattccttc agtcattata aagttcttaa aatacaaaaag aaattaaaac tgtaagaaaag 180  
 tctagtagac cagatgctgt tgtcaagact tgtatgttgg tgtttttgct ttcagtacat 240  
 cccacgccat ccacctccac tycatgccgc cttgcccata gtaacctcca ctgcctccac 300  
 caccacggcc ataaccaccc aaaccatcag gattaccata tcctccactg taattgttcc 360  
 ccattcccat tcttccaact ggattccata ggccytccct ggattatttt tnaaaaggaa 420  
 aaa 423

<210> 28  
 <211> 76  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> UNSURE  
 <222> (66)

<220>  
 <221> UNSURE  
 <222> (68)

<220>  
 <221> UNSURE  
 <222> (73)

<400> 28  
 Met Leu Leu Ser Arg Leu Val Cys Trp Cys Phe Cys Phe Gln Tyr Ile  
 1 5 10 15  
 Pro Arg His Pro Pro Pro Leu His Ala Ala Leu Pro Ile Val Thr Ser  
 20 25 30  
 Thr Ala Ser Thr Thr Thr Ala Ile Thr Thr Gln Thr Ile Arg Ser Thr  
 35 40 45  
 Ile Ser Ser Thr Val Ile Val Pro His Ser His Ser Ser Asn Trp Ile  
 50 55 60  
 Pro Xaa Ala Xaa Pro Gly Leu Phe Xaa Lys Arg Lys  
 65 70 75

<210> 29  
 <211> 294  
 <212> DNA  
 <213> Homo sapiens

<220>  
<221> unsure  
<222> (20)

<220>  
<221> unsure  
<222> (23)

<220>  
<221> unsure  
<222> (28)

<220>  
<221> unsure  
<222> (46)

<220>  
<221> unsure  
<222> (64)

<220>  
<221> unsure  
<222> (66)

<220>  
<221> unsure  
<222> (69)

<220>  
<221> unsure  
<222> (86)

<220>  
<221> unsure  
<222> (92)

<220>  
<221> unsure  
<222> (101)

<220>  
<221> unsure  
<222> (103)

<220>  
<221> unsure  
<222> (107)

<220>  
<221> unsure  
<222> (185)

<220>  
<221> unsure  
<222> (208)

<220>  
<221> unsure  
<222> (214)

&lt;400&gt; 29

taaaaacccc tttttcctcn tangggnta tcatagggtc cgggtngctg tcccagcaat 60  
 tttntngngn gatcataaaa tccttngatt tnactcgtga nanttgngaa gatctcaata 120  
 tacctattta aaaatgtttt aagggtacagg tttcagcata aatgtattag tgtaaattag 180  
 atacngggca aaatgcagta agtttttnta tatntagata cataacccaa tttaaattgc 240  
 ctaaatacac cgtaagttaa cagtttaaac ctacaaactt aattaagcgg ccgc 294

&lt;210&gt; 30

&lt;211&gt; 514

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (462)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (467)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (481)

&lt;400&gt; 30

aagcttggca cgtggctgat tggagctgta aaacatcatc aggtgttgct attwttttat 60  
 atgattattc tgttacttgt atttattggt cagttttctg tatcttgcgc ttgttttagcc 120  
 ctgaaccagg agcaacaggg tcagcttctg gaggttggtt ggaacaatac ggcaagtgc 180  
 cgaaatgaca tccagagaaa tctaaactgc tgtgggttcc gaagtgttaa cccaaatgac 240  
 acctgtctgg ctagtctgtg taaaagtac cactcgtgct cgccatgtgc tccaatcata 300  
 ggagaatatg ctggagaggt tttagattt gttgggtggca ttggcctgtt cttcagtttt 360  
 acagagatcc tgggggtgtt ggctgacct cagatacagg aaccagaaag acccccgcgc 420  
 gaatcctagt gcattccttt ggatgaggaa aacaaggga gnttcnntt cgtattatgg 480  
 ncttgtttca ctttctgtaa tttttctgtt aagg 514

&lt;210&gt; 31

&lt;211&gt; 151

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; UNSURE

&lt;222&gt; (123)

&lt;220&gt;

&lt;221&gt; UNSURE

&lt;222&gt; (134)

&lt;220&gt;

&lt;221&gt; UNSURE

&lt;222&gt; (136)

&lt;220&gt;

&lt;221&gt; UNSURE

&lt;222&gt; (141)

&lt;400&gt; 31

Met Ile Ile Leu Leu Leu Val Phe Ile Val Gln Phe Ser Val Ser Cys

1

5

10

15

Ala Cys Leu Ala Leu Asn Gln Glu Gln Gln Gly Gln Leu Leu Glu Val  
 20 25 30

Gly Trp Asn Asn Thr Ala Ser Ala Arg Asn Asp Ile Gln Arg Asn Leu  
 35 40 45

Asn Cys Cys Gly Phe Arg Ser Val Asn Pro Asn Asp Thr Cys Leu Ala  
 50 55 60

Ser Cys Val Lys Ser Asp His Ser Cys Ser Pro Cys Ala Pro Ile Ile  
 65 70 75 80

Gly Glu Tyr Ala Gly Glu Val Leu Arg Phe Val Gly Gly Ile Gly Leu  
 85 90 95

Phe Phe Ser Phe Thr Glu Ile Leu Gly Cys Leu Ala Asp Leu Gln Ile  
 100 105 110

Gln Glu Pro Glu Arg Pro Pro Arg Glu Ser Xaa Cys Ile Pro Leu Asp  
 115 120 125

Glu Glu Asn Lys Gly Xaa Phe Xaa Phe Val Leu Trp Xaa Cys Phe Thr  
 130 135 140

Phe Cys Asn Phe Ser Val Lys  
 145 150

<210> 32

<211> 204

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> (53)

<220>

<221> unsure

<222> (106)

<220>

<221> unsure

<222> (111)

<220>

<221> unsure

<222> (128)

<400> 32

acgtagcaaa aagatatttg attatcttaa aaattgtaa ataccgtttt cangaaagtt 60  
 ctcagtattg taacagcaac ttgtcaaac taagcatatt tgaatntgat ntcccataat 120  
 ttgaaatnga aatcgtatgg tgtggctctg tatattctgt taaaaaatta agggaccaga 180  
 aaccttaaaa aaaaaaaaaa aaaa 204

<210> 33

<211> 391

<212> DNA

<213> Homo sapiens

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (5)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (20)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (336)

&lt;400&gt; 33

```

gaacntgggc cgcattgtatn tcttctatgg caacaagacc tcggtgcagt tccagaattt 60
ctcaccacct gtggttcacc cgggagacct ccagactcag ctggctgtgc agaccaagcg 120
cgtggcggcg caggtggacg gcggcgcgca ggtgcagcag gtgctcaata tcgagtgcct 180
gcgggacttc ctgacgcccc cgctgctgtc cgtgcgcttc cggtagcgtg gcgcccccca 240
ggccctcacc ctgaagctcc cagtgacct caacaagttc ttccagccca ccgagatggc 300
ggcccaggat ttcttccagc gctggaagca gctgancctc cctcaacagg aggcgcagaa 360
aatcttcaaa gcccaaccacc ccatggacgc a 391

```

&lt;210&gt; 34

&lt;211&gt; 126

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; UNSURE

&lt;222&gt; (3)

&lt;220&gt;

&lt;221&gt; UNSURE

&lt;222&gt; (108)

&lt;400&gt; 34

```

Met Tyr Xaa Phe Tyr Gly Asn Lys Thr Ser Val Gln Phe Gln Asn Phe
  1             5             10             15

Ser Pro Thr Val Val His Pro Gly Asp Leu Gln Thr Gln Leu Ala Val
          20             25             30

Gln Thr Lys Arg Val Ala Ala Gln Val Asp Gly Gly Ala Gln Val Gln
  35             40             45

Gln Val Leu Asn Ile Glu Cys Leu Arg Asp Phe Leu Thr Pro Pro Leu
  50             55             60

Leu Ser Val Arg Phe Arg Tyr Gly Gly Ala Pro Gln Ala Leu Thr Leu
  65             70             75             80

Lys Leu Pro Val Thr Ile Asn Lys Phe Phe Gln Pro Thr Glu Met Ala
          85             90             95

Ala Gln Asp Phe Phe Gln Arg Trp Lys Gln Leu Xaa Leu Pro Gln Gln
          100             105             110

Glu Ala Gln Lys Ile Phe Lys Ala Asn His Pro Met Asp Ala
          115             120             125

```

<210> 35  
<211> 177  
<212> DNA  
<213> *Homo sapiens*

<220>  
<221> unsure  
<222> (7)

<220>  
<221> unsure  
<222> (11)

<220>  
<221> unsure  
<222> (66)

<220>  
<221> unsure  
<222> (135)

<400> 35  
ccctctntcc ntttcccccc caagcacaga ggggagaggg gccagggaag tggatgtttc 60  
ttccctntccc accccaccct gttgtagecc ctcttaccct ctccccatcc aggggctgtg 120  
tattattgtg agcgnataaa cagagagacg ctaaaaaaaa aaaaaaaaaa aaaaaaa 177

<210> 36  
<211> 655  
<212> DNA  
<213> *Homo sapiens*

<220>  
<221> unsure  
<222> (5)

<220>  
<221> unsure  
<222> (27)

<220>  
<221> unsure  
<222> (29)..(30)

<220>  
<221> unsure  
<222> (153)

<220>  
<221> unsure  
<222> (431)

<220>  
<221> unsure  
<222> (610)

<220>  
<221> unsure  
<222> (638)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (655)

&lt;400&gt; 36

```

caatnataaa atgtcagctt ttaaggnann cctgtggaat atattttcca gcaataaaaa 60
gagatccagg cagatattta catagtgttc cctgaatctg tgaaaaaatg gcttcgacag 120
ctaaagaatg ctgggaaaat tcttctgtta atnaccagtt ctcacagtga ttactgtaga 180
cttctctgcg aatatattct tgggaatgat ttacagacc tttttgacat tgtgattaca 240
aatgcattga agcctgggtt cttctccac ttaccaagtc agagaccttt ccggacactc 300
gagaatgatg aggagcagga ggcactgcc aactctggata aacctggctg gtactcccaa 360
gggaacgctg tccaccteta tgaacttctg aagaaaatga ctggcaaacc tgaacccaag 420
gttstttatt nwtggtgwca gcatgcawtc agatattttc ccagctcgtc actatagtaa 480
ttggggagac agtctctatc cgkgaagga actcagaggg ggatgaargg gcacgagggg 540
gttcagaggg cttgagggag ttcagagcct cttagaagaa ggaaagggaa attttgaggg 600
gacaaaaagn caaaccttt aattatttca ttttaaanat ggggggtttt ttttn 655

```

&lt;210&gt; 37

&lt;211&gt; 199

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; UNSURE

&lt;222&gt; (32)

&lt;220&gt;

&lt;221&gt; UNSURE

&lt;222&gt; (123)

&lt;220&gt;

&lt;221&gt; UNSURE

&lt;222&gt; (125)

&lt;220&gt;

&lt;221&gt; UNSURE

&lt;222&gt; (127)

&lt;220&gt;

&lt;221&gt; UNSURE

&lt;222&gt; (131)

&lt;220&gt;

&lt;221&gt; UNSURE

&lt;222&gt; (140)..(141)

&lt;220&gt;

&lt;221&gt; UNSURE

&lt;222&gt; (149)

&lt;220&gt;

&lt;221&gt; UNSURE

&lt;222&gt; (166)

&lt;220&gt;

&lt;221&gt; UNSURE

&lt;222&gt; (180)

&lt;220&gt;

&lt;221&gt; UNSURE



&lt;222&gt; (185)

&lt;220&gt;

&lt;221&gt; UNSURE

&lt;222&gt; (193)..(194)

&lt;400&gt; 37

Lys Glu Ile Gln Ala Asp Ile Tyr Ile Val Val Pro Glu Ser Val Lys  
 1 5 10 15

Lys Trp Leu Arg Gln Leu Lys Asn Ala Gly Lys Ile Leu Leu Leu Xaa  
 20 25 30

Thr Ser Ser His Ser Asp Tyr Cys Arg Leu Leu Cys Glu Tyr Ile Leu  
 35 40 45

Gly Asn Asp Phe Thr Asp Leu Phe Asp Ile Val Ile Thr Asn Ala Leu  
 50 55 60

Lys Pro Gly Phe Phe Ser His Leu Pro Ser Gln Arg Pro Phe Arg Thr  
 65 70 75 80

Leu Glu Asn Asp Glu Glu Gln Glu Ala Leu Pro Ser Leu Asp Lys Pro  
 85 90 95

Gly Trp Tyr Ser Gln Gly Asn Ala Val His Leu Tyr Glu Leu Leu Lys  
 100 105 110

Lys Met Thr Gly Lys Pro Glu Pro Lys Val Xaa Tyr Xaa Trp Xaa Gln  
 115 120 125

His Ala Xaa Arg Tyr Phe Pro Ser Ser Ser Leu Xaa Xaa Leu Gly Arg  
 130 135 140

Gln Ser Ser Ser Xaa Glu Gly Thr Gln Arg Gly Met Lys Gly His Glu  
 145 150 155 160

Gly Val Gln Arg Pro Xaa Gly Ser Ser Glu Pro Leu Arg Arg Arg Lys  
 165 170 175

Gly Lys Phe Xaa Gly Asp Gln Lys Xaa Lys Pro Leu Ile Ile Ser Phe  
 180 185 190

Xaa Xaa Trp Gly Phe Phe Phe  
 195

&lt;210&gt; 38

&lt;211&gt; 265

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (11)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (47)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (49)

&lt;400&gt; 38

```

tcctccactg ntcttatcaa gtgatgagac actgatatcc aaataantng tatttactga 60
aaaatgaagt gaagacccat atatgcagtt aaaaaaaagt taattttcaa aaaatactgt 120
aaaagacttt aaggaacaag ttttattgac caataagttg atattttgtcc ataggtctcc 180
tttctataaa tcattttgat gtttaacaac ttttattata ttaaaatctc agtatccctaa 240
aacttaaaaa aaaaaaaaaa aaaaaa                                     265

```

&lt;210&gt; 39

&lt;211&gt; 377

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 39

```

gaattcggca cgaggattct catcagcttt tcctgggttt gcccaattaca tccgagctgg 60
gactctaatc atggctctgc atgactcttc cgattacctg ctggaktcag ccaagatggt 120
taactacgcg ggatggaaga acacctgcaa caacatcttc atcgtcttcg ccattgtttt 180
tatcatcacc cgactgggtca tcctgccctt ctggatccctg cattgcaccc tgggtgtacc 240
cactgggagct ctatcctgcc ttctttgggc tattactttct ttcaattcca tgatgggagt 300
tctacagctg ctgcatatct tctgggsecta cctcattttg cgsatgggcc cacaagttca 360
taactgggaa agctggtt                                     377

```

&lt;210&gt; 40

&lt;211&gt; 102

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; UNSURE

&lt;222&gt; (12)

&lt;220&gt;

&lt;221&gt; UNSURE

&lt;222&gt; (86)

&lt;400&gt; 40

```

Met Ala Leu His Asp Ser Ser Asp Tyr Leu Leu Xaa Ser Ala Lys Met
  1             5             10             15

```

```

Phe Asn Tyr Ala Gly Trp Lys Asn Thr Cys Asn Asn Ile Phe Ile Val
      20             25             30

```

```

Phe Ala Ile Val Phe Ile Ile Thr Arg Leu Val Ile Leu Pro Phe Trp
      35             40             45

```

```

Ile Leu His Cys Thr Leu Gly Val Pro Thr Gly Ala Leu Ser Cys Leu
      50             55             60

```

```

Leu Trp Ala Ile Thr Ser Phe Asn Ser Met Met Gly Val Leu Gln Leu
      65             70             75             80

```

```

Leu His Ile Phe Trp Xaa Tyr Leu Ile Leu Arg Met Gly Pro Gln Val
      85             90             95

```

```

His Asn Trp Glu Ser Trp
      100

```

<210> 41  
<211> 359  
<212> DNA  
<213> Homo sapiens

<220>  
<221> unsure  
<222> (38)

<220>  
<221> unsure  
<222> (42)

<220>  
<221> unsure  
<222> (49)

<220>  
<221> unsure  
<222> (82)

<220>  
<221> unsure  
<222> (106)

<220>  
<221> unsure  
<222> (111)

<220>  
<221> unsure  
<222> (125)

<220>  
<221> unsure  
<222> (127)

<220>  
<221> unsure  
<222> (138)

<220>  
<221> unsure  
<222> (175)

<220>  
<221> unsure  
<222> (229)

<220>  
<221> unsure  
<222> (246)

<220>  
<221> unsure  
<222> (263)

<220>  
<221> unsure

<222> (268)

<220>

<221> unsure

<222> (273)

<220>

<221> unsure

<222> (276)

<220>

<221> unsure

<222> (278)

<220>

<221> unsure

<222> (281)

<220>

<221> unsure

<222> (302)

<220>

<221> unsure

<222> (314)..(316)

<400> 41

```

aaaaagtggg ggctgtactg gggactgctc ggatgatntt tnttagtgnt actttttttca 60
gctgtccctg tagcgacagg tntaagatct gactgcctcc ttttntggc ntcttcccc 120
ttccntnttc tcttcagnta ggctagctgg tttggagtag aatggcaact aattntaatt 180
tttatttatt aaatatttgg ggttttgggt ttaaagccag aattacggnt agcacctagc 240
atttengcag agggaccatt ttngaccnaa atntantntt natgggtttt tttttaaaat 300
tnaaagatta aatnnnaaat attaaataaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 359

```

<210> 42

<211> 332

<212> DNA

<213> Homo sapiens

<400> 42

```

gaattcggca cgagcttgat tgctccaggg cccacaacgg cagtgtccta catgtcgggtg 60
aaatgtgtgg atgcccgtaa gaaccatcac aagacaaaat gggtcgtgcc ttggggaccc 120
aatcattgtg acaagatccg agacattgaa gaggcaattc caagggaaat tgaagccaat 180
gacatcgtgt tttctgttca cattccccct cccacatgg gagatgagtc cttggttcca 240
attcatgmtg tttatcctgg cagctgggac attgcctttc aagctaaaaca accaaatcag 300
gggaaaatgc aggaagtctc catgggacgt tt 332

```

<210> 43

<211> 110

<212> PRT

<213> Homo sapiens

<220>

<221> UNSURE

<222> (83)

<400> 43

```

Glu Phe Gly Thr Ser Leu Ile Ala Pro Gly Pro Thr Thr Ala Val Ser
  1             5             10            15

```

Tyr Met Ser Val Lys Cys Val Asp Ala Arg Lys Asn His His Lys Thr  
                   20                  25                  30  
 Lys Trp Phe Val Pro Trp Gly Pro Asn His Cys Asp Lys Ile Arg Asp  
                   35                  40                  45  
 Ile Glu Glu Ala Ile Pro Arg Glu Ile Glu Ala Asn Asp Ile Val Phe  
                   50                  55                  60  
 Ser Val His Ile Pro Leu Pro His Met Gly Asp Glu Ser Leu Val Pro  
                   65                  70                  75                  80  
 Ile His Xaa Val Tyr Pro Gly Ser Trp Asp Ile Ala Phe Gln Ala Lys  
                   85                  90                  95  
 Gln Pro Asn Gln Gly Lys Met Gln Glu Val Ser Met Gly Arg  
                   100                  105                  110

<210> 44  
 <211> 314  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> unsure  
 <222> (78)..(79)

<220>  
 <221> unsure  
 <222> (85)..(86)

<220>  
 <221> unsure  
 <222> (126)

<400> 44  
 tcactcctaa tccatgacca ctgttttttt cctattttata tcaccaggta gcctactgag 60  
 ttaatatatta agttgtcnnt gggtnngtgt ccctgttttg tggcataata taactgaatt 120  
 tcatgngaag atttattcca ccagggttat ttcagctttg aaaccaaate tgtgtatcta 180  
 atactaacca atctgttga tgtggatttt aaaaaatgtt tgctaaacta cccaagtaag 240  
 atttactgta ttaaattgcc ttcgggtctg aaaagctttt ttaaaaaaaaa aaaaaaaaaa 300  
 aaaaaaaaaa aaaa 314

<210> 45  
 <211> 1089  
 <212> DNA  
 <213> Homo sapiens

<400> 45  
 gggctactac gatggcgatg agtttcgagt ggccgtggca gtatcgcttc ccacccttct 60  
 ttacgttaca accgaatgtg gacactcggc agaagcagct ggccgccttg tgctcgctgg 120  
 tcctgtcctt ctgcccctg cacaacacgt ccagcatgac ggtgatggaa gctcaggaga 180  
 gcccgtctct caacaacgtc aagctacagc gaaagcttcc tgtggagtcg atccagattg 240  
 tattagagga actgaggaag aaagggaacc tcgagtgggt ggataagagc aagtcagct 300  
 tcctgatcat gtggcgagg ccagaagaat gggggaaact catctatcag tgggtttcca 360  
 ggagtggcca gaacaactcc gtctttaccc tgtatgaact gactaatggg gaagacacag 420  
 aggatgagga gttccacggg ctggatgaag ccactctact gcgggctctg caggccctac 480  
 agcaggagca caaggccgag atcatcactg tcagcgatgg ccgaggcgtc aagttcttct 540  
 agcaggggacc tgtctcctt tactttctac ctcccacctt tccagggtt tcaaaggag 600

acagacccag tgtcccccaa agactggatc tgtgactcca ccagactcaa aaggactcca 660  
 gtccctgaagg ctgggacctg gggatgggtt tctcacaccc catatgtctg tcccttggat 720  
 aggggtgaggc tgaagcacca gggagaaaat atgtgcttct tctcgcccta cctcctttcc 780  
 catcctagac tgtccttgag ccagggtctg taaacctgac acttttatatg tgttcacaca 840  
 tgtaagtaca tacacacatg cgcctgcagc acatgcttct gtctcctcct cctcccaccc 900  
 ctttagctgc tgttgctcc cttctcaggc tgggtgctga tccttcctag gggatggggg 960  
 aagccctggc tgcaggcagc cttccaggca atatgaagat aggaggccca cgggcctggc 1020  
 agtgagaggt gtggcccccac accgatttat gatattaaaa tctcaactcc caaaaaaaaa 1080  
 aaaaaaaaaa 1089

&lt;210&gt; 46

&lt;211&gt; 176

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 46

Met Ala Met Ser Phe Glu Trp Pro Trp Gln Tyr Arg Phe Pro Pro Phe  
 1 5 10 15

Phe Thr Leu Gln Pro Asn Val Asp Thr Arg Gln Lys Gln Leu Ala Ala  
 20 25 30

Trp Cys Ser Leu Val Leu Ser Phe Cys Arg Leu His Lys Gln Ser Ser  
 35 40 45

Met Thr Val Met Glu Ala Gln Glu Ser Pro Leu Phe Asn Asn Val Lys  
 50 55 60

Leu Gln Arg Lys Leu Pro Val Glu Ser Ile Gln Ile Val Leu Glu Glu  
 65 70 75 80

Leu Arg Lys Lys Gly Asn Leu Glu Trp Leu Asp Lys Ser Lys Ser Ser  
 85 90 95

Phe Leu Ile Met Trp Arg Arg Pro Glu Trp Gly Lys Leu Ile Tyr  
 100 105 110

Gln Trp Val Ser Arg Ser Gly Gln Asn Asn Ser Val Phe Thr Leu Tyr  
 115 120 125

Glu Leu Thr Asn Gly Glu Asp Thr Glu Asp Glu Glu Phe His Gly Leu  
 130 135 140

Asp Glu Ala Thr Leu Leu Arg Ala Leu Gln Ala Leu Gln Gln Glu His  
 145 150 155 160

Lys Ala Glu Ile Ile Thr Val Ser Asp Gly Arg Gly Val Lys Phe Phe  
 165 170 175

&lt;210&gt; 47

&lt;211&gt; 632

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 47

agcttcggaa taataatttt ggcaaatcta tcttctgaac cactcatttc tgtggtctta 60  
 atggctccaa tttggggacc aataatgttc attgtctcag gatccctgtc aattgcagca 120  
 ggagtgaac ctacaaaaag cctgatcatc agcagtctaa ctctgaacac tatcacctct 180  
 gtgttggtc caactgcaag cataatgggt gtatgcagtg tggctgtggg ttcacagttt 240

ccgtttcggg ataattatac aatcaccaag ggtttggata ttttgatgtt aattttaaat 300  
 atgctagaat tctgcattgc tgtgtccatc tctgcttttg gatgtaaagc ttctgttgt 360  
 aactccagcg aggttcttgt agtgctacca tcaaattcctg ctgtgactgt gatggcacc 420  
 cccacaccac ttaatgaagg tttagggcca ccaaaagatc aacagacaaa tgctccagaa 480  
 atctatgctg actgtgacac aagaagcctc acatgaagaa attaccagta tccaacttcg 540  
 atactgatag acttgttgat attattatta tatgtaatcc aattatgaac tgtgtgtgta 600  
 tagagagata ataaattcaa aattatgttc tc 632

<210> 48

<211> 151

<212> PRT

<213> Homo sapiens

<400> 48

Met Ala Pro Ile Trp Gly Pro Ile Met Phe Ile Val Ser Gly Ser Leu  
 1 5 10 15

Ser Ile Ala Ala Gly Val Lys Pro Thr Lys Ser Leu Ile Ile Ser Ser  
 20 25 30

Leu Thr Leu Asn Thr Ile Thr Ser Val Leu Ala Ala Thr Ala Ser Ile  
 35 40 45

Met Gly Val Val Ser Val Ala Val Gly Ser Gln Phe Pro Phe Arg Tyr  
 50 55 60

Asn Tyr Thr Ile Thr Lys Gly Leu Asp Ile Leu Met Leu Ile Leu Asn  
 65 70 75 80

Met Leu Glu Phe Cys Ile Ala Val Ser Ile Ser Ala Phe Gly Cys Lys  
 85 90 95

Ala Ser Cys Cys Asn Ser Ser Glu Val Leu Val Val Leu Pro Ser Asn  
 100 105 110

Pro Ala Val Thr Val Met Ala Pro Pro Thr Pro Leu Asn Glu Gly Leu  
 115 120 125

Arg Pro Pro Lys Asp Gln Gln Thr Asn Ala Pro Glu Ile Tyr Ala Asp  
 130 135 140

Cys Asp Thr Arg Ser Leu Thr  
 145 150

<210> 49

<211> 365

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> (18)

<220>

<221> unsure

<222> (25)

<220>

<221> unsure

<222> (75)

<220>

<221> unsure

<222> (78)

<220>

<221> unsure

<222> (134)

<220>

<221> unsure

<222> (137)

<220>

<221> unsure

<222> (141)

<220>

<221> unsure

<222> (144)

<220>

<221> unsure

<222> (206)

<220>

<221> unsure

<222> (215)

<220>

<221> unsure

<222> (238)

<220>

<221> unsure

<222> (241)

<220>

<221> unsure

<222> (260)

<400> 49

```
ctatggggac caaagtgnnt tttcnttcag gaagtggaga tgcattggcca tctccccctc 60
ccttttttct tctcntgnnt ttctttcccc atagaaagta ccttggaagta gcacagtcgg 120
tccttgcatt tgcncgngct ntcttttgag taaaagtata catggagtaa aaatcatatt 180
aagcatcaga ttcaacttat atttnttatt tcatnttctt cctttccctt ctcccacntt 240
ntactgggca taattatatn ttaatcatat atggaaatgt gcaacatatg gtatttggtta 300
aatacgtttg tttttattgc agagcaaaaa taaatcaaat tagaagcaaa aaaaaaaaaa 360
aaaaa 365
```

<210> 50

<211> 689

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> (5)



<220>  
 <221> unsure  
 <222> (10)

<220>  
 <221> unsure  
 <222> (413)

<400> 50  
 cccanagagn cctaggaaga tgaacaaacg acagctctac taccagggtt taaactttgc 60  
 catgatcgtg tcttctgctg tcatgatctg gaaaggcctg attgttctca cgggcagcga 120  
 gagtcccatc gtggwgtac tcagtggcag tatggagccg gccttcacac gaggagatct 180  
 bctgttcctc acgaatttcc gggaggaccc catcagagct ggtgaaatag ttgtttttaa 240  
 ggttgaagga agagacattc cgatagttca cagagtaatc aagggttcag aaaaagataa 300  
 tgggtgacatc aartttctga ctaaaggaga taataatgaa gtygatgata gaggcttgta 360  
 caaagaaggc cagaactggc tggaaaagaa ggacgtggtg ggaagagcaa ganggttttt 420  
 accatatggt ggtatggtca ccataataat gaatgactat ccaaaattca aktatgctct 480  
 ttgtgctgta atgggtgcat atgtgttact aaaacgtgaa tcctaaaatg agaagcagtt 540  
 cctgggacca gattgaaatg aattctgttg aaaaagagaa aaactaatat atttgagatg 600  
 ttccattttc tgtataaaaag ggaacagtgt ggagatgttt ttgtctgtc caaataaaaag 660  
 attcatcagt aaaaaaaaaa aaaaaaaaaa 689

<210> 51  
 <211> 168  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> UNSURE  
 <222> (39)

<220>  
 <221> UNSURE  
 <222> (132)

<220>  
 <221> UNSURE  
 <222> (151)

<400> 51  
 Met Asn Lys Arg Gln Leu Tyr Tyr Gln Val Leu Asn Phe Ala Met Ile  
 1 5 10 15  
 Val Ser Ser Ala Leu Met Ile Trp Lys Gly Leu Ile Val Leu Thr Gly  
 20 25 30  
 Ser Glu Ser Pro Ile Val Xaa Val Leu Ser Gly Ser Met Glu Pro Ala  
 35 40 45  
 Phe His Arg Gly Asp Leu Leu Phe Leu Thr Asn Phe Arg Glu Asp Pro  
 50 55 60  
 Ile Arg Ala Gly Glu Ile Val Val Phe Lys Val Glu Gly Arg Asp Ile  
 65 70 75 80  
 Pro Ile Val His Arg Val Ile Lys Val His Glu Lys Asp Asn Gly Asp  
 85 90 95  
 Ile Lys Phe Leu Thr Lys Gly Asp Asn Asn Glu Val Asp Asp Arg Gly  
 100 105 110

Leu Tyr Lys Glu Gly Gln Asn Trp Leu Glu Lys Lys Asp Val Val Gly  
 115 120 125

Arg Ala Arg Xaa Phe Leu Pro Tyr Val Gly Met Val Thr Ile Ile Met  
 130 135 140

Asn Asp Tyr Pro Lys Phe Xaa Tyr Ala Leu Leu Ala Val Met Gly Ala  
 145 150 155 160

Tyr Val Leu Leu Lys Arg Glu Ser  
 165

<210> 52

<211> 309

<212> DNA

<213> Homo sapiens

<400> 52

ctctccccc cccctctctc tctctctcgc atactaacta ggtttgactg tattactcgt 60  
 accagattta aaattagact agccttgcca caacgcccta ctgagaggta ctgtcgaact 120  
 gtagacagca tgatgttctt tgatggtgaa agtctaaatc tggaccgtgt tcagagatac 180  
 caaatgatga ggctgaaaag gggaaagggg gttcttcagt ctcttcttct tcttcttttt 240  
 attttttttt ccatgatgtt ttctctatgg ccagtgc aaa tgggtgtgtc acccttgcat 300  
 gttgccaac 309

<210> 53

<211> 60

<212> PRT

<213> Homo sapiens

<400> 53

Met Met Phe Phe Asp Gly Glu Ser Leu Asn Leu Asp Arg Val Gln Arg  
 1 5 10 15

Tyr Gln Met Met Arg Leu Lys Arg Gly Lys Gly Val Leu Gln Ser Leu  
 20 25 30

Leu Leu Leu Leu Phe Ile Phe Phe Ser Met Met Phe Ser Leu Trp Pro  
 35 40 45

Val Gln Met Val Leu Ser Pro Leu His Val Ala Asn  
 50 55 60

<210> 54

<211> 257

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> (97)

<220>

<221> unsure

<222> (170)

<220>

&lt;221&gt; unsure

&lt;222&gt; (222)

&lt;400&gt; 54

```

aggctctctt ggttctttct atatcatcat tttattatta tgtcctaata taaagtactg 60
gctcataggg ccagggtatt attatagaat attattntcg catgtaaaca aagatatctt 120
tgctttaaga tgtgagaaga aatgaattta ctttgtttgc attaagttan ggaagagttg 180
taatatatac tttagaaaag aagagaagaa aactagtatc tntaagcggc aaaaaaaaaa 240
aaaaaaaaaa aaaaaaa                                     257

```

&lt;210&gt; 55

&lt;211&gt; 467

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (32)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (84)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (87)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (89)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (96)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (149)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (246)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (248)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (250)..(251)

&lt;400&gt; 55

```

cacgaggatt gatttccatc ttgcctctcc anaaggcaaa accttagttt ttgaacaaag 60
aaaatcagat ggagttcaca ctgntanana ctgaanttgg tgattacatg ttctgctttg 120
acaatacatt cagcaccatt tctgagaang tgattttctt tgaattaatc ctggataata 180
tgggagaaca ggcacaagaa caagaagatt ggaagaaata tattactggc acagatatat 240
tggaatntnan nctggaagac atcctggaat ccatcaacag catcaagtcc agactaagca 300
aaagtgggca catacaaact ctgcttagag catttgaagc tcgtgatcga aacatacaag 360
aaagcaactt tgatagagtc aatttctggt ctatggttaa tttagtggtc atgggtgggtg 420
gttcagccat tcaagtttat atgctgaaga gtctgtttga agataag 467

```

<210> 56  
 <211> 133  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> UNSURE  
 <222> (6)..(7)

<220>  
 <221> UNSURE  
 <222> (10)

<220>  
 <221> UNSURE  
 <222> (27)

<220>  
 <221> UNSURE  
 <222> (60)..(61)

<400> 56  
 Met Glu Phe Thr Leu Xaa Xaa Thr Glu Xaa Gly Asp Tyr Met Phe Cys  
           1                  5                  10                  15  
 Phe Asp Asn Thr Phe Ser Thr Ile Ser Glu Xaa Val Ile Phe Phe Glu  
                   20                  25                  30  
 Leu Ile Leu Asp Asn Met Gly Glu Gln Ala Gln Glu Gln Glu Asp Trp  
           35                  40                  45  
 Lys Lys Tyr Ile Thr Gly Thr Asp Ile Leu Asp Xaa Xaa Leu Glu Asp  
           50                  55                  60  
 Ile Leu Glu Ser Ile Asn Ser Ile Lys Ser Arg Leu Ser Lys Ser Gly  
           65                  70                  75                  80  
 His Ile Gln Thr Leu Leu Arg Ala Phe Glu Ala Arg Asp Arg Asn Ile  
                   85                  90                  95  
 Gln Glu Ser Asn Phe Asp Arg Val Asn Phe Trp Ser Met Val Asn Leu  
           100                  105                  110  
 Val Val Met Val Val Val Ser Ala Ile Gln Val Tyr Met Leu Lys Ser  
           115                  120                  125  
 Leu Phe Glu Asp Lys  
           130

<210> 57  
 <211> 387  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> unsure  
 <222> (48)

<220>  
 <221> unsure  
 <222> (113)

<220>  
 <221> unsure  
 <222> (116)

<220>  
 <221> unsure  
 <222> (178)

<400> 57  
 tgtttgaaga taagaggaaa agtagaactt aaaactccaa actagagnac gtaacattga 60  
 aaaatgaggc ataaaaatgc aataaaactgt tacagtcaag accattaatg gtntntntcca 120  
 aaatattttg agatataaaa gtaggaaaca ggtataattt taatgtgaaa attaatgtntt 180  
 cactttctgt gcaagtaatc ctgctgatcc agttgtactt aagtgtgtaa caggaatatt 240  
 ttgcagaata taggtttaac tgaatgaagc catattaata actgcatttt cctaactttg 300  
 aaaaattttg caaatgtctt aggtgattta aataaatgag tattgggcct aattgcacaaa 360  
 aaaaaaaaaa aaaaaaaaaa aaaaaaa 387

<210> 58  
 <211> 1150  
 <212> DNA  
 <213> Homo sapiens

<400> 58  
 ggcggtgac attcagccgg cggttcgggg cgacggactc tccattccag aaccatggcc 60  
 caatttgacc gtaacctgt ggagaagacc cggcgctgg tgaacgctgc tgtgacttac 120  
 tcgaagccctc gattggccac attttggtac tacgccaagg ttgagctggc tccctcccacc 180  
 cctgctgaga tccctagagc tattcagagc ctgaaaaaaa tagtcaatag tgcctcagact 240  
 ggtagcttca aacagctcac agttaaggaa gctgtgctga atggtttggg ggccactgag 300  
 gtgttgatgt ggttttatgt cggagagatt ataggcaagc ggggcatcat tggctatgat 360  
 gtttgaagac caatctttaa catctgatta tatttgattt attatttgag tgttgttggg 420  
 ccatgtgtga tcagactgct atctgaataa aataagattt gtcaaaactc agtgttttct 480  
 ccatcagaca ctccatgaaa ggtcaccaatt tctcttgata ttaagctggg ttgtctttaa 540  
 acaaccctaa atacacgtct gtttagcccg caattggaaa ggatatatgt ggcaatatta 600  
 acctgggtaca tgaatatatg gggataacat ttaatttga aggtttggaa tatatatatt 660  
 taagctttat ttccagaaca gtgagggtta ggtcttgga aaactataac ttgccaaagt 720  
 agaagaaata gtagtaccat atgccaaagt gatagagatg aatcatgtca gtagttagaa 780  
 taacatttca actgttttct ttgctaaaat cacagaaaga ccctattgac aacatctatg 840  
 tctgtaaaaa tgttagagta cttgtcatct tgaatatagc ctcccagaaga gagaacaggg 900  
 tggatattcta agtatgtttc ttgttaacat ctttagcagt aggacagatc catacatgtg 960  
 aaatctgatt tttatgtgtg ttattcgttt gtctggtttt actacctttg caaaaacaaa 1020  
 ataccccaaa gatattttaa caagggtata atttagcatc ttccctggat ctaaatagta 1080  
 tattatatcc tgaaataaat gaaatgattg ctataaaaaa aaaaaaaaaa aaaaaaaaaa 1140  
 aaaaaaaaaa 1150

<210> 59  
 <211> 103  
 <212> PRT  
 <213> Homo sapiens

<400> 59  
 Met Ala Gln Phe Val Arg Asn Leu Val Glu Lys Thr Pro Ala Leu Val  
 1 5 10 15  
 Asn Ala Ala Val Thr Tyr Ser Lys Pro Arg Leu Ala Thr Phe Trp Tyr  
 20 25 30

Tyr Ala Lys Val Glu Leu Val Pro Pro Thr Pro Ala Glu Ile Pro Arg  
 35 40 45

Ala Ile Gln Ser Leu Lys Lys Ile Val Asn Ser Ala Gln Thr Gly Ser  
 50 55 60

Phe Lys Gln Leu Thr Val Lys Glu Ala Val Leu Asn Gly Leu Val Ala  
 65 70 75 80

Thr Glu Val Leu Met Trp Phe Tyr Val Gly Glu Ile Ile Gly Lys Arg  
 85 90 95

Gly Ile Ile Gly Tyr Asp Val  
 100

<210> 60  
 <211> 456  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> unsure  
 <222> (269)

<220>  
 <221> unsure  
 <222> (271)

<400> 60  
 agagattcag gacctgcaga gtcgccagaa gcatgaaatt gaatctttgt atactaaact 60  
 gggcaaggtt cccctgctg tcattattcc cccagctgct cctctgtcgg ggagaagaag 120  
 gagaccact aaaagcaaag gcagcaagtc tagtcgcagc agctcattgg gcaataaaaag 180  
 cccacagctt tcaggcaacc tgtctggta gagtggaaact tcagtcttac accccaaca 240  
 gacctccac cctctggca acatccana ntccgggcag aatcagctgt tacagccct 300  
 taagccatct cctccagtg acaacctcta ttcagccttc accagtgatg gtgccatttc 360  
 agtaccagc cttctgctc caggtcaagg aaccagcagc acaaactctg ttggggcaac 420  
 agtgaacagc caagccgcc aagctcagcc tcctgc 456

<210> 61  
 <211> 130  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> UNSURE  
 <222> (79)..(80)

<400> 61  
 Met Lys Leu Asn Leu Cys Ile Leu Asn Trp Ala Arg Phe Pro Leu Leu  
 1 5 10 15

Ser Leu Phe Pro Gln Leu Leu Leu Cys Arg Gly Glu Glu Gly Asp Pro  
 20 25 30

Leu Lys Ala Lys Ala Ala Ser Leu Val Ala Ala Ala His Trp Ala Ile  
 35 40 45

Lys Ala His Ser Phe Gln Ala Thr Cys Leu Val Arg Val Glu Leu Gln  
 50 55 60

Ser Tyr Thr Pro Asn Arg Pro Ser Thr Leu Leu Ala Thr Ser Xaa Xaa  
65 70 75 80

Pro Gly Arg Ile Ser Cys Tyr Ser Pro Leu Ser His Leu Pro Pro Val  
85 90 95

Thr Thr Ser Ile Gln Pro Ser Pro Val Met Val Pro Phe Gln Tyr Gln  
100 105 110

Ala Phe Leu Leu Gln Val Lys Glu Pro Ala Ala Gln Thr Leu Leu Gly  
115 120 125

Gln Gln  
130

<210> 62  
<211> 188  
<212> DNA  
<213> Homo sapiens

<220>  
<221> unsure  
<222> (24)..(25)

<220>  
<221> unsure  
<222> (171)

<400> 62  
taccctgccc tcctcccttt tttmnacccc tctctttttt attttttctt tgctcttttag 60  
aaccacagtga aaaataaccag ggtactgggg tgcaactctt tcttatgata ggtcattagt 120  
gctttaagca aaagatatta gcagctttga ctgcagcatt agcaattagg naaaaaaaaa 180  
aaaaaaaaa 188

<210> 63  
<211> 752  
<212> DNA  
<213> Homo sapiens

<220>  
<221> unsure  
<222> (151)

<220>  
<221> unsure  
<222> (616)

<220>  
<221> unsure  
<222> (636)

<220>  
<221> unsure  
<222> (662)

<220>  
<221> unsure  
<222> (711)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (722)

&lt;400&gt; 63

```

ccttatggcc tactttaaaa aaaaaccaat accaaagaag cctacaatgt tggccttagc 60
caaaattctg ttgatttcaa cgttggttta ttcacttcta tcggggagcc atggaaaaga 120
aaatcaagac ataaacacaa cacagaacat ngcagaagtt tttaaaacaa tggaaaataa 180
acctatttct ttggaagtg aagcaaactt aaactcagat aaagaaaata taaccacctc 240
aaatctcaag gcgagtcatt cccctccttt gaatctaccc aacaacagcc acggaataac 300
agatttctcc agtaactcat cagcagagca ttctttgggc agtctaaaac ccacatctac 360
catttcacac agccctccct tgatccatag ctttgtttct aaagtgcctt ggaatgcacc 420
tatagcagat gaagatcttt tgcccatctc agcacatccc aatgstacac ctgctctgty 480
ttcaraaaac ttcacttggc ctttgtcaat gacaccgtga aaactcctga taacagttcc 540
attacagtta gcatactcty ttcaraacca acttctccat ctgtgacccc cttgatagtg 600
gaaccaagtg gatggnttac cacaaacagt gatagnntca ctggggttac cccttatcaa 660
gnaaaaacaa ctttcagacc taccttaaaa ttcaccaata attcaaaact ntttccaaat 720
angtcagatc ccccaaaaaa aaaaaaaaaa aa 752

```

&lt;210&gt; 64

&lt;211&gt; 157

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; UNSURE

&lt;222&gt; (35)

&lt;220&gt;

&lt;221&gt; UNSURE

&lt;222&gt; (140)

&lt;220&gt;

&lt;221&gt; UNSURE

&lt;222&gt; (145)

&lt;220&gt;

&lt;221&gt; UNSURE

&lt;222&gt; (147)

&lt;400&gt; 64

```

Met Leu Ala Leu Ala Lys Ile Leu Leu Ile Ser Thr Leu Phe Tyr Ser
  1             5             10            15

Leu Leu Ser Gly Ser His Gly Lys Glu Asn Gln Asp Ile Asn Thr Thr
  20            25            30

Gln Asn Xaa Ala Glu Val Phe Lys Thr Met Glu Asn Lys Pro Ile Ser
  35            40            45

Leu Glu Ser Glu Ala Asn Leu Asn Ser Asp Lys Glu Asn Ile Thr Thr
  50            55            60

Ser Asn Leu Lys Ala Ser His Ser Pro Pro Leu Asn Leu Pro Asn Asn
  65            70            75            80

Ser His Gly Ile Thr Asp Phe Ser Ser Asn Ser Ser Ala Glu His Ser
  85            90            95

```



Leu Gly Ser Leu Lys Pro Thr Ser Thr Ile Ser Thr Ser Pro Pro Leu  
 100 105 110

Ile His Ser Phe Val Ser Lys Val Pro Trp Asn Ala Pro Ile Ala Asp  
 115 120 125

Glu Asp Leu Leu Pro Ile Ser Ala His Pro Asn Xaa Thr Pro Ala Leu  
 130 135 140

Xaa Ser Xaa Asn Phe Thr Trp Ser Leu Ser Met Thr Pro  
 145 150 155

<210> 65

<211> 417

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> (69)

<400> 65

aagcttggca cgaggtcttt agaagaacta caaaacctga atggaaaact tcgaagtga 60  
 ggacaaggna atatgggctt tactaggcag aatcacaggg cagaagttga atataccggc 120  
 aattttgaga gcacccaag agagaaaacc aagtaaaaaa agaaggaggc acacaaaaga 180  
 catctactct tcttcagta ctttatagtt gtgggatttg taagaagaac catgatcagc 240  
 atcttctttt attgtgtgat acctgtaaac tacattacca ttttggatgt ctggatcctc 300  
 ctctaacaag gatgccaaga aagacccaaa acagttattg gcagtgtctg gaatgtgacc 360  
 aggcaggagg cagtgcacatg gaagcagata tggccatgga aaccctacca gatggaa 417

<210> 66

<211> 35

<212> PRT

<213> Homo sapiens

<400> 66

Met Pro Arg Lys Thr Gln Asn Ser Tyr Trp Gln Cys Ser Glu Cys Asp  
 1 5 10 15

Gln Ala Gly Ser Asp Met Glu Ala Asp Met Ala Met Glu Thr Leu  
 20 25 30

Pro Asp Gly  
 35

<210> 67

<211> 359

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> (90)

<220>

<221> unsure

<222> (156)..(157)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (160)

&lt;400&gt; 67

tctgtgttca gtataat ttttttctca accttaata tgaacttagg aaataaggag 60  
 ggaagtacaa agattattga ctatacaacn taccagctga aagaagatc ttcacaaaca 120  
 tctgtatctt tccagaggta tacagaatta aaattnnatn ttcaagcttt aatgatccag 180  
 ttttaagtca acggcagaag tatgttgaat atttcatcac tcaatcttga actgatttag 240  
 aagagactct ttgctgaaat tgaattgcac ttatacatgt aaattgtcaa catgtaattt 300  
 ggaattttct gattaataaa tgtggttttg gacatctaaa aaaaaaaaaa aaaaaaaaaa 359

&lt;210&gt; 68

&lt;211&gt; 656

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 68

aacgaacggc ttgggcgcgg actgggtatcc ggggactgtg acttgcaggg tccgccatgg 60  
 agccagagca gatgctggag ggacaaacgc aggttgcaga aaatcctcac tctgagtacg 120  
 gtctcacaga caacgttgag agaatagtag aaaatgagaa gattaatgca gaaaagtcac 180  
 caaagcagaa ggtagatctc cagtctttgc caactcgtgc ctacctggat cagacagttg 240  
 tgcctatctt attacagga cttgctgtgc ttgcaaagga aagaccacca aatcccattg 300  
 aatttctagc atcttatctt ttaaaaaaca aggcacagtt tgaagatcga aactgactta 360  
 atgggaagaa cagaaaaatt tagttgctac ttagatttta catgattaag aggcagcttt 420  
 aattgccatg atcattccct ctttttggat gtataagaac cttccggaca acagaacctc 480  
 tttctggaat tgcagaagat aacatatctc ctttattttg atttaatcac cataaacccat 540  
 acctatttaa tgagtgtatt ctgtgcaatt tttttctcag attgtcttta actttgtttt 600  
 taaatgacc ttcaaaataa actgtcaaaa caccattaaa aaaaaaaaaa aaaaaa 656

&lt;210&gt; 69

&lt;211&gt; 99

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 69

Met Glu Pro Glu Gln Met Leu Glu Gly Gln Thr Gln Val Ala Glu Asn  
 1 5 10 15

Pro His Ser Glu Tyr Gly Leu Thr Asp Asn Val Glu Arg Ile Val Glu  
 20 25 30

Asn Glu Lys Ile Asn Ala Glu Lys Ser Ser Lys Gln Lys Val Asp Leu  
 35 40 45

Gln Ser Leu Pro Thr Arg Ala Tyr Leu Asp Gln Thr Val Val Pro Ile  
 50 55 60

Leu Leu Gln Gly Leu Ala Val Leu Ala Lys Glu Arg Pro Pro Asn Pro  
 65 70 75 80

Ile Glu Phe Leu Ala Ser Tyr Leu Leu Lys Asn Lys Ala Gln Phe Glu  
 85 90 95

Asp Arg Asn

&lt;210&gt; 70

&lt;211&gt; 979

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 70

```

ggtttgggtga ggaaattacc agagaactat taaagacttg gatgctcttc tcggctttgc 60
tattaagtaa gttggacaag ttgtttggct tctttgagcc tctgttttct ccattctaaa 120
attctaaaaat gggagtgttg aattagatca gtggcttttcg aactttctgc tcctagtagt 180
gagaaataca ttttactcca ctccctggta tgtacacgca ttctgtgtt ttgtgaaaac 240
ctgacaccat gtccttccct cactacatgt aaaacacttt tattcattaa aaagaaaact 300
gactggcttg gacctacaaa ttagtttcat tatttgtaa tgtttgaaag ccattaaaag 360
atgaatatta aggtttcttt atactcaata cttgtagttt tgtttggggg aatgagagga 420
tgcccttggg acctttgtga ggctctcca ctgaggggca atcatgactt ctgttttaaa 480
ccagcccatc catcttctc cagctgtctc ccttatgtct tgcttctctc cctccaacc 540
ttctcagcat aaggactcaa tcctaggctc ctaccccaga cgggtgcctt ccaacgttcc 600
tggtgccagt ggcctccat cagccatgtt cccccagccc ctctccaca ctctgtcac 660
ccagggcccc atccttcagt gcattgcaca ctttgcacgc tgggtcaggg aagattgtgg 720
agagaggaca gtgcacctgg tttccccac atagactgcg tgggggtatg tcctgcttcc 780
gccacttcca actgtggcac ttgggcacgc cctctcagg gcacctccc ttttgtttc 840
cgcaaaatga ggttgtaata gtgcctgceg cactgtctgg cacacagtaa gctctcaaga 900
aatgttagct gttgttgceg ttagaacacc atagctagaa taccatacct ggcattcact 960
taaaaaaaaa aaaaaaaaaa
979

```

&lt;210&gt; 71

&lt;211&gt; 96

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 71

```

Met Thr Ser Val Leu Asn Gln Pro Ile Pro Ser Ser Pro Ala Ala Leu
  1             5             10             15

Leu Met Ser Cys Phe Ser Pro Leu Gln Pro Ser Gln His Lys Asp Ser
             20             25             30

Ile Leu Gly Ser Tyr Pro Arg Arg Val Pro Ser Asn Val Pro Gly Ala
             35             40             45

Ser Gly Pro Pro Ser Ala Met Phe Pro Gln Pro Leu Leu His Thr Leu
             50             55             60

Val Thr Gln Gly Pro Ile Leu Gln Cys Ile Ala His Phe Ala Cys Trp
             65             70             75             80

Val Arg Glu Asp Cys Gly Glu Arg Thr Val His Leu Val Ser Pro Thr
             85             90             95

```

&lt;210&gt; 72

&lt;211&gt; 310

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 72

```

attgaggaaa accacaaaaa acttcaaaac agctacaacg ggaaaaagag agttttgtcc 60
cacagtacgc aggccactag tttattaact tccagtcacc ttgatttttg ctaaaatgaa 120
gactctgcag tctacacttc tcctgttact gcttgtgcct ctgataaagc cagcaccacc 180
aaccacgacg gactcacgca ttatctatga ttatggaaca gataattttg aagaatccat 240
attagccaa gattatgagg ataaatacct ggatggaaaa aatattaagg aaaaagaaac 300
tgtgataata
310

```

<210> 73  
<211> 65  
<212> PRT  
<213> Homo sapiens

<400> 73  
Met Lys Thr Leu Gln Ser Thr Leu Leu Leu Leu Leu Val Pro Leu  
1 5 10 15

Ile Lys Pro Ala Pro Pro Thr Gln Gln Asp Ser Arg Ile Ile Tyr Asp  
20 25 30

Tyr Gly Thr Asp Asn Phe Glu Glu Ser Ile Phe Ser Gln Asp Tyr Glu  
35 40 45

Asp Lys Tyr Leu Asp Gly Lys Asn Ile Lys Glu Lys Glu Thr Val Ile  
50 55 60

Ile  
65

<210> 74  
<211> 303  
<212> DNA  
<213> Homo sapiens

<220>  
<221> unsure  
<222> (7)

<220>  
<221> unsure  
<222> (10)

<220>  
<221> unsure  
<222> (17)

<220>  
<221> unsure  
<222> (29)

<220>  
<221> unsure  
<222> (40)

<220>  
<221> unsure  
<222> (49)

<220>  
<221> unsure  
<222> (63)

<220>  
<221> unsure  
<222> (65)

<220>

<221> unsure  
<222> (73)

<220>  
<221> unsure  
<222> (78)

<220>  
<221> unsure  
<222> (83)

<220>  
<221> unsure  
<222> (88)

<220>  
<221> unsure  
<222> (93)

<220>  
<221> unsure  
<222> (130)

<220>  
<221> unsure  
<222> (132)

<220>  
<221> unsure  
<222> (153)

<220>  
<221> unsure  
<222> (155)

<220>  
<221> unsure  
<222> (180)

<220>  
<221> unsure  
<222> (200)

<220>  
<221> unsure  
<222> (211)

<220>  
<221> unsure  
<222> (241)

<220>  
<221> unsure  
<222> (266)

<220>  
<221> unsure  
<222> (279)

<400> 74

```

cccaagnaan tttcaanttt ttgcctttnc tggcctttan tggatccna aagcatttaa 60
ggnanagtgt ccnaaaanttt tgnaaagnta aangtttccc atgatcgctc attttttttt 120
tatgattcan angttattcc ttataaagta agnantttgt tttcctccta tcaaggcagn 180
tattttatta aatttttcan ttagtttgag naatagcaga tagtttcata tttagggaaa 240
ntttccaaat aaaataaatg ttattntttg ataaagagnt aaaaaaaaaa aaaaaaaaaa 300
aaa 303

```

&lt;210&gt; 75

&lt;211&gt; 1823

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 75

```

gawagcttgg cactgaggcac aagtagctac gactgcaagc acctgccacc ataaagggct 60
gcatttttgc accataaagg gctgcatttt tttaaaaagc ctaggcagct ctaacatcat 120
ctgatattga cacaaggcca acagtttccct tatttacatc cttacctcta aaagataact 180
caaaagtgaca aaaacgtggt ccttccccac ttagagacaa tgattaacag ggccctatat 240
gttctttacca catacagagg atgcatttat ttttgctcta tgacacttgc aaaaatctct 300
actgtaatta atttgggtct attattaact ctctgttcca tcatagaatg tggccaggcc 360
ttacaatgga gagccagagt taaaacttca agttgcatct gtttttgggc tgagtcacca 420
cctttgcttc atgctccttt gtctgcaaag gcctaggatt ctttctttaa atgaaatgct 480
taggactttg tggcttggtta cttttgtcat ttaactgcag tgctattctt tgaaagctgc 540
tatgtgtatt ttctctgaag tctgcatttt actaaaattt acaacagtct gatgattgat 600
tgattactgt ccaggtacat tttagaaaaa gtgttcttct tccagtttgt tttactaagc 660
aaactttgag taaatccttt gtcttatatt gaatccagtc ccaaagtgtt caggtgagtt 720
tctctagttc cataaacaac acatacatag tgggaactcc ctggtatgcc atagagcaca 780
caagaacccc aatattaatg ctaacaatta taccagtcca tttgtttat tctgtggaat 840
tgacttgaca aagcatgaag atattcccag tgtctgtctg ataataattt gcatctaaga 900
atgggtttga ctcaagatct tgggtttacca agatgtctta aatgttcagt aaatatcttt 960
cttacagtc agtagcttag agcatgtttg ctgattgata ttacatttaa acttggggct 1020
acagcttggt acctagaatt ttgagatact aagagaatgc aattttaaat gccactggt 1080
tttatttgggt ttggagagag ggtctcattc tgtcaccag gctagagtac agtgggaaca 1140
gtcatggctc actgcagctt aagctcactg cctgggctca agtgatctta ctccagcctc 1200
ccaagcagct gggactacac caccacaggt atgcaccacc atacctggtt gattttttaa 1260
attttttcta gagatgaggt ctactatgt tggccagctg gtctcactc cctcagttca 1320
cgcatctctc ccacctccac ctcccaaact gctgggatta caggtgtgag ccaccatgcc 1380
caacacccac tgatctttaa ctctcacatg ttgggcataa gaagtcacta tataattgtt 1440
actggaaagc aagacttaac gaacaattct gactatgaaa aatgtctctt tcagtttgtt 1500
ctgtaaatat ttagaaaagt gacagctgtc aacytcagrg taactatttc taaaaatgta 1560
aatatgtatt aatccttgta tcttttatgg taattttgca tattgatag aattatataa 1620
aattgtttta aataaaaggt gtccctgaat tactgaccac ccatagatgt ctactgttac 1680
caggttttac aatgcaaatt ttcactaata cctgggttta atacagctca catcactgaa 1740
tgttacacat gagtttaaat gggttaatat acaggttttg ttataataaa gttactgatt 1800
aaattaaaaa aaaaaaaaaa aaa 1823

```

&lt;210&gt; 76

&lt;211&gt; 78

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 76

```

Met Ile Asn Arg Ala Leu Tyr Val Leu Thr Thr Tyr Arg Gly Cys Ile
  1             5             10             15

```

```

Tyr Phe Cys Ser Met Thr Leu Ala Lys Ile Ser Thr Val Ile Asn Leu
      20             25             30

```

```

Gly Leu Leu Leu Thr Leu Cys Ser Ile Ile Glu Cys Gly Gln Ala Leu
      35             40             45

```

Gln Trp Arg Ala Arg Val Lys Thr Ser Ser Cys Ile Cys Phe Trp Ala  
50 55 60

Glu Ser Pro Pro Leu Pro His Ala Pro Leu Ser Ala Lys Ala  
65 70 75

<210> 77  
<211> 583  
<212> DNA  
<213> Homo sapiens

<220>  
<221> unsure  
<222> (5)

<220>  
<221> unsure  
<222> (217)

<400> 77  
cacgnggggtg aggcgcgactg ctgaagacag ctcgccaccc tccttgccctc cactccaatc 60  
caggggctgg ggccacattc ttgacctca ttatccctca gatcaggtga gatcgacagg 120  
aggtgttgat ggcagtgcca gcaattattg ctaatccgtt tgcacacctta tgcatagata 180  
tgaattcaga ctttgtgaat ttccagaggt gtgggtnata taatagaatt cagtgaagg 240  
gcatggctga tcttgtgcaa attaaaagtt atggggcata agaatagcaa aagttgaact 300  
tcttttaaaa aggaaagtac cctgagagcc agtattgggt gaggcctctc agtatgcccc 360  
ggttggcagc actgagaacc gcaggaacgg cctgttggtta caaaaaggag attgactcag 420  
ctgcccttgg tgcacttgac tgactatgac tgctgagaga ttccaaggac ccttaatgcc 480  
agggctaacc tctccatgtg cagtgaagac tctggaggaa gtgtcatcct ctggccttgt 540  
gtggtactca ttatggtgca gtgcgggcat gaaatgaaga cac 583

<210> 78  
<211> 29  
<212> PRT  
<213> Homo sapiens

<400> 78  
Met Cys Ser Glu Thr Ser Gly Gly Ser Val Ile Leu Trp Leu Cys Val  
1 5 10 15

Val Leu Ile Met Val Gln Cys Gly His Glu Met Lys Thr  
20 25

<210> 79  
<211> 311  
<212> DNA  
<213> Homo sapiens

<220>  
<221> unsure  
<222> (40)

<220>  
<221> unsure  
<222> (234)

<220>  
<221> unsure

&lt;222&gt; (247)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (251)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (260)

&lt;400&gt; 79

```

cccaaatagg cttacagata cgatatgttt taaatgtttt gtatttaaca aaaacatact 60
gacactgttt ggaaatggca acaggaagat agcaaaatga atactaacat tacgaaaaga 120
tgaacaggta catgttccaa ggcagggtggc tgtgaacttc ctctgagtga aggcattccc 180
tcagcacct ttcagcctgc tagttaggac gacccgcgcg caccctccag gacntccagc 240
cctgcantgc nttttttttt ttttaaataa ttcttcattg agttctaata tgtaaaaaaa 300
aaaaaaaaa a 311

```

&lt;210&gt; 80

&lt;211&gt; 405

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 80

```

aagcttgcca cgagggcggt tgaggccttc ggtggtgaac gagtctccag caccatgtct 60
ggtttgtctg gccaccagc ccggcgcggc ccttttccgt tagcgttget gcttttgttc 120
ctgctcgggc ccagattggt ccttgccatc tccttccatc tgcccattaa ctctcgcaag 180
tgctccgtg aggagattca caaggacctg ctagtgactg gcgcgtacga gatctccgac 240
cagtctgggg gcgctggcgg cctgcgcagc cacctcraga tcacagattc tgctggccat 300
attctctact ccaaagagga tgcaaccaag gggaaatttg cctttaccac tgaagattat 360
gacatgtttg aagtgtgttt tgagagcaag ggaacagggc ggata 405

```

&lt;210&gt; 81

&lt;211&gt; 117

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; UNSURE

&lt;222&gt; (75)

&lt;400&gt; 81

```

Met Ser Gly Leu Ser Gly Pro Pro Ala Arg Arg Gly Pro Phe Pro Leu
  1           5           10          15

Ala Leu Leu Leu Leu Phe Leu Leu Gly Pro Arg Leu Val Leu Ala Ile
  20           25           30

Ser Phe His Leu Pro Ile Asn Ser Arg Lys Cys Leu Arg Glu Glu Ile
  35           40           45

His Lys Asp Leu Leu Val Thr Gly Ala Tyr Glu Ile Ser Asp Gln Ser
  50           55           60

Gly Gly Ala Gly Gly Leu Arg Ser His Leu Xaa Ile Thr Asp Ser Ala
  65           70           75           80

Gly His Ile Leu Tyr Ser Lys Glu Asp Ala Thr Lys Gly Lys Phe Ala
  85           90           95

```



Phe Thr Thr Glu Asp Tyr Asp Met Phe Glu Val Cys Phe Glu Ser Lys  
 100 105 110

Gly Thr Gly Arg Ile  
 115

<210> 82  
 <211> 225  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> unsure  
 <222> (42)

<220>  
 <221> unsure  
 <222> (61)

<220>  
 <221> unsure  
 <222> (71)

<220>  
 <221> unsure  
 <222> (99)

<400> 82  
 tctttcaatt taccttgatg aaacaccctt aactttttct tnacccttag ctgaaatggt 60  
 nacatagctt ntggatgatg cttttcatga tttatatnt cttaaaatgg tgatggatgt 120  
 gacacctcat aaaagtgggc ttgaaactgt agataactct taaagaaaat gtcatttttag 180  
 acaattaaaa tatttggtgt caaaaaaaaa aaaaaaaaaa aaaaa 225

<210> 83  
 <211> 1711  
 <212> DNA  
 <213> Homo sapiens

<400> 83  
 cgagggcagg tcagtcaggt tcctgggggc tctgtttcac aagcaagata cagccagccc 60  
 cacctaattt tgtttccctg gcacctctct gctcagtgcg acattgtcac acttaaccga 120  
 tctgttttct ctaatgcacg acagattcct ttcagacagg acaactgtga tatttcagtt 180  
 cctgattgta aatacctcct aagcctgaag cttctgttac tagccattgt gagcttcagt 240  
 ttcttcatct gcaaaatggg cataatacaa tctattcttg ccacatcaag ggattgttat 300  
 tcctttaaaa aaaaaaccaa taccaaagaa gcctacaatg ttggccttag ccaaaattct 360  
 gttgatttca acgttggttt attcacttct atcggggagc catggaaaag aaaatcaaga 420  
 cataaacaca acacagaaca ttcagaagtt tttaaaaaaa tggaaaataa acctatttct 480  
 ttggaaagtg aagcaaaact aaactcagat aaagaaaata taaccacctc aaatctcaag 540  
 gcgagtcatt cccctccttt gaattctacc aacaacagcc acggaataac agatttctcc 600  
 agtaactcgt cagcagagca ttctttgggc agtctaaaac ccacatctac catttccaca 660  
 agccctccct tgatccatag ctttgtttct aaagtgcctt ggaatgcacc tatagcagat 720  
 gaagatcttt tgcccatctc agcacatccc aatgctacac ctgctctgtc ttcagaaaac 780  
 ttcacttggt ctttgggtcaa tgacaccgtg aaaactcctg ataacagttc cattacagtt 840  
 agcatcctct cttcagaacc aacttctcca tctgtgacct ccttgatagt ggaaccaagt 900  
 ggatggctta ccacaaacag tgatagcttc actgggttta tcccttatca agaaaaaaca 960  
 actctacagc ctaccttaaa attcaccat aattcaaaa tctttccaaa tacgtcagat 1020  
 ccccaaaaag aaaatagaaa tacaggaata gtattcgggg ccatttttagg tgctattctg 1080  
 ggtgtctcat tgcttactct tgtgggctac ttgttgtgtg gaaaaaggaa aacggattca 1140  
 ttttcccatc ggcgacttta tgacgacaga aatgaaccag ttctgcgatt agacaatgca 1200

```

ccggaacctt atgatgtgag ttttgggaat tctagctact acaatccaac tttgaatgat 1260
tcagccatgc cagaaagtga agaaaatgca cgtgatggca ttcctatgga tgacatacct 1320
ccacttcgta cttctgtata gaactaacag caaaaaggcg ttaaacagca agtgtcatct 1380
acatcctagc cttttgacaa attcatcttt caaaagggtta cacaaaatta ctgtcacgtg 1440
gattttgtca aggagaaatca taaaagcagg agaccagtag cagaaatgta gacaggatgt 1500
atcatccaaa ggttttcttt cttacaattt ttggccatcc tgaggcattt actaagtagc 1560
cttaatttgt attttagtag tattttctta gtagaaaata tttgtggaat cagataaaac 1620
taaaagattt caccattaca gccctgcctc ataactaaat aataaaaatt attccaccaa 1680
aaaattctaa aacaaaaaaa aaaaaaaaaa a 1711

```

&lt;210&gt; 84

&lt;211&gt; 361

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 84

```

Met Gly Ile Ile Gln Ser Ile Leu Ala Thr Ser Arg Asp Cys Tyr Ser
  1             5             10             15

Phe Lys Lys Lys Thr Asn Thr Lys Glu Ala Tyr Asn Val Gly Leu Ser
      20             25             30

Gln Asn Ser Val Asp Phe Asn Val Val Leu Phe Thr Ser Ile Gly Glu
      35             40             45

Pro Trp Lys Arg Lys Ser Arg His Lys His Asn Thr Glu His Ser Glu
      50             55             60

Val Phe Lys Thr Met Glu Asn Lys Pro Ile Ser Leu Glu Ser Glu Ala
      65             70             75             80

Asn Leu Asn Ser Asp Lys Glu Asn Ile Thr Thr Ser Asn Leu Lys Ala
      85             90             95

Ser His Ser Pro Pro Leu Asn Leu Pro Asn Asn Ser His Gly Ile Thr
      100            105            110

Asp Phe Ser Ser Asn Ser Ser Ala Glu His Ser Leu Gly Ser Leu Lys
      115            120            125

Pro Thr Ser Thr Ile Ser Thr Ser Pro Pro Leu Ile His Ser Phe Val
      130            135            140

Ser Lys Val Pro Trp Asn Ala Pro Ile Ala Asp Glu Asp Leu Leu Pro
      145            150            155            160

Ile Ser Ala His Pro Asn Ala Thr Pro Ala Leu Ser Ser Glu Asn Phe
      165            170            175

Thr Trp Ser Leu Val Asn Asp Thr Val Lys Thr Pro Asp Asn Ser Ser
      180            185            190

Ile Thr Val Ser Ile Leu Ser Ser Glu Pro Thr Ser Pro Ser Val Thr
      195            200            205

Pro Leu Ile Val Glu Pro Ser Gly Trp Leu Thr Thr Asn Ser Asp Ser
      210            215            220

Phe Thr Gly Phe Ile Pro Tyr Gln Glu Lys Thr Thr Leu Gln Pro Thr
      225            230            235            240

```

Leu Lys Phe Thr Asn Asn Ser Lys Leu Phe Pro Asn Thr Ser Asp Pro  
 245 250 255

Gln Lys Glu Asn Arg Asn Thr Gly Ile Val Phe Gly Ala Ile Leu Gly  
 260 265 270

Ala Ile Leu Gly Val Ser Leu Leu Thr Leu Val Gly Tyr Leu Leu Cys  
 275 280 285

Gly Lys Arg Lys Thr Asp Ser Phe Ser His Arg Arg Leu Tyr Asp Asp  
 290 295 300

Arg Asn Glu Pro Val Leu Arg Leu Asp Asn Ala Pro Glu Pro Tyr Asp  
 305 310 315 320

Val Ser Phe Gly Asn Ser Ser Tyr Tyr Asn Pro Thr Leu Asn Asp Ser  
 325 330 335

Ala Met Pro Glu Ser Glu Glu Asn Ala Arg Asp Gly Ile Pro Met Asp  
 340 345 350

Asp Ile Pro Pro Leu Arg Thr Ser Val  
 355 360

<210> 85  
 <211> 565  
 <212> DNA  
 <213> Homo sapiens

<400> 85  
 ttccagacca gcttggtgtca atagggtcct acagagcagc tgatatacagc agttttacta 60  
 gtatgcagga cctgaaagaa tatctcaaag ggaaaacaat gtttcataat gttcaggaag 120  
 ttatctatag agcagctaag gggaaataat cttgtaacag ggtctgggtg attctgaggt 180  
 aataggccccc aaacaacccat ggggaagcag gtcagagggc aagctggcct agtgtttaac 240  
 attgaatggg ctgaaagttt ggtttatttt tgtttcttgt ttctccccct cccttettac 300  
 ctgaataaatt ttatgaagtt tatagggatg gtttcaggac ctccattcta tctgttcctg 360  
 aaatattaca aaaagattat tattgtagca ctcactaat tgtgttttat ctggttgttt 420  
 gcattgtctgt ttcttccccca gtgagttgta aattgcttaa gggcaaacag acgcataccta 480  
 tttatctgtc tgtcactaac attaagcaca gcatttggtg tacagtcatac actctaataa 540  
 agtttgaaaa aaaaaaaaaa aaaaa 565

<210> 86  
 <211> 66  
 <212> PRT  
 <213> Homo sapiens

<400> 86  
 Met Gly Lys Gln Val Arg Gly Gln Ala Gly Leu Val Phe Asn Ile Glu  
 1 5 10 15

Trp Ala Glu Ser Leu Val Tyr Phe Cys Phe Leu Phe Leu Pro Leu Pro  
 20 25 30

Ser Tyr Leu Asn Asn Phe Met Lys Phe Ile Gly Met Val Ser Gly Pro  
 35 40 45

Pro Phe Tyr Leu Phe Leu Lys Tyr Tyr Lys Lys Ile Ile Ile Val Ala  
 50 55 60

Leu Ile  
65

<210> 87  
<211> 636  
<212> DNA  
<213> Homo sapiens

<220>  
<221> unsure  
<222> (528)

<220>  
<221> unsure  
<222> (530)

<220>  
<221> unsure  
<222> (580)

<400> 87  
cacgaggga aaaaagagtt ttttttttag atcatcagct attgttagtg tttgtgtatg 60  
ttatgtgtgg ctcaagacaa ctttgcttct tttaatatag gcagggaagt caaaagattg 120  
gatatccctg ctttatacca agaaagacaa caccacacat ttgcagtgcc tgaaaacact 180  
accagccatc tgaaaaacat gtgacttcta acttctgttc tttttgttag cagtgggaatc 240  
ccacggtgat atctgaggga tgtggttacc ttttgaggga ggttgacggt ttctaaggat 300  
gattctttct gagtgaata ttgtcagtgt cattgacctt ttcattattt caactattat 360  
tattccaggt tatcaatact ctggctgacc atcatcatcg tgagactgac tttggtgtag 420  
gagttcgaga ccaccctggc caacatggca aaaccccatc tccacaaaaa ttggataatt 480  
tgataattat cattattggg tttctgagac gttacacatt taacattntn ttctgcacaa 540  
gttgcccttg tgtgagtata ctaactttct gtagagggtan acttgtaatc acaaataaga 600  
ataaattata taaaacaaaa aaaaaaaaaa aaaaaa 636

<210> 88  
<211> 105  
<212> PRT  
<213> Homo sapiens

<220>  
<221> UNSURE  
<222> (76)

<220>  
<221> UNSURE  
<222> (93)

<400> 88  
Phe Phe Leu Ser Glu Ile Leu Ser Val Ser Leu Thr Phe Ser Leu Phe  
1 5 10 15  
Gln Leu Leu Leu Phe Gln Val Ile Asn Thr Leu Ala Asp His His His  
20 25 30  
Arg Glu Thr Asp Phe Gly Val Gly Val Arg Asp His Pro Gly Gln His  
35 40 45  
Gly Lys Thr Pro Ser Pro Gln Lys Leu Asp Asn Leu Ile Ile Ile Ile  
50 55 60

Ile Gly Phe Leu Arg Arg Tyr Thr Phe Asn Ile Xaa Phe Cys Thr Ser  
65 70 75 80

Cys Leu Cys Val Ser Ile Leu Thr Phe Cys Arg Gly Xaa Leu Val Ile  
85 90 95

Thr Asn Lys Asn Lys Leu Tyr Lys Thr  
100 105

<210> 89

<211> 861

<212> DNA

<213> Homo sapiens

<400> 89

```

ggccggaggg cagatgactc tgagaagggc aagcacttta accttttaag cccaaccaga 60
tgagttgcct gcagtttttg aggccttcag agcatttcac tagacctctg tctgtgtcgg 120
tccagtgtct ttageccaagc ttgtattaaa gatgacttcc ttgtttgctc aagaaattcg 180
cctttctaaa agacatgaag aaatagtatc acaaagatta atgttacttc aacaaatgga 240
gaataaattg ggtgatcaac acacagaaaa ggcatctcaa ctccaaactg ttgagactgc 300
ttttaaaagg aaccttagtc ttttaaagga tatagaagca gcagaaaagt cactacagac 360
caggattcac ccacttcac ggcctgaggt ggtttctctt gagactcgtt actgggcatc 420
agtagaagaa tatattccca aatgggaaca gtttctttta ggaagagcac catatccttt 480
tgctgttgaa aatcaaaatg aagcagaaaa taccattcaa aatgaggcac agcgataact 540
tcttcacatg ctatttcaaa aagcctggtt aataaagctg aatgttaagg tgtatgtagg 600
ttattgcagg aacttttaga attaaatatg ttcattttct tcgattatct cctaagtgac 660
agtgaagata tgagaattta ctggcaagtc acatgttatc acctactact atttcaaggt 720
catgaatttg ctttctacca aaccataga tgtgttaaac acgaatatta aaaggtggac 780
tctttattaa ttaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 840
aaaaaaaaaa aaaaaaaaaa a

```

861

<210> 90

<211> 128

<212> PRT

<213> Homo sapiens

<400> 90

Met Thr Ser Leu Phe Ala Gln Glu Ile Arg Leu Ser Lys Arg His Glu  
1 5 10 15

Glu Ile Val Ser Gln Arg Leu Met Leu Leu Gln Gln Met Glu Asn Lys  
20 25 30

Leu Gly Asp Gln His Thr Glu Lys Ala Ser Gln Leu Gln Thr Val Glu  
35 40 45

Thr Ala Phe Lys Arg Asn Leu Ser Leu Leu Lys Asp Ile Glu Ala Ala  
50 55 60

Glu Lys Ser Leu Gln Thr Arg Ile His Pro Leu Pro Arg Pro Glu Val  
65 70 75 80

Val Ser Leu Glu Thr Arg Tyr Trp Ala Ser Val Glu Glu Tyr Ile Pro  
85 90 95

Lys Trp Glu Gln Phe Leu Leu Gly Arg Ala Pro Tyr Pro Phe Ala Val  
100 105 110

Glu Asn Gln Asn Glu Ala Glu Asn Thr Ile Gln Asn Glu Ala Gln Arg  
 115 120 125

<210> 91  
 <211> 709  
 <212> DNA  
 <213> Homo sapiens

<400> 91  
 gtcacttcgc ccacgtggt catcacttca ggcatcgcag ccacgtggt gtcacgctac 60  
 ctccctagca ccccccctgc ctggacagt tttagctcga gcgtggcctg tgctctcctt 120  
 tctctgacct gtgcctcgc cctcttgcc tccatcgcca tgacctttgc caccagggc 180  
 aaggcactgc tggctgctg cacttttggg agctctgaac tactggcct cgcacctgac 240  
 tgctccctgc accccacacg catttatagc tccagcctgt gcctctgggg catcgcccta 300  
 gtgctctgcg tggcggagaa cgtgtttgct gtacgctgtg ctcagctcac ccaccagctg 360  
 ctggagctga ggccctggtg ggggaaaagc agccaccaca tgatgcggga gaaccagag 420  
 ctggtggagg gccgtgacct gctgagctgc accagctctg agcctctgac cctctgagag 480  
 atgatgtcct gccaggccc gatggccact aggaccctgc aagcaactct gctctgtgac 540  
 caggccagga ttctggagc tggcctgaga gggctcaatg gaccctcggg gacccaagt 600  
 gggctttcaa cctctcccc caccacccag cccactgcac tgaaatgaga ctttattctg 660  
 aaattattaa aaagaacaga gatgctcaaa aaaaaaaaaa aaaaaaaaaa 709

<210> 92  
 <211> 105  
 <212> PRT  
 <213> Homo sapiens

<400> 92  
 Met Thr Phe Ala Thr Gln Gly Lys Ala Leu Leu Ala Ala Cys Thr Phe  
 1 5 10 15  
 Gly Ser Ser Glu Leu Leu Ala Leu Ala Pro Asp Cys Pro Phe Asp Pro  
 20 25 30  
 Thr Arg Ile Tyr Ser Ser Ser Leu Cys Leu Trp Gly Ile Ala Leu Val  
 35 40 45  
 Leu Cys Val Ala Glu Asn Val Phe Ala Val Arg Cys Ala Gln Leu Thr  
 50 55 60  
 His Gln Leu Leu Glu Leu Arg Pro Trp Trp Gly Lys Ser Ser His His  
 65 70 75 80  
 Met Met Arg Glu Asn Pro Glu Leu Val Glu Gly Arg Asp Leu Leu Ser  
 85 90 95  
 Cys Thr Ser Ser Glu Pro Leu Thr Leu  
 100 105

<210> 93  
 <211> 419  
 <212> DNA  
 <213> Homo sapiens

<400> 93  
 aggaaggaca ccaatgaatc agcttgggac ctcttttaggc cttccccctt tctccaccc 60  
 cgatgtcct tagtgatgct ctgaggcgtg gccacgatct cctcccagg tggatcgcc 120  
 cacctgaaaa aatcctgaga atttctccca tcttggcctc tccagaaac cggccaggca 180

aggaaagagg cccgtcacca gaagccagca ggcgtgggggt gtgatactct ctatagccac 240  
 tacagggcgc gcgcaggtcg cggatctccc cagttgctaa tcccggctct gccactcaat 300  
 cctatcccta gttcccgcgc gcgggtcccc cgccttgacg tctccagccg tgcggggccg 360  
 ggagcaggcc tccggcctcc cagacttcta gagcccgcgc ggcccatctt tgtactcat 419

<210> 94

<211> 93

<212> PRT

<213> Homo sapiens

<400> 94

Glu Phe Leu Pro Ser Trp Pro Leu Pro Glu Thr Gly Gln Ala Arg Lys  
 1 5 10 15

Glu Ala Gly His Gln Lys Pro Ala Gly Val Gly Cys Asp Thr Leu Tyr  
 20 25 30

Ser His Tyr Arg Ala Arg Ala Gly Arg Gly Ser Pro Gln Leu Leu Ile  
 35 40 45

Pro Ala Leu Pro Leu Asn Pro Ile Pro Ser Ser Arg Ala Arg Val Pro  
 50 55 60

Arg Leu Ala Val Ser Ser Arg Ala Gly Pro Gly Ala Gly Leu Arg Pro  
 65 70 75 80

Pro Arg Leu Leu Glu Pro Ala Gly Pro Ile Phe Val Leu  
 85 90

<210> 95

<211> 220

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> (37)

<220>

<221> unsure

<222> (55)

<220>

<221> unsure

<222> (69)

<220>

<221> unsure

<222> (86)

<220>

<221> unsure

<222> (103)

<220>

<221> unsure

<222> (155)

<220>

<221> unsure  
<222> (157)

<220>  
<221> unsure  
<222> (167)

<220>  
<221> unsure  
<222> (178)

<220>  
<221> unsure  
<222> (193)

<400> 95  
tttgtttgcc atttattagc catgtgaaat tggccanac aattcacctc cctgngcctc 60  
agtttcctna tttgtcaaat gggggnttat aaacacctac ctngcagggt tggttgagag 120  
atttaatgcg ataatgtatg taaagcgctt tgcanantgc ctggcanaca gtaggcgntc 180  
aataaattta agnttcctt taaaaaaaaa aaaaaaaaaa 220

<210> 96  
<211> 431  
<212> DNA  
<213> Homo sapiens

<220>  
<221> unsure  
<222> (29)

<220>  
<221> unsure  
<222> (82)

<220>  
<221> unsure  
<222> (106)

<220>  
<221> unsure  
<222> (118)

<220>  
<221> unsure  
<222> (167)..(168)

<220>  
<221> unsure  
<222> (182)

<220>  
<221> unsure  
<222> (190)

<220>  
<221> unsure  
<222> (204)

<220>  
<221> unsure



&lt;222&gt; (214)

&lt;400&gt; 96

cacgaggggc agtgtttgct tttgccganc tgggtgccga cagctccctg ggtgtccggg 60  
 gtgggagAAC tgttgacaga anctctccgg gccctcagg gcttanatcc cacttgantc 120  
 gtaagccttc ttgcttttga taacacagta ttatttctct tactgttnaa aaaaaaattt 180  
 tnttaccaan caagaatttt tttnggaaag aaanggacaa acctataaat taactcaacc 240  
 tatactctccc ttgaaaatac tttcaggctc caccaaaacg tagaactgaa agcatgtatt 300  
 ttggaagaaa gagatacatt ttgtatgctt tcttttcctt ttgtagattc ccagtttatt 360  
 ttctaagact gcaaagatca ctttgtcacc agccctggga cctgagacca aggggggtgtc 420  
 ttgtgggcag t 431

&lt;210&gt; 97

&lt;211&gt; 46

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 97

Met Tyr Phe Gly Arg Lys Arg Tyr Ile Leu Tyr Ala Phe Phe Ser Phe  
 1 5 10 15

Cys Arg Phe Pro Val Tyr Phe Leu Arg Leu Gln Arg Ser Leu Cys His  
 20 25 30

Gln Pro Trp Asp Leu Arg Pro Arg Gly Cys Leu Val Gly Ser  
 35 40 45

&lt;210&gt; 98

&lt;211&gt; 341

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (82)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (216)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (219)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (222)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (311)

&lt;400&gt; 98

tgtgtcttgg agctgactta ctaagtggaa tgagccgagg atttgaatat cagttctaac 60  
 cttgatagaa gaaccttggg tncatgtgg ttcacattaa gaggatagaa tcctttggaa 120  
 tcttatggca accaaatgtg gcttgacgaa gtcgtggttt tattttttaa acaccggtgt 180  
 gtaaatattat tcaactaacg atgggaaatg tattantntn gnacacagtg gactgaagtg 240  
 caatttgttg aaagggaaac agtcattgaa gagaaaaaaa aaagcccaat acttagagtc 300  
 ccaattttgt ntcatttgcc aaaaaaaaaa aaaaaaaaaa a 341

<210> 99  
 <211> 1491  
 <212> DNA  
 <213> Homo sapiens

<400> 99  
 ggacacctct tggagtcac ttggcatgat taccggttga gccacatatt ggaaaaatga 60  
 ttcttttttg agcactgttc tctgtcttag acccagtact cactattgct gctagtctca 120  
 gtttcaaaga tccatttgtc attccactgg gaaaagaaaa gattgcagat gcaagaagaa 180  
 aggaattggc aaaggatact agaagtgatc acttaacagt tgtgaatgcg tttgagggct 240  
 gggaagaggc taggcgacgt ggtttcagat acgaaaagga ctattgtctgg gaatattttc 300  
 tgtcttcaaa cacactgcag atgctgcata acatgaaagg acagtttgcg gagcatcttc 360  
 ttggagctgg atttgtaagc agtagaaatc ctaaagatcc agaattctaat ataaattcag 420  
 ataattgaga gataattaaa gctgtcatct gtgctgggtt atatcccaaa gttgctaaaa 480  
 ttcgactaaa tttgggtaaa aaaagaaaaa tggtaaaagt ttacacaaaa accgatggcc 540  
 tgggtgctgt tcatcctaaa tctgttaatg tggagcaaac agactttcac tacaactggc 600  
 ttatctatca cctaagatg agaacaagca gtatatactt gtatgactgc acagagggtt 660  
 cccatactg tctcttgttt tttggaggtg acattttccat ccagaaggat aacgatcagg 720  
 aaactattgc tgtagatgag tggattgtat ttcagtctcc agcaagaatt gccatcttg 780  
 ttaagggaatt aagaaaggaa ctagatatcc ttctgcaaga gaagattgaa agtcctcctc 840  
 ctgtagactg gaatgacact aaatccagag actgtgcagt actgtcagct attatagact 900  
 tgatcaaaac acaggaaaag gcaactccca ggaactttcc gccacgattc caggatggat 960  
 attacagctg acagcttttc aggggtgggc tgaaaagcca gtttgacagc cattcttcat 1020  
 cattgtttaa attttggctg gatgccaaac cctgggacat gaacaatttt catgtgtgtaag 1080  
 gtagaagcct tcagtaggta gtaaagactt aatgtgcagt acttgatgtt atatgtagag 1140  
 atatatatat atatatatat ataccataaa agcaatatgt tctctgatca tatactctgc 1200  
 tgtggtcatg cccactcttt gggagtatat tccctttata tatattgagt attgtaccac 1260  
 ttgagaaatt cctttgttct gttatacaaa attaatcttt ctgctcataa tgattgatga 1320  
 taccaccagt aaaaatagga tgtttacccc aaaacaagtg tcaattgaaga atttgaacac 1380  
 aaccacattt tttaaaatga aacttctatc ggaagtaaat taatttgttg taataaagtc 1440  
 cagtatttaa taaaatgtac aatgttaaat ctcaaaaaaa aaaaaaaaaa a 1491

<210> 100  
 <211> 304  
 <212> PRT  
 <213> Homo sapiens

<400> 100  
 Met Ile Leu Phe Gly Ala Leu Phe Cys Cys Leu Asp Pro Val Leu Thr  
 1 5 10 15  
 Ile Ala Ala Ser Leu Ser Phe Lys Asp Pro Phe Val Ile Pro Leu Gly  
 20 25 30  
 Lys Glu Lys Ile Ala Asp Ala Arg Arg Lys Glu Leu Ala Lys Asp Thr  
 35 40 45  
 Arg Ser Asp His Leu Thr Val Val Asn Ala Phe Glu Gly Trp Glu Glu  
 50 55 60  
 Ala Arg Arg Arg Gly Phe Arg Tyr Glu Lys Asp Tyr Cys Trp Glu Tyr  
 65 70 75 80  
 Phe Leu Ser Ser Asn Thr Leu Gln Met Leu His Asn Met Lys Gly Gln  
 85 90 95  
 Phe Ala Glu His Leu Leu Gly Ala Gly Phe Val Ser Ser Arg Asn Pro  
 100 105 110

Lys Asp Pro Glu Ser Asn Ile Asn Ser Asp Asn Glu Lys Ile Ile Lys  
 115 120 125  
 Ala Val Ile Cys Ala Gly Leu Tyr Pro Lys Val Ala Lys Ile Arg Leu  
 130 135 140  
 Asn Leu Gly Lys Lys Arg Lys Met Val Lys Val Tyr Thr Lys Thr Asp  
 145 150 155 160  
 Gly Leu Val Ala Val His Pro Lys Ser Val Asn Val Glu Gln Thr Asp  
 165 170 175  
 Phe His Tyr Asn Trp Leu Ile Tyr His Leu Lys Met Arg Thr Ser Ser  
 180 185 190  
 Ile Tyr Leu Tyr Asp Cys Thr Glu Val Ser Pro Tyr Cys Leu Leu Phe  
 195 200 205  
 Phe Gly Gly Asp Ile Ser Ile Gln Lys Asp Asn Asp Gln Glu Thr Ile  
 210 215 220  
 Ala Val Asp Glu Trp Ile Val Phe Gln Ser Pro Ala Arg Ile Ala His  
 225 230 235 240  
 Leu Val Lys Glu Leu Arg Lys Glu Leu Asp Ile Leu Leu Gln Glu Lys  
 245 250 255  
 Ile Glu Ser Pro His Pro Val Asp Trp Asn Asp Thr Lys Ser Arg Asp  
 260 265 270  
 Cys Ala Val Leu Ser Ala Ile Ile Asp Leu Ile Lys Thr Gln Glu Lys  
 275 280 285  
 Ala Thr Pro Arg Asn Phe Pro Pro Arg Phe Gln Asp Gly Tyr Tyr Ser  
 290 295 300

<210> 101  
 <211> 220  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> unsure  
 <222> (139)

<220>  
 <221> unsure  
 <222> (208)..(209)

<400> 101  
 cagaagctgt accacaagaa gatcttccgg actgccatgc tgttccagtt tgtgaacgtg 60  
 ctgctccagg tctgtgtcca caagtcccat gatcttctgc aggaggagat tggcatcgcc 120  
 atctacaaca tggcctcant cgactttgat ggcttctttg ccgccttcct cccagagttc 180  
 ctgaccagct gtgatggtgt ggatgccnnc cagaaaagtt 220

<210> 102  
 <211> 61  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> UNSURE  
 <222> (35)

<220>  
 <221> UNSURE  
 <222> (58)

<400> 102  
 Met Leu Phe Gln Phe Val Asn Val Leu Leu Gln Val Leu Val His Lys  
           1                  5                  10                  15  
 Ser His Asp Leu Leu Gln Glu Glu Ile Gly Ile Ala Ile Tyr Asn Met  
                   20                          25                          30  
 Ala Ser Xaa Asp Phe Asp Gly Phe Phe Ala Ala Phe Leu Pro Glu Phe  
                   35                          40                          45  
 Leu Thr Ser Cys Asp Gly Val Asp Ala Xaa Gln Lys Ser  
           50                          55                          60

<210> 103  
 <211> 251  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> unsure  
 <222> (17)

<220>  
 <221> unsure  
 <222> (72)

<220>  
 <221> unsure  
 <222> (77)

<220>  
 <221> unsure  
 <222> (100)

<220>  
 <221> unsure  
 <222> (108)

<220>  
 <221> unsure  
 <222> (132)

<220>  
 <221> unsure  
 <222> (134)

<220>  
 <221> unsure  
 <222> (171)

<220>  
 <221> unsure  
 <222> (179)

<220>  
 <221> unsure  
 <222> (227)

<400> 103  
 acgacagtgc caccttntca ccattccagc caaggagaga tgtgacgttg gaactgcttt 60  
 ggcaattttg tnaagcntcc cccgccccaa ttgccttgan atttttgntt ttgtgcagag 120  
 atttgcaaaag antnaagttt ttgttgtttt ttcattcattc cattgtgata ntaagaaant 180  
 aagaagctta atgaaaagaa ataaaatgcc tatgttggtg ttttagnaaa aaaaaaaaaa 240  
 aaaaaaaaaa a 251

<210> 104  
 <211> 422  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> unsure  
 <222> (9)

<220>  
 <221> unsure  
 <222> (11)

<220>  
 <221> unsure  
 <222> (355)

<400> 104  
 gaattcggna ngaggcttga ckggactttc tattaretca actccaccag ctgtcagtag 60  
 tgttctcagt acaggtgtac caacagtacc gttattgcca ccacaagtaa accagtccct 120  
 cacttctgtg ccaccaatga atccagctac tacattacca ggtctgatgc ctttaccagc 180  
 aggactgccc aacctcccca acctcaacct caacctccca gcaccacaca tcatgccagg 240  
 ggttggctta ccagaacttg taaacccagg tctgccacct cttcettcca tgcctccccg 300  
 aaacttacct gggcattgca cctcttcccc ctggccatcc gagttcctcc cgttnatttc 360  
 ccttggggtt ccagaggagg ttttttttgc aggcaaggtt caggagagat tgggtggttt 420  
 ta 422

<210> 105  
 <211> 140  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> UNSURE  
 <222> (3) .. (4)

<220>  
 <221> UNSURE  
 <222> (12)

<220>  
 <221> UNSURE  
 <222> (118)

<400> 105

Asn Ser Xaa Xaa Gly Leu Thr Gly Leu Ser Ile Xaa Ser Thr Pro Pro  
 1 5 10 15  
 Ala Val Ser Ser Val Leu Ser Thr Gly Val Pro Thr Val Pro Leu Leu  
 20 25 30  
 Pro Pro Gln Val Asn Gln Ser Leu Thr Ser Val Pro Pro Met Asn Pro  
 35 40 45  
 Ala Thr Thr Leu Pro Gly Leu Met Pro Leu Pro Ala Gly Leu Pro Asn  
 50 55 60  
 Leu Pro Asn Leu Asn Leu Asn Leu Pro Ala Pro His Ile Met Pro Gly  
 65 70 75 80  
 Val Gly Leu Pro Glu Leu Val Asn Pro Gly Leu Pro Pro Leu Pro Ser  
 85 90 95  
 Met Pro Pro Arg Asn Leu Pro Gly His Cys Thr Ser Ser Pro Trp Pro  
 100 105 110  
 Ser Glu Phe Leu Pro Xaa Ile Ser Leu Gly Phe Pro Glu Glu Val Phe  
 115 120 125  
 Phe Ala Gly Lys Val Gln Gly Glu Leu Val Val Phe  
 130 135 140

<210> 106  
 <211> 328  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> unsure  
 <222> (76)

<400> 106  
 caggttgagt ggggctcaca cgctagggtg agatgtcaga aagcgcttgt attttaaaca 60  
 accaaaaaga attgtngggg tggcttgctg ccaggcttgc actgccgttc ctgggggtgt 120  
 gcatcttcgg gaaaggtggt ggcggggcgt ccactagggtt tcctgtcccc tgctgtcct 180  
 tccgtaagaa aatgaaatat tctatgccta atactcacac gcaacatttc ttgtactttg 240  
 taagtcgttt gcgagaatgc agaccacctc actaaactgt aaacggtaaa gagattttta 300  
 cttttggtca aaaaaaaaaa aaaaaaaa 328

<210> 107  
 <211> 896  
 <212> DNA  
 <213> Homo sapiens

<400> 107  
 cagcaggcgg tggactgcaa ggaccagat gatgtggtac cagttggcca aagaagagcc 60  
 tgggtgttgt gcatgtgctt tggactagca tttatgcttg caggtgttat tctaggagga 120  
 gcatacttgt acaaatatct tgcacttcaa ccagatgacg tgtactactg tggataaag 180  
 tacatcaaag atgatgtcat cttaaatgag ccctctgcag atgccccagc tgctctctac 240  
 cagacaattg aagaaaatat taaaatcttt gaagaagaag aagttgaatt tatcagtgtg 300  
 cctgtcccag agtttgcaga tagtgatcct gccaacattg ttcattgact taacaagaaa 360  
 cttacagcct atttagatct taacctggat aagtgtctat tgatccctct gaacacttcc 420  
 attgttatgc caccagaaa cctactggag ttacttatta acatcaaggc tggaaacctat 480  
 ttgcctcagt cctatctgat tcatgagcac atggttatta ctgatcgat tgaaaacatt 540

gatcacctgg gtttctttat ttatcgactg tgcctatgaca aggaaactta caaactgcaa 600  
 cgcagagaaa ctattaaagg tattcagaaa cgtgaagcca gcaattgttt cgcaattcgg 660  
 cattttgaaa acaaatttgc cgtggaaact ttaatttggtt cttgaacagt caagaaaaac 720  
 attattgagg aaaattaata tcacagcata accccaccct ttacattttg tgcagtgtatt 780  
 attttttaa gtcttctttc atgtaagtag caaacagggc ttactatct tttcatctca 840  
 ttaattcaat taaaaccatt accttaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaa 896

<210> 108

<211> 210

<212> PRT

<213> Homo sapiens

<400> 108

Met Cys Phe Gly Leu Ala Phe Met Leu Ala Gly Val Ile Leu Gly Gly  
 1 5 10 15

Ala Tyr Leu Tyr Lys Tyr Phe Ala Leu Gln Pro Asp Asp Val Tyr Tyr  
 20 25 30

Cys Gly Ile Lys Tyr Ile Lys Asp Asp Val Ile Leu Asn Glu Pro Ser  
 35 40 45

Ala Asp Ala Pro Ala Ala Leu Tyr Gln Thr Ile Glu Glu Asn Ile Lys  
 50 55 60

Ile Phe Glu Glu Glu Glu Val Glu Phe Ile Ser Val Pro Val Pro Glu  
 65 70 75 80

Phe Ala Asp Ser Asp Pro Ala Asn Ile Val His Asp Phe Asn Lys Lys  
 85 90 95

Leu Thr Ala Tyr Leu Asp Leu Asn Leu Asp Lys Cys Tyr Val Ile Pro  
 100 105 110

Leu Asn Thr Ser Ile Val Met Pro Pro Arg Asn Leu Leu Glu Leu Leu  
 115 120 125

Ile Asn Ile Lys Ala Gly Thr Tyr Leu Pro Gln Ser Tyr Leu Ile His  
 130 135 140

Glu His Met Val Ile Thr Asp Arg Ile Glu Asn Ile Asp His Leu Gly  
 145 150 155 160

Phe Phe Ile Tyr Arg Leu Cys His Asp Lys Glu Thr Tyr Lys Leu Gln  
 165 170 175

Arg Arg Glu Thr Ile Lys Gly Ile Gln Lys Arg Glu Ala Ser Asn Cys  
 180 185 190

Phe Ala Ile Arg His Phe Glu Asn Lys Phe Ala Val Glu Thr Leu Ile  
 195 200 205

Cys Ser  
 210

<210> 109

<211> 268

<212> DNA

<213> Homo sapiens

&lt;400&gt; 109

gtttgacctg gctggaataa cgtgtgggca cttccttgaa cttttctgga ccttctttgg 60  
 tgcaaccctg attgggaaag caatcattaa aatgcataac cagaaaaat ttgttatagt 120  
 aactttcagc aagcacatcg tggagcagat ggtgactttc attggtgctg tccccggcat 180  
 aggtccgtct ctgcagaagc cttttcaaga gtacctggag gcgcagcggc agaagcttca 240  
 tcacagaagt gaagcgggca caccgcag 268

&lt;210&gt; 110

&lt;211&gt; 59

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 110

Met His Ile Gln Lys Ile Phe Val Ile Val Thr Phe Ser Lys His Ile  
 1 5 10 15

Val Glu Gln Met Val Thr Phe Ile Gly Ala Val Pro Gly Ile Gly Pro  
 20 25 30

Ser Leu Gln Lys Pro Phe Gln Glu Tyr Leu Glu Ala Gln Arg Gln Lys  
 35 40 45

Leu His His Arg Ser Glu Ala Gly Thr Pro Gln  
 50 55

&lt;210&gt; 111

&lt;211&gt; 138

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (33)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (44)

&lt;400&gt; 111

gagacagtat aaggaaaatc tgggttggtgt ctnacaagtg agcngacacc attttttatt 60  
 ctgtgtattt agaatgaagt cttgaaaaaa acttaaaaaa gacaacttta atcattccaa 120  
 aaaaaaaaaa aaaaaaaaaa 138

&lt;210&gt; 112

&lt;211&gt; 415

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (211)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (225)

&lt;220&gt;

&lt;221&gt; unsure



<222> (234)

<220>

<221> unsure

<222> (238)

<220>

<221> unsure

<222> (243)

<220>

<221> unsure

<222> (251)

<220>

<221> unsure

<222> (255)

<220>

<221> unsure

<222> (262)

<220>

<221> unsure

<222> (269)

<220>

<221> unsure

<222> (274)

<220>

<221> unsure

<222> (285)

<220>

<221> unsure

<222> (289)

<220>

<221> unsure

<222> (298)

<220>

<221> unsure

<222> (300)

<220>

<221> unsure

<222> (320)

<220>

<221> unsure

<222> (323)

<220>

<221> unsure

<222> (337)

<220>

<221> unsure

<222> (350)

<220>

<221> unsure

<222> (353)

<220>

<221> unsure

<222> (355)

<220>

<221> unsure

<222> (378)

<220>

<221> unsure

<222> (382)

<220>

<221> unsure

<222> (393) .. (394)

<220>

<221> unsure

<222> (403)

<220>

<221> unsure

<222> (405)

<220>

<221> unsure

<222> (413)

<400> 112

```

aacagagaaa gaaaccaccc aagagtatat cagaatcggg atttccgagg tcacaacaga 60
ggctatagaa ggccctatta ttcccggtgg cgtaacagag gcttttatcc atggggccaa 120
tataaccgag gaggctatgg aaactaccgc tcaaattggc agaattaccg gcaagcatac 180
agtccctcgtc gaggccgttc aagatcccg nccccaaaaa aaagntcccc tccnccangg 240
tcnagaaccc ntcnnaaaac cnctaataant tctnctccta accgntcang gccccccncc 300
cccccccttc ctccccccan ccntacccaa tttaatnctc ctaaccccan ttntncaaag 360
aaaaaaaaatt cccctccnaa gnatacccg cennctcagg ctncngggaa tance 415

```

<210> 113

<211> 92

<212> PRT

<213> Homo sapiens

<220>

<221> UNSURE

<222> (30)

<220>

<221> UNSURE

<222> (33) .. (34)

<220>

<221> UNSURE

<222> (36)

<220>  
 <221> UNSURE  
 <222> (39)..(40)

<220>  
 <221> UNSURE  
 <222> (45)

<220>  
 <221> UNSURE  
 <222> (50)..(51)

<220>  
 <221> UNSURE  
 <222> (55)

<220>  
 <221> UNSURE  
 <222> (62)..(63)

<220>  
 <221> UNSURE  
 <222> (67)

<220>  
 <221> UNSURE  
 <222> (72)..(73)

<220>  
 <221> UNSURE  
 <222> (81)..(82)

<220>  
 <221> UNSURE  
 <222> (86)

<220>  
 <221> UNSURE  
 <222> (90)

<400> 113  
 Met Glu Thr Thr Ala Gln Ile Gly Arg Ile Thr Gly Lys His Thr Val  
   1                  5                  10                  15  
 Leu Val Glu Ala Val Gln Asp Pro Gly Pro Gln Lys Lys Xaa Pro Leu  
                   20                  25                  30  
 Xaa Xaa Gly Xaa Glu Pro Xaa Xaa Lys Pro Leu Ile Xaa Leu Leu Leu  
           35                  40                  45  
 Thr Xaa Xaa Gly Pro Pro Xaa Pro Pro Phe Leu Pro Pro Xaa Xaa Pro  
   50                  55                  60  
 Asn Leu Xaa Leu Leu Thr Pro Xaa Xaa Gln Arg Lys Lys Ile Pro Leu  
   65                  70                  75                  80  
 Xaa Xaa Ile Pro Gly Xaa Leu Arg Leu Xaa Gly Ile  
                   85                  90

<210> 114  
<211> 268  
<212> DNA  
<213> Homo sapiens

<220>  
<221> unsure  
<222> (37)

<220>  
<221> unsure  
<222> (71)

<220>  
<221> unsure  
<222> (73)

<220>  
<221> unsure  
<222> (88)

<220>  
<221> unsure  
<222> (99)

<220>  
<221> unsure  
<222> (103)

<220>  
<221> unsure  
<222> (106)

<220>  
<221> unsure  
<222> (122)

<220>  
<221> unsure  
<222> (127)

<220>  
<221> unsure  
<222> (132)

<220>  
<221> unsure  
<222> (134)

<220>  
<221> unsure  
<222> (142)

<220>  
<221> unsure  
<222> (146)

<220>  
<221> unsure  
<222> (151)

<220>  
<221> unsure  
<222> (158)

<220>  
<221> unsure  
<222> (165)

<220>  
<221> unsure  
<222> (168)

<220>  
<221> unsure  
<222> (179)

<220>  
<221> unsure  
<222> (184)

<220>  
<221> unsure  
<222> (194)

<220>  
<221> unsure  
<222> (223)

<220>  
<221> unsure  
<222> (232)

<220>  
<221> unsure  
<222> (240)

<220>  
<221> unsure  
<222> (248)

<220>  
<221> unsure  
<222> (250)

<400> 114  
aatattgccga gtggtgccgg gtatcagttt gggaaanacc aaggtcagtt tgaccatggt 60  
tttgggtccc ngngtccatc caaaaagngc cctgtgggna agngtncacc atccaatggg 120  
tncaaanatg gntnatttca gnagngggag ngtgctgntt caggngggngc agcctatana 180  
aagnggtatt tagnagagca gaagacagag gatgggaaag atnagggaca gnaacaaacn 240  
aataccgntn aaaaaaaaaa aaaaaaaaaa 268

<210> 115  
<211> 323  
<212> DNA  
<213> Homo sapiens

<220>  
<221> unsure  
<222> (245)..(247)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (275)

&lt;400&gt; 115

```

acagagcctc tacctggcag gaaacaagtc cgggatactt tggcagcaat ctcagaagtt 60
ctttatgttg atttgctaga aggggataca gaatgccatg ctagatttaa aactcctgag 120
gatgctcaag cagtaataaa tgcctataca gaaattaaca agaaacactg ctggaaactc 180
gagatccttt ctggtgatca cgaacaaagg tattggcaga agattttggt tgatagaaag 240
gcaannntta atcagcctcg ggaaaagaaa agagnngtga aaagttaatc accagagctg 300
aaaagattag actggcaaag act                                     323

```

&lt;210&gt; 116

&lt;211&gt; 95

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; UNSURE

&lt;222&gt; (82)..(83)

&lt;220&gt;

&lt;221&gt; UNSURE

&lt;222&gt; (92)

&lt;400&gt; 116

```

Thr Glu Pro Leu Pro Gly Arg Lys Gln Val Arg Asp Thr Leu Ala Ala
  1              5              10              15

Ile Ser Glu Val Leu Tyr Val Asp Leu Leu Glu Gly Asp Thr Glu Cys
          20              25              30

His Ala Arg Phe Lys Thr Pro Glu Asp Ala Gln Ala Val Ile Asn Ala
          35              40              45

Tyr Thr Glu Ile Asn Lys Lys His Cys Trp Lys Leu Glu Ile Leu Ser
          50              55              60

Gly Asp His Glu Gln Arg Tyr Trp Gln Lys Ile Leu Val Asp Arg Lys
          65              70              75              80

Ala Xaa Xaa Asn Gln Pro Arg Glu Lys Lys Arg Xaa Val Lys Ser
          85              90              95

```

&lt;210&gt; 117

&lt;211&gt; 190

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (15)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (18)

&lt;220&gt;

WO 00/37630

PCT/US99/31005

<221> unsure  
<222> (31)

<220>  
<221> unsure  
<222> (40)

<220>  
<221> unsure  
<222> (58)

<220>  
<221> unsure  
<222> (62)

<220>  
<221> unsure  
<222> (65)

<220>  
<221> unsure  
<222> (70)

<220>  
<221> unsure  
<222> (72)

<220>  
<221> unsure  
<222> (74)

<220>  
<221> unsure  
<222> (80)

<220>  
<221> unsure  
<222> (93)

<220>  
<221> unsure  
<222> (120)

<220>  
<221> unsure  
<222> (124)

<220>  
<221> unsure  
<222> (133)

<220>  
<221> unsure  
<222> (149)

<220>  
<221> unsure  
<222> (161)

<400> 117

tttttaatta aaagnaanat ttttgttccct naaattgtan ataagaattt ttttttagnga 60  
 cnaanatgan gnanaccacn atttttttta aanattttat ttgttgaaat tattttagan 120  
 gtcngtgtca gnggatttag taaataaang tgttttggac nttaaaaaa aaaaaaaaaa 180  
 aaaaaaaaaa 190

<210> 118  
 <211> 294  
 <212> DNA  
 <213> Homo sapiens

<400> 118  
 ggcattctgca acctgctcct ttacttcgcc ttctacatca tcatgaagct ccggagtggg 60  
 gagaggatca agctcatccc cctgctctgc atcgtttgca cctccgtggt ctggggcttc 120  
 gcgctcttct tcttcttcca gggactcagc acctggcaga aaaccctgc agagtcgagg 180  
 ggcacaacc gggactgcat cctcctcgac ttctttgacg accacgacat ctggcacttc 240  
 ctctctcca tcgccatgtt tcgggtcctt cctgggtgtt gctgacactg gatg 294

<210> 119  
 <211> 80  
 <212> PRT  
 <213> Homo sapiens

<400> 119  
 Met Lys Leu Arg Ser Gly Glu Arg Ile Lys Leu Ile Pro Leu Leu Cys  
 1 5 10 15  
 Ile Val Cys Thr Ser Val Val Trp Gly Phe Ala Leu Phe Phe Phe Phe  
 20 25 30  
 Gln Gly Leu Ser Thr Trp Gln Lys Thr Pro Ala Glu Ser Arg Glu His  
 35 40 45  
 Asn Arg Asp Cys Ile Leu Leu Asp Phe Phe Asp Asp His Asp Ile Trp  
 50 55 60  
 His Phe Leu Ser Ser Ile Ala Met Phe Arg Val Leu Pro Gly Val Cys  
 65 70 75 80

<210> 120  
 <211> 230  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> unsure  
 <222> (38)

<220>  
 <221> unsure  
 <222> (46)

<400> 120  
 accccagatg ctgaggatgg gggagctcag gcggggcctc tgcttngggg atgggaatgt 60  
 gttttttccc caaacttggt tttatagctc tgcttgaagg gctgggagat gaggtgggtc 120  
 tggatctttt ctccagcgt ctccatgcta tggttgcatt tccgttttct atgaatgaat 180  
 ttgcattcaa taaacaacca gactcaaaaa aaaaaaaaaa aaaaaaaaaa 230

<210> 121  
 <211> 495



<212> DNA  
<213> Homo sapiens

<220>  
<221> unsure  
<222> (429)

<220>  
<221> unsure  
<222> (467)

<400> 121  
gacctgcctt cctgctcttc taggtagtca cacttcacta aagtgtcatc caccagtgtg 60  
ttgaatccga agaatgacaa ttttctacca ctggtgtaaa aaacaaacat ttgaagaccc 120  
ttgtgcattg tgtgtcacaa agctaaatac atggaaatcg ttaatatcgc tgatattaag 180  
taatttcccc actctgagtg aatactttga tgattgccaa cagtggctaa taaaatgacg 240  
gctaccacac tcatgggtca ctggggctgc gcagggctct ttgaggtggg tggcttcttt 300  
tggaagtac tatgaacgtc tcgaagcagt attctagtga taagaattct taacatagcc 360  
aagcgcccca cgtttgttcc ccacgtttgt tccccctttc tgtttgaaaa acctgttctg 420  
gtagctccnc aagagagatg atactgactt tttaaatttt ttacaanagt ctgtattcct 480  
gatatgccta ttttt 495

<210> 122  
<211> 41  
<212> PRT  
<213> Homo sapiens

<400> 122  
Thr Ser Arg Ser Ser Ile Leu Val Ile Arg Ile Leu Asn Ile Ala Lys  
1 5 10 15

Arg Pro Thr Phe Val Pro His Val Cys Ser Pro Phe Leu Phe Glu Lys  
20 25 30

Pro Val Leu Val Ala Pro Gln Glu Arg  
35 40

<210> 123  
<211> 18  
<212> DNA  
<213> Homo sapiens

<400> 123  
gaaaaaaaaa aaaaaaaaa 18

<210> 124  
<211> 285  
<212> DNA  
<213> Homo sapiens

<220>  
<221> unsure  
<222> (135)

<220>  
<221> unsure  
<222> (234)

<400> 124

cacgaagggt ttcaagggtct gtcttagttc tcattctcaa gattgtttcc agttgcaagt 60  
 tagaggcaag ccagctagct gccagcctt aactctgttc agtgccctgt tactaacatt 120  
 ttttaacaga ttggnnttcta catgtttaaa gtatccagcg ttggatttta cctcttgcta 180  
 gttccatttg tccctgggtgc tgctttttaa ggtaggggc cctgtgaagt ggantatgta 240  
 cgcagttggc ctgggtgatgt atctgtgcct gttttatctt ctccc 285

&lt;210&gt; 125

&lt;211&gt; 48

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; UNSURE

&lt;222&gt; (31)

&lt;400&gt; 125

Met Phe Lys Val Ser Ser Val Gly Phe Tyr Leu Leu Leu Val Pro Phe  
 1 5 10 15

Val Pro Gly Ala Ala Phe Lys Gly Ile Gly Pro Cys Glu Val Xaa Tyr  
 20 25 30

Val Arg Ser Trp Pro Gly Asp Val Ser Val Pro Val Leu Ser Ser Pro  
 35 40 45

&lt;210&gt; 126

&lt;211&gt; 350

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (5)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (215)

&lt;400&gt; 126

ttcctatgt aagatgtcat actgcagatt taaaatatag actatcaata aaatgcatga 60  
 agtgatcatt tgtgcttgat catctctcct tgggtttttc tttaaaaagg ggaatctgct 120  
 ataaagggtt tggtgcttca aaccaatgtc aaatagactt gattttttaga gtcattggaat 180  
 tacagtgcga ccttgatttt tattcccctc actgntatga gtgtgggcag gtactgggtt 240  
 atatgttata acttccgttt tatctgtgtt gtgtagttag atggcttaac cgttgagtgg 300  
 taaaataaaa gattatattc caatacaagg aaaaaaaaaa aaaaaaaaaa 350

&lt;210&gt; 127

&lt;211&gt; 517

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (63)

&lt;400&gt; 127

cacgaggcct cgtgccaaaca ggaaagttgc tttgttttgc ttcgagatgg ctgcggggat 60  
 gtntttggaa cattatctgg acagtattga aaacctccc tttgaattac agagaaactt 120  
 tcagctcatg agggacctag accaaaggac agaggacctg aaggctgaaa ttgacaagtt 180

ggccactgaa tatatgagta gcgcccgcag cctgagctcc gaggagaagc tggcccttct 240  
 cagacagatc caggaggcct atggcaagtg caaggaattt ggtgacgaca aggtgcagct 300  
 ggccatgcag acctatgaga tggtagacaa acacattcgg cggctggaca cagacctggc 360  
 ccgttttgag gctgatctga aggagaaaca gatcgagtcc agtgactatg acagctcttc 420  
 tagcaaaggc aaaaagagcc ggacccaaaa ggagaaaaaa gctgccagag cccgttccaa 480  
 agggaaaaac tcagatgaag aagcccccaa ggctgcc 517

&lt;210&gt; 128

&lt;211&gt; 157

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; UNSURE

&lt;222&gt; (6)

&lt;400&gt; 128

Met Ala Ala Gly Met Xaa Leu Glu His Tyr Leu Asp Ser Ile Glu Asn  
 1 5 10 15

Leu Pro Phe Glu Leu Gln Arg Asn Phe Gln Leu Met Arg Asp Leu Asp  
 20 25 30

Gln Arg Thr Glu Asp Leu Lys Ala Glu Ile Asp Lys Leu Ala Thr Glu  
 35 40 45

Tyr Met Ser Ser Ala Arg Ser Leu Ser Ser Glu Glu Lys Leu Ala Leu  
 50 55 60

Leu Arg Gln Ile Gln Glu Ala Tyr Gly Lys Cys Lys Glu Phe Gly Asp  
 65 70 75 80

Asp Lys Val Gln Leu Ala Met Gln Thr Tyr Glu Met Val Asp Lys His  
 85 90 95

Ile Arg Arg Leu Asp Thr Asp Leu Ala Arg Phe Glu Ala Asp Leu Lys  
 100 105 110

Glu Lys Gln Ile Glu Ser Ser Asp Tyr Asp Ser Ser Ser Ser Lys Gly  
 115 120 125

Lys Lys Ser Arg Thr Gln Lys Glu Lys Lys Ala Ala Arg Ala Arg Ser  
 130 135 140

Lys Gly Lys Asn Ser Asp Glu Glu Ala Pro Lys Ala Ala  
 145 150 155

&lt;210&gt; 129

&lt;211&gt; 246

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (24)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (27)..(28)

<220>  
<221> unsure  
<222> (30)

<220>  
<221> unsure  
<222> (33)

<220>  
<221> unsure  
<222> (87)

<220>  
<221> unsure  
<222> (122)

<400> 129  
tcctgtggtg agggctaggt gtgntcnnch ctnctattct ccattccctt cctgcttttt 60  
tcattggtggg ggatccacca ggtcatntag gctctggccc tagttgaagg ggcacccctt 120  
cncctgtgcc aagaggattc atcctgggag aggggggcaag gtggaatgca gataactcac 180  
atgtaaaagg aacttgggta ggtaaataaa agctatacat gttgaaaaaa aaaaaaaaaa 240  
aaaaaa 246

<210> 130  
<211> 694  
<212> DNA  
<213> Homo sapiens

<220>  
<221> unsure  
<222> (17)

<220>  
<221> unsure  
<222> (62)

<220>  
<221> unsure  
<222> (363)

<220>  
<221> unsure  
<222> (368)

<220>  
<221> unsure  
<222> (413)

<220>  
<221> unsure  
<222> (416)

<220>  
<221> unsure  
<222> (424)

<220>  
<221> unsure  
<222> (465)

<220>  
 <221> unsure  
 <222> (473)

<220>  
 <221> unsure  
 <222> (495)

<220>  
 <221> unsure  
 <222> (499)

<220>  
 <221> unsure  
 <222> (522)

<220>  
 <221> unsure  
 <222> (548)

<220>  
 <221> unsure  
 <222> (566)

<220>  
 <221> unsure  
 <222> (584)

<220>  
 <221> unsure  
 <222> (620)

<220>  
 <221> unsure  
 <222> (651) .. (652)

<220>  
 <221> unsure  
 <222> (667)

<220>  
 <221> unsure  
 <222> (679)

<400> 130  
 aagcttggca cgaggggncaa acctctatgg atatataaag ggaagcttga ggaggaattt 60  
 cncagttaca gtgcagaagc acaagcaaaa gaattaacca gctcttcagt caagcaaatc 120  
 ctctactcac catgcttctt cctgccattc atttctatct ccttcccctt gcatgcatcc 180  
 taatgaaaag ctgttttggt tttaaaaatg atgccacaga aatcctttat tcacatgtgg 240  
 ttaaacctgt tccagcacac cccagcagca acagcacgtt gaatcaagcc agaaatggtt 300  
 gcaggcattt cagtaacact ggactggatc ggaacactcg ggttcaagtg gggtgccggg 360  
 aankgcgntc ccaccaata catctctgat ggccagtgc cagcatcag ccntangaag 420  
 gagntgggtg gtgctggcga gtgacttgcc cctgccagt ctcentaatt ggnttgagg 480  
 aggcgtgtga acaangtant ggagcaggag gagctcccag gngtggcggg gtgtcaatga 540  
 caaaaccngt acccagagaa tccagntgca gttccaagat ggcngcacac gcacgtacaa 600  
 aatcacagta gtcggtgcn gcaagtgcaa gaggtacac cggcagcaca nngagtcag 660  
 tcacganttt gagagcatnt cacgtgcca gcca 694

<210> 131

WO 00/37630

PCT/US99/31005

<211> 102  
<212> PRT  
<213> Homo sapiens

<220>  
<221> UNSURE  
<222> (9)..(10)

<220>  
<221> UNSURE  
<222> (12)

<220>  
<221> UNSURE  
<222> (26)

<220>  
<221> UNSURE  
<222> (29)

<220>  
<221> UNSURE  
<222> (36)..(37)

<220>  
<221> UNSURE  
<222> (45)

<220>  
<221> UNSURE  
<222> (54)

<220>  
<221> UNSURE  
<222> (60)

<220>  
<221> UNSURE  
<222> (66)

<220>  
<221> UNSURE  
<222> (78)

<220>  
<221> UNSURE  
<222> (88)

<220>  
<221> UNSURE  
<222> (93)

<220>  
<221> UNSURE  
<222> (97)

<400> 131  
Met Ala Ser Ala Pro Ala Ser Ala Xaa Xaa Arg Xaa Trp Cys Val Leu  
1 5 10 15

Ala Ser Asp Leu Pro Leu Pro Val Leu Xaa Asn Trp Xaa Gly Gly Gly  
 20 25 30

Cys Gly Thr Xaa Xaa Trp Ser Arg Arg Ser Ser Gln Xaa Trp Arg Cys  
 35 40 45

Val Asn Asp Lys Thr Xaa Thr Gln Arg Ile Gln Xaa Gln Phe Gln Asp  
 50 55 60

Gly Xaa Thr Arg Thr Tyr Lys Ile Thr Val Val Gly Ala Xaa Lys Cys  
 65 70 75 80

Lys Arg Tyr Thr Arg Gln His Xaa Glu Ser Ser His Xaa Phe Glu Ser  
 85 90 95

Xaa Ser Arg Ala Lys Pro  
 100

<210> 132  
 <211> 243  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> unsure  
 <222> (53)

<400> 132  
 atatgaaata catgtttag atatgtaaaa tgaatatattt agtctccctt atnacatata 60  
 tgttcattggt gaactttatc aatagtatgg atctttttta atcaataaga tgctttgtaa 120  
 agttgaaata agtaatactt tcttgtttaa tctgtgcaat cagaaggtgt cttgaccttc 180  
 aattcaattg gtttctttta acaaaaaata acactgctaa aagttaaaaa aaaaaaaaaa 240  
 aaa 243

<210> 133  
 <211> 1187  
 <212> DNA  
 <213> Homo sapiens

<400> 133  
 tgtcagagtt ggtctgttac tcggtggtgg cggagctctac ggaagccggt ttcgcttcac 60  
 ttttcctggc ttagagcgc tttcccccgt gcgggtgaga gtgcagagac gaaggtgcga 120  
 gatgagcact atgttcgcgg acactctcct catcggtttt atctctgtgt gcacggctct 180  
 gctgcagag ggcataacct gggtcctggt ttacaggaca gacaagtaca agagactgaa 240  
 ggcagaagt gaaaaacaga gtaaaaaatt ggaaaagaag aaggaaacaa taacagagtc 300  
 agctggtcga caacagaaaa agaaaaataga gagacaagaa gagaactga agaataacaa 360  
 cagagatcta tcaatgggtc gaatgaaatc catgtttgct attggctttt gttttactgc 420  
 cctaattgga atgttcaatt ccatatttga tggtagagt gtggcaaagc ttccttttac 480  
 cctcttttct tacatccaag gactgtctca tcgaaatctg ctgggagatg acaccacaga 540  
 ctgttccttc attttcctgt atattctctg tactatgtcg attcgacaga acattcagaa 600  
 gattctcggc cttgccctt caagagccgc caccaagcag gcaggtggat ttcttgcccc 660  
 accacctcct tctgggaagt tctcttgaac tcaagaactc tttattttct atcattcttt 720  
 ctgacacac acacatcaga ctggcaactg tttgttagca agagccatag gtacgcttac 780  
 tacttgggcc tctttctagt ttgaattat ttctaagcct tttgggtatg attagagtga 840  
 aaatggcagc cagcaaacct gatagtgtt ttggtcctag atgattttta tcaaataggt 900  
 ggattgatta gttaagtcca ggtaatgttt atgtaatgaa aaacaaatag catccttctt 960  
 gtttcattta cataagtatt ttctgtggga ccgactctca aggcactgtg tatgccctgc 1020  
 aagttggctg tctatgagca tttagagatt tagaagaaaa atttagtttg ttttaaccctt 1080  
 gtaactgttt gttttgttgt tgtttttttt tcaagccaaa tacatgacat aagatcaata 1140

aagaggccaa atttttagct gttttatgta aaaaaaaaaa aaaaaaa

1187

&lt;210&gt; 134

&lt;211&gt; 188

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 134

Met Ser Thr Met Phe Ala Asp Thr Leu Leu Ile Val Phe Ile Ser Val  
 1 5 10 15

Cys Thr Ala Leu Leu Ala Glu Gly Ile Thr Trp Val Leu Val Tyr Arg  
 20 25 30

Thr Asp Lys Tyr Lys Arg Leu Lys Ala Glu Val Glu Lys Gln Ser Lys  
 35 40 45

Lys Leu Glu Lys Lys Lys Glu Thr Ile Thr Glu Ser Ala Gly Arg Gln  
 50 55 60

Gln Lys Lys Lys Ile Glu Arg Gln Glu Glu Lys Leu Lys Asn Asn Asn  
 65 70 75 80

Arg Asp Leu Ser Met Val Arg Met Lys Ser Met Phe Ala Ile Gly Phe  
 85 90 95

Cys Phe Thr Ala Leu Met Gly Met Phe Asn Ser Ile Phe Asp Gly Arg  
 100 105 110

Val Val Ala Lys Leu Pro Phe Thr Pro Leu Ser Tyr Ile Gln Gly Leu  
 115 120 125

Ser His Arg Asn Leu Leu Gly Asp Asp Thr Thr Asp Cys Ser Phe Ile  
 130 135 140

Phe Leu Tyr Ile Leu Cys Thr Met Ser Ile Arg Gln Asn Ile Gln Lys  
 145 150 155 160

Ile Leu Gly Leu Ala Pro Ser Arg Ala Ala Thr Lys Gln Ala Gly Gly  
 165 170 175

Phe Leu Gly Pro Pro Pro Ser Gly Lys Phe Ser  
 180 185

&lt;210&gt; 135

&lt;211&gt; 1300

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 135

tactgactcg aggccaaagaa ttcggcacga ggcaattttt aatttttggt aatatcaaca 60  
 gcaaaagcct agtgatttgg gagatgtgca acctccctga aaatcttttc tgtttctgga 120  
 gtacttcagg ggtggcctct ggccccagag cctttgccac agtgctccca ccagccccc 180  
 cctcatcctg ctgtttgcag agcctcatct acaggtcccc acgctgcctt ctttactcac 240  
 tctgcgcttg gccgttttgt tatttggtt agtctacatt gggcggaagt ctgtgtgcac 300  
 agagtgggtg ttccttcgag cccctccac tcagagggcc acaccagcg atgccagtga 360  
 aggtggcaca gcctctcttc agtttctcct gactgtgac tcaactgggtt agaattcccc 420  
 tgagagaatt cctcactca cggctccctt tgccagagtc agttcaatca ggtctgatgt 480  
 gagcaattta cacacttgc tcagaaagtc cctcagggtt ttagaggac tgcagggggg 540



```

catccgctgc agactcagcc ttctcttgc gccatcctgc agtgggggtg agcgggcaca 600
ggctgagaac tgcctttggg tgggtggaagc aggtgtcacg gtgcaagtct cccctctcac 660
ccctccccca gcttgagccg tgtcaccccc ctctccctcc agcatgggcc tgtgtctcag 720
gctctcttga aggtggccct gccccggacc ctcttgacag tgccttggtt tgacttgga 780
ctagatggcc atctttccag gcttttggtg cccaagagca gtctgggtgg atggaagtgg 840
ctgtcccttc ctctccagcc cctgcccacc cactggtgga ggtgctaact agcagggacg 900
tggcatagga tgggagctgg gcgtgaggtg cttgggggtc attctttgtc cctcagcttc 960
tcagagtccg gccagccctt gtgttcccg gccccacact ttctctctcc cactgcagt 1020
gaggcaatag tccgggggtg ggcttgccct cctgcccctg attggggact caggaggtga 1080
ggcctggggg gcttcttgc cctccttgc ccactgcct gccccgggc agcacgggag 1140
ggagagcagg gtgagcacgc ttgttggtt cagatgcact ttctgcttgc attgccgtat 1200
ctgtgcgttc cttcatctg gtcttggtt tatggaacac catgttttta gcatgttttt 1260
aaataaaaac ggataaagtg tcaaaaaaaa aaaaaaaaaa 1300

```

&lt;210&gt; 136

&lt;211&gt; 62

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 136

```

Met Cys Asn Leu Pro Glu Asn Leu Phe Cys Phe Trp Ser Thr Ser Gly
  1             5             10            15

```

```

Val Ala Ser Gly Pro Arg Ala Phe Ala Thr Val Leu Pro Pro Ala Pro
      20             25            30

```

```

Thr Ser Ser Val Cys Leu Gln Ser Leu Ile Tyr Arg Ser Pro Arg Cys
    35             40            45

```

```

Leu Leu Tyr Ser Leu Cys Ala Trp Pro Phe Cys Tyr Leu Ala
    50             55            60

```

&lt;210&gt; 137

&lt;211&gt; 1330

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 137

```

aagcttgga cgaggaagc ccatccaggt catgtgctac gactatgaca atgacggggg 60
ccatgacttc atcgcgaggt tccagacctc agtgtcacag atgtgtgagg ctcgagacag 120
cgtcccgtcg gagttcgagt gcatcaaccc caagaagcag aggaagaaga agaactataa 180
aaactcgggc atcatcatcc tgcgatcctg caagataaac cgagactact ctttcttga 240
ctacatcctg ggaggtgcc agtcatggt caccgttga atagacttta cagcctccaa 300
cggaatccc ctgcacctt cctctttgca ctatatcaac cctatgggca ccaacgaata 360
tctgtcggcc atctgggctg ttgggcagat cattcaggac tacgacagtg ataagatgtt 420
tccagctctg ggattcgggg ccagttacc ccagactgg aagtctccca tgagtttgc 480
atcaacttca accccaccaa ccccttctgc tcaggtgttg atggtattgc ccaggcgtac 540
tcagcttgcc tgccccacat ccgcttctac ggtctacca atttctcccc catcgtcaac 600
cacgtggccc ggtttgcggc ccaggccaca caacagcga cggccacgca gtacttcac 660
ctcctcatca tcacggacgg ggtcatcagt gacatggagg agacacggca tgccgtgggtg 720
caggcttcca agctgccc atccatcatc atcgtggcg tgggcaatgc ggacttcgct 780
gccatggagt tcttgatgg ggacagccgc atgctgcgt cccacacggg ggaggaggca 840
gcccgcgata ttgtgcagtt cgttccctt cgagagttcc gcaacgcagc aaaagagacc 900
ttggccaaag ctgtgctggc ggagctgccc caacaagtg tcagtatatt caagcataaa 960
aacctgcccc ccaccaactc ggagcccgcc tgagctccag tgcccagcag cagcatgtca 1020
gctgagcttc ctgcccctcc ccaggaacat gcacgctcac tctgcttcc tgtgggtggc 1080
ctttttttac cgatccccct ttttattttt tacaaccgga cctccacccc caacttcttc 1140
cagcccagct gggcttctct tgttgagtc aactgttgat gcttcaggc caaactggct 1200
tcctctcttc ctctccccc ctttgccatt cttaagtatt gaatgtactt tgtataattt 1260

```

tagtggaatt gttattgaga ataaaaatttt tacaatcata aaaaaaaaaa aaaaaaaaaa 1320  
 aaaaaaaaaa 1330

<210> 138  
 <211> 423  
 <212> PRT  
 <213> Homo sapiens

<400> 138

Met Cys Tyr Asp Tyr Asp Asn Asp Gly Gly His Asp Phe Ile Gly Glu  
 1 5 10 15  
 Phe Gln Thr Ser Val Ser Gln Met Cys Glu Ala Arg Asp Ser Val Pro  
 20 25 30  
 Leu Glu Phe Glu Cys Ile Asn Pro Lys Lys Gln Arg Lys Lys Lys Asn  
 35 40 45  
 Tyr Lys Asn Ser Gly Ile Ile Ile Leu Arg Ser Cys Lys Ile Asn Arg  
 50 55 60  
 Asp Tyr Ser Phe Leu Asp Tyr Ile Leu Gly Gly Cys Gln Leu Met Phe  
 65 70 75 80  
 Thr Val Gly Ile Asp Phe Thr Ala Ser Asn Gly Asn Pro Leu Asp Pro  
 85 90 95  
 Ser Ser Leu His Tyr Ile Asn Pro Met Gly Thr Asn Glu Tyr Leu Ser  
 100 105 110  
 Ala Ile Trp Ala Val Gly Gln Ile Ile Gln Asp Tyr Asp Ser Asp Lys  
 115 120 125  
 Met Phe Pro Ala Leu Gly Phe Gly Ala Gln Leu Pro Pro Asp Trp Lys  
 130 135 140  
 Ser Pro Met Ser Leu Pro Ser Thr Ser Thr Pro Pro Thr Pro Ser Ala  
 145 150 155 160  
 Gln Val Trp Met Val Leu Pro Arg Arg Thr Gln Leu Ala Cys Pro Thr  
 165 170 175  
 Ser Ala Ser Thr Val Leu Pro Ile Ser Pro Pro Ser Ser Thr Thr Trp  
 180 185 190  
 Pro Gly Leu Arg Pro Arg Pro His Asn Ser Gly Arg Pro Arg Ser Thr  
 195 200 205  
 Ser Ser Ser Ser Ser Ser Arg Thr Gly Ser Ser Val Thr Trp Arg Arg  
 210 215 220  
 His Gly Met Pro Trp Cys Arg Leu Pro Ser Cys Pro Cys Pro Ser Ser  
 225 230 235 240  
 Ser Trp Ala Trp Ala Met Arg Thr Ser Leu Pro Trp Ser Ser Trp Met  
 245 250 255  
 Gly Thr Ala Ala Cys Cys Ala Pro Thr Arg Gly Arg Arg Gln Pro Ala  
 260 265 270

Ile Leu Cys Ser Ser Phe Pro Phe Glu Ser Ser Ala Thr Gln Gln Lys  
 275 280 285

Arg Pro Trp Pro Lys Leu Cys Trp Arg Ser Cys Pro Asn Lys Leu Cys  
 290 295 300

Ser Ile Ser Ser Ile Lys Thr Cys Pro Pro Pro Thr Arg Ser Pro Pro  
 305 310 315 320

Glu Leu Gln Cys Pro Ala Ala Ala Cys Gln Leu Ser Leu Leu Pro Ser  
 325 330 335

Pro Arg Asn Met His Ala His Ser Ala Ser Leu Trp Val Ala Phe Phe  
 340 345 350

Tyr Arg Ser Pro Phe Leu Phe Phe Thr Thr Gly Pro Pro Pro Pro Thr  
 355 360 365

Ser Ser Ser Pro Ala Gly Leu Pro Leu Leu Glu Ser Thr Val Asp Ala  
 370 375 380

Ser Arg Pro Asn Trp Leu Pro Leu Leu Leu Ser Pro Pro Leu Pro Phe  
 385 390 395 400

Leu Ser Ile Glu Cys Thr Leu Tyr Asn Phe Ser Gly Ile Val Ile Glu  
 405 410 415

Asn Lys Ile Phe Thr Ile Ile  
 420

&lt;210&gt; 139

&lt;211&gt; 1920

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 139

aagcttgcca cgaggcgcca gacggcgagg cgggcctttt ggcgctccact gcgcggtctgc 60  
 accctgcccc atcctgcccgg gatcatggtc tgcggcgagcc cgggagggat gctgctgctg 120  
 cgggcccgggc tgcttgccct ggctgctctc tgctgctcc ggggtgcccgg ggctcgggct 180  
 gcagcctgtg agcccgctcc catccccctg tgcaagtccc tgccctggaa catgactaag 240  
 atgcccaccc acctgcacca cagcactcag gccaacgcca tcctggccat cgagcagttc 300  
 gaaggtctgc tgggcaccca ctgcagcccc gatctgctct tcttctctctg tgccatgtac 360  
 gcgcccactc gcaccattga cttccagcac gagcccatca agccctgtaa gtctgtgtgc 420  
 gagcggggccc ggcagggctg tgagcccata ctcatcaagt accgccactc gtggccggag 480  
 aacctggcct gcgaggagct gccagtgtac gacaggggag tgtgcatctc tcccaggagg 540  
 atcgttactg cggacggagc tgattttcct atggattcta gtaacggaaa ctgtagaggg 600  
 gcaagcagtg aacgctgtaa atgtaagcct attagagcta cacagaagac ctatttccgg 660  
 aacaattaca actatgtcat tcgggctaaa gttaaagaga taaagactaa gtgccatgat 720  
 gtgactgcag tagtgagggt gaaggagatt ctaaagtcct ctctggtaaa cattccacag 780  
 gacactgtca acctctatac cagctctggc tgccctctgcc ctccacttaa tgtaattgag 840  
 gaatatatca tcatgggcta tgaagatgag gaacgttcca gattactctt ggtggaaggc 900  
 tctatagctg agaagtggaa ggatcgactc ggtaaaaaag ttaagcgtg ggatatgaag 960  
 cttcgtctac ttggactcag taaaagtgat tctagcaata gtgattccac tcagagtcag 1020  
 aagttctgcca ggaactcgaa cccccggcaa gcacgcaact aaatcccgaa atacaaaaag 1080  
 taacacagtg gacttctat taagacttac ttgcattgct ggactagcaa aggaaaattg 1140  
 cactattgca catcatattc tattgtttac tataaaaaac atgtgataac tgattattac 1200  
 ttctgtttct cttttgggtt ctgcttctct cttctctcaa cccctttgta atggtttggg 1260  
 ggcagactct taagtatat gtgagttttc tatttacta atcatgagaa aaactgttct 1320  
 ttgcaataa taataaatta aacatgctgt taccagagcc tctttgctgg agtctccaga 1380

tgttaattta ctttctgcac cccaattggg aatgcaatat tggatgaaaa gagagggttc 1440  
 tggatttcac agaaagctag atatgcctta aaacatactc tgccgatcta attacagcct 1500  
 tatttttgta tgccttttgg gcattctcct catgcttaga aagttccaaa tgtttataaa 1560  
 ggtaaaatgg cagtttgaag tcaaatgtca cataggcaaa gcaatcaagc accaggaagt 1620  
 gtttatgagg aaacaacacc caagatgaat tatttttgag actgtcagga agtaaaataa 1680  
 ataggagctt aagaagaac attttgcctg attgagaagc acaactgaaa ccagtagccg 1740  
 ctgggggtgtt aatggtagca ttcttctttt ggcaatacat ttgatttggt catgaatata 1800  
 ttaatcagca ttagagaaat gaattataac tagacatctg ctgttatcac catagttttg 1860  
 ttttaattgc ttctttttaa ataaacccat tggtgaaagt caaaaaaaaa aaaaaaaaaa 1920

&lt;210&gt; 140

&lt;211&gt; 325

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 140

Met Val Cys Gly Ser Pro Gly Gly Met Leu Leu Leu Arg Ala Gly Leu  
 1 5 10 15

Leu Ala Leu Ala Ala Leu Cys Leu Leu Arg Val Pro Gly Ala Arg Ala  
 20 25 30

Ala Ala Cys Glu Pro Val Arg Ile Pro Leu Cys Lys Ser Leu Pro Trp  
 35 40 45

Asn Met Thr Lys Met Pro Asn His Leu His His Ser Thr Gln Ala Asn  
 50 55 60

Ala Ile Leu Ala Ile Glu Gln Phe Glu Gly Leu Leu Gly Thr His Cys  
 65 70 75 80

Ser Pro Asp Leu Leu Phe Phe Leu Cys Ala Met Tyr Ala Pro Ile Cys  
 85 90 95

Thr Ile Asp Phe Gln His Glu Pro Ile Lys Pro Cys Lys Ser Val Cys  
 100 105 110

Glu Arg Ala Arg Gln Gly Cys Glu Pro Ile Leu Ile Lys Tyr Arg His  
 115 120 125

Ser Trp Pro Glu Asn Leu Ala Cys Glu Glu Leu Pro Val Tyr Asp Arg  
 130 135 140

Gly Val Cys Ile Ser Pro Glu Ala Ile Val Thr Ala Asp Gly Ala Asp  
 145 150 155 160

Phe Pro Met Asp Ser Ser Asn Gly Asn Cys Arg Gly Ala Ser Ser Glu  
 165 170 175

Arg Cys Lys Cys Lys Pro Ile Arg Ala Thr Gln Lys Thr Tyr Phe Arg  
 180 185 190

Asn Asn Tyr Asn Tyr Val Ile Arg Ala Lys Val Lys Glu Ile Lys Thr  
 195 200 205

Lys Cys His Asp Val Thr Ala Val Val Glu Val Lys Glu Ile Leu Lys  
 210 215 220

Ser Ser Leu Val Asn Ile Pro Gln Asp Thr Val Asn Leu Tyr Thr Ser  
 225 230 235 240

Ser Gly Cys Leu Cys Pro Pro Leu Asn Val Asn Glu Glu Tyr Ile Ile  
 245 250 255

Met Gly Tyr Glu Asp Glu Glu Arg Ser Arg Leu Leu Leu Val Glu Gly  
 260 265 270

Ser Ile Ala Glu Lys Trp Lys Asp Arg Leu Gly Lys Lys Val Lys Arg  
 275 280 285

Trp Asp Met Lys Leu Arg His Leu Gly Leu Ser Lys Ser Asp Ser Ser  
 290 295 300

Asn Ser Asp Ser Thr Gln Ser Gln Lys Ser Gly Arg Asn Ser Asn Pro  
 305 310 315 320

Arg Gln Ala Arg Asn  
 325

<210> 141  
 <211> 1469  
 <212> DNA  
 <213> Homo sapiens

<400> 141  
 gatagcttgg cacgaggctg ggaagtagag gtgttgtgct gagcggcgct cggcgaactg 60  
 tgtggaccgt ctgctgggac tccggccctg cgtccgctca gcccctgggc cccgcgcacc 120  
 tactgccatg gagacgcggc ctgctctcgg ggccacctgt ttgctgggct tcagtttctt 180  
 gctcctcgtc atctcttctg atggacataa tgggcttgga aagggttttg gagatcatat 240  
 tcattggagg acactggaag atgggaagaa agaagcagct gccagtggac tgcccttgat 300  
 ggtgattatt cataaatcct ggtgtggagc ttgcaaagct cttaaagccca aatttgca 360  
 atctacggaa atttcagaac tctcccataa tttgtttatg gtaaatcttg aggatgaaga 420  
 ggaacccaaa gatgaagatt tcagccctga cgggggttat attccacgaa tcttttttct 480  
 ggatcccagt ggcaagggtc atcctgaaat catcaatgag aatggaaacc ccagctacaa 540  
 gtatttttat gtcagtgccg agcaagtgtt tcaggggatg aaggaaagctc aggaaggct 600  
 gacgggtgat gccttcagaa agaaacatct tgaagatgaa ttgtaacatg aatgtgcccc 660  
 ttctttcatc agagttagtg ttctggaagg aaagcagcag ggaagggaat attgaggaat 720  
 catctagaac aattaagccg accaggaaac ctcatctcta cctacactgg aaggagcgct 780  
 ctcaactgtg aagagtcttg ctaacagaag ctggtctgca tgtttgtgga tccagcggag 840  
 agtggcagac tttcttctcc ttttccctct cacctaaatg tcaacttgtc attgaatgta 900  
 aagaatgaaa ccttctgaca caaaacttga gccacttgga tgtttactcc tgcacttaa 960  
 gtattttagt cttttcccat ttctccccc tttactcacc ttagtgtgta aaggagacta 1020  
 gtagcatctt ttctacaacg ttaaaattgc agaagtagct tatcattaaa aaacaacaac 1080  
 aacaacaata acaataaatc ctaagtgtaa atcagttatt ctacccccta ccaaggatat 1140  
 cagcctgttt tttccctttt ttctcctggg aataattgtg ggcttcttcc caaatttcta 1200  
 cagcctcttt cctcttctca tgcttgagct tccctgtttg cagcctatgc tgtgcaggac 1260  
 tggctgtgtg cttggactcg gctccagggt gaagcatgct tccctgtgtt actgttggag 1320  
 aaactcaaac cttcaagccc taggtgtagc cattttgtca agtcatcaac tgtatttttg 1380  
 tactggcatt aacaaaaaaa gagataaaat attgtaccat taaactttta taaaacttta 1440  
 aaaggaaaaa aaaaaaaaaa aaaaaaaaaa 1469

<210> 142  
 <211> 172  
 <212> PRT  
 <213> Homo sapiens

<400> 142  
 Met Glu Thr Arg Pro Arg Leu Gly Ala Thr Cys Leu Leu Gly Phe Ser  
 1 5 10 15

Phe Leu Leu Leu Val Ile Ser Ser Asp Gly His Asn Gly Leu Gly Lys  
                   20                  25                  30  
 Gly Phe Gly Asp His Ile His Trp Arg Thr Leu Glu Asp Gly Lys Lys  
           35                  40                  45  
 Glu Ala Ala Ala Ser Gly Leu Pro Leu Met Val Ile Ile His Lys Ser  
           50                  55                  60  
 Trp Cys Gly Ala Cys Lys Ala Leu Lys Pro Lys Phe Ala Glu Ser Thr  
           65                  70                  75                  80  
 Glu Ile Ser Glu Leu Ser His Asn Phe Val Met Val Asn Leu Glu Asp  
                   85                  90                  95  
 Glu Glu Glu Pro Lys Asp Glu Asp Phe Ser Pro Asp Gly Gly Tyr Ile  
                   100                  105                  110  
 Pro Arg Ile Leu Phe Leu Asp Pro Ser Gly Lys Val His Pro Glu Ile  
           115                  120                  125  
 Ile Asn Glu Asn Gly Asn Pro Ser Tyr Lys Tyr Phe Tyr Val Ser Ala  
           130                  135                  140  
 Glu Gln Val Val Gln Gly Met Lys Glu Ala Gln Glu Arg Leu Thr Gly  
           145                  150                  155                  160  
 Asp Ala Phe Arg Lys Lys His Leu Glu Asp Glu Leu  
                   165                  170

&lt;210&gt; 143

&lt;211&gt; 1458

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 143

tctagaacta gtggatcccc cggggtgcag gaattcggca cgagcttcaa gtgaccattc 60  
 tttttcttct gccagctatt tgcagcagta acagcacagg tgtttttagag gcagctaata 120  
 attcacttgt tactacaaca aaacctctta taacaacacc aaacacagaa tcattacaga 180  
 aaaatgttgt cacaccaaca actggaacaa ctctctaaagg aacaatcacc aatgaattac 240  
 ttaaaatgtc tctgatgtca acagctactt ttttaacaag taaagatgaa ggattgaaag 300  
 ccacaaccac tgatgtcagg aagaatgact ccatcatttc aaacgtaaca gtaacaagtg 360  
 ttacacttcc aaatgctgtt tcaacattac aaagttccaa acccaagact gaaactcaga 420  
 gttcaattaa aacaacagaa ataccaggta gtgttctaca accagatgca tcaccttcta 480  
 aaactggtac attaacctca ataccagtta caattccaga aaacacctca cagtctcaag 540  
 taataggcac tgagggtgga aaaaatgcaa gcacttcagc aaccagccgg tcttattcca 600  
 gtattatatt gccgggtggt attgctttga ttgtaataac actttcagta ttgttctgg 660  
 tgggtttgta ccgaatgtgc tggaaggcag atccgggcac accagaaaat ggaatgac 720  
 aacctcagtc tgataaagag agcgtgaagc ttcttaccgt taagacaatt tctcatgagt 780  
 ctggtgagca ctctgcacaa ggaaaaacca agaactgaca gcttgaggaa ttctctccac 840  
 acctaggcaa taattacgct taattctcag ctctctatgca ccaagcgtgg aaaaggagaa 900  
 agtcttcgag aatcaatccc gacttccata cctgctgctg gactgtacca gacgtctgtc 960  
 ccagtaaaagt gatgtccagc tgacatgcaa taatttgatg gaatcaaaaa gaaccccg 1020  
 gctctctctgt tctctcacat ttaaaaattc cattactcca ttacaggag cgttcttagg 1080  
 aaaaggaatt ttaggaggag aatttgtag cagtgaatct gacagcccag gaggtgggct 1140  
 cgctgatagg catgactttc cttaattgtt aaagttttcc gggccaagaa tttttatcca 1200  
 tgaagacttt cctacttttc tcggtgttct tatattacct actgttagta tttattgttt 1260  
 accactatgt taatgcaggg aaaagttgca cgtgtattat taaatattag gtagaaatca 1320

taccatgcta ctttgtagat ataagtattt tattcctgct ttcgtgttac ttttaataaa 1380  
 taactactgt actcaatact ctaaaaaatac tataacatga ctgtgaaaaa aaaaaaaaaat 1440  
 aaaaaaaaaa aaaaaaaaaa 1458

<210> 144

<211> 255

<212> PRT

<213> Homo sapiens

<400> 144

Val Thr Ile Leu Phe Leu Leu Pro Ser Ile Cys Ser Ser Asn Ser Thr  
 1 5 10 15

Gly Val Leu Glu Ala Ala Asn Asn Ser Leu Val Thr Thr Thr Lys Pro  
 20 25 30

Ser Ile Thr Thr Pro Asn Thr Glu Ser Leu Gln Lys Asn Val Val Thr  
 35 40 45

Pro Thr Thr Gly Thr Thr Pro Lys Gly Thr Ile Thr Asn Glu Leu Leu  
 50 55 60

Lys Met Ser Leu Met Ser Thr Ala Thr Phe Leu Thr Ser Lys Asp Glu  
 65 70 75 80

Gly Leu Lys Ala Thr Thr Thr Asp Val Arg Lys Asn Asp Ser Ile Ile  
 85 90 95

Ser Asn Val Thr Val Thr Ser Val Thr Leu Pro Asn Ala Val Ser Thr  
 100 105 110

Leu Gln Ser Ser Lys Pro Lys Thr Glu Thr Gln Ser Ser Ile Lys Thr  
 115 120 125

Thr Glu Ile Pro Gly Ser Val Leu Gln Pro Asp Ala Ser Pro Ser Lys  
 130 135 140

Thr Gly Thr Leu Thr Ser Ile Pro Val Thr Ile Pro Glu Asn Thr Ser  
 145 150 155 160

Gln Ser Gln Val Ile Gly Thr Glu Gly Gly Lys Asn Ala Ser Thr Ser  
 165 170 175

Ala Thr Ser Arg Ser Tyr Ser Ser Ile Ile Leu Pro Val Val Ile Ala  
 180 185 190

Leu Ile Val Ile Thr Leu Ser Val Phe Val Leu Val Gly Leu Tyr Arg  
 195 200 205

Met Cys Trp Lys Ala Asp Pro Gly Thr Pro Glu Asn Gly Asn Asp Gln  
 210 215 220

Pro Gln Ser Asp Lys Glu Ser Val Lys Leu Leu Thr Val Lys Thr Ile  
 225 230 235 240

Ser His Glu Ser Gly Glu His Ser Ala Gln Gly Lys Thr Lys Asn  
 245 250 255

<210> 145

&lt;211&gt; 839

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 145

```

ggaacagcgt aagaggagag agacacattc agcagccaaa ggactcgggtg gaaagagcag 60
aacaccatag acaatatgtc gctcttggga cccaagggtgc tgctgtttct tgctgcattc 120
atcatcacct ctgactggat acccctgggg gtcaatagtc aacgaggaga cgatgtgact 180
caagcgactc cagaaacatt cacagaagat cctaactctgg tgaatgatcc cgctacagat 240
gaaacagtgt tggctgtttt ggctgatatt gcaccttcca cagatgactt ggctccctc 300
agtgaaaaaa ataccactgc agagtgtctgg gatgagaaat ttacctgcac aaggctctac 360
tctgtgcacg ggccggttaa acaatgcatt catcagttat gcttcaccag tttacgacgt 420
atgtacatcg tcaacaagga gatctgtctc cgtcttgtct gtaaggaaca cgaagctatg 480
aaagatgagc tttgccgtca gatggctggg ctgcccccta ggagactccg tcgctccaat 540
tacttccgac ttcttccctg tgaaaatgtg gatttgcaga gacccaatgg tctgtgatca 600
ttgaaaaaga ggaaagaaga aaaaatgtat gggtagagag aaggaggatc tccttcttct 660
ccaaccattg acagctaacc cttagacagt atttcttaaa ccaatccttt tgcaatgtcc 720
agcttttacc cctactctct actttttcac ccaaactgat aacattttat tcattttcta 780
gcacttaaaa tacaaagtct atattattgc aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 839

```

&lt;210&gt; 146

&lt;211&gt; 173

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 146

```

Met Ser Leu Leu Gly Pro Lys Val Leu Leu Phe Leu Ala Ala Phe Ile
 1             5             10            15

Ile Thr Ser Asp Trp Ile Pro Leu Gly Val Asn Ser Gln Arg Gly Asp
      20             25            30

Asp Val Thr Gln Ala Thr Pro Glu Thr Phe Thr Glu Asp Pro Asn Leu
      35             40            45

Val Asn Asp Pro Ala Thr Asp Glu Thr Val Leu Ala Val Leu Ala Asp
      50             55            60

Ile Ala Pro Ser Thr Asp Asp Leu Ala Ser Leu Ser Glu Lys Asn Thr
      65             70            75            80

Thr Ala Glu Cys Trp Asp Glu Lys Phe Thr Cys Thr Arg Leu Tyr Ser
      85             90            95

Val His Arg Pro Val Lys Gln Cys Ile His Gln Leu Cys Phe Thr Ser
      100            105            110

Leu Arg Arg Met Tyr Ile Val Asn Lys Glu Ile Cys Ser Arg Leu Val
      115            120            125

Cys Lys Glu His Glu Ala Met Lys Asp Glu Leu Cys Arg Gln Met Ala
      130            135            140

Gly Leu Pro Pro Arg Arg Leu Arg Arg Ser Asn Tyr Phe Arg Leu Pro
      145            150            155            160

Pro Cys Glu Asn Val Asp Leu Gln Arg Pro Asn Gly Leu
      165            170

```



&lt;210&gt; 147

&lt;211&gt; 2227

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 147

```

tgagacagaa ggttggtgtg ggaactgaa ggagtttctg tgggaggaca gggaaccctg 60
cgtttctact gtgtgattct gccaccttcc tggcccgacg ccatgggagt gacttgtgtg 120
tcccagatgc ctgtggccga gggcaagagt gttcagcaaa ccgtagagct ccttaccg 180
aaatttgaga tgcttggggc agagaagcaa ggaacatttt gtgtggactg tgagacttac 240
catacggcgc cctctacctt tggcagccaa ggtggagtat ggccctgtg tggtagctag 300
tgactgctgg agtctgctgc tgcagttcct acagagtttt ctaggcagcc acacaccagg 360
ggctcccgca gtgtttggga acagacatga tgcggtctac ggcccagcag ataccatggt 420
ccagtacatg gaactcttca acaagatccg caagcagcag caggtgccgg tggctgggat 480
tcgttagtga tgagcagctg ccagctgagc tctgtcacca ggggtactcc acaggaggag 540
caggtgctga ctttcaggtc cctggacccc agaaacccag ggtagacatg gactctactc 600
tcttctctct gttctcagct gtggcttttg ttctggggct gagtccctc cccaaccccc 660
tgactctcac acatagcccc ccacagctg ttttactccg tgccttactg gatttggcct 720
gtcctcaaga atggttaatta tgaagagtgg agaagtttgg accttgccct ctttaaggga 780
aaggtaggac acgaatgtcc ctgtctggta ctggtggggg aaatagtccg tatccccaac 840
tattaaggat ttgtcccagg ccaatgggtga acatgctgca ttttatgttt ggattgtgct 900
gtaataagag cctcttcctt cctcagagg atgtggctgg gcttctatcc tagagatgga 960
gtagaaggca caactggttt gatagttact attttctgac ctgtttgctg aagtgtattg 1020
cgaattgact ttcttagggt gttgctgag tcttttgaaa tctcttctg acatctttcc 1080
cttctgttgt gaaatatgtt aagccacagg ccaaactgct ataagatcga agtttgtttt 1140
tttctcacct aataatagag aggtaatcag atggtctagg gcagatttaa ctctacaagg 1200
ttatcaggaa ctttgtttcc ttctgtctta ttcagccatt cccagtatgt tttcttacc 1260
ataagatcaa agcagactca tcagctccat atctgcattc cactggggaa agagagagag 1320
aaaagagcag attttaaaaa aaagatatga ctgagcattg ctcagatcac tgtggccaca 1380
ctttgctgaa agagatgctg ggacatagaa cctcttatct gggcagtcac gtgctgagtt 1440
aaaactcagg gattctgtga ctacaagatg aaagacaatt tggttgctga gatgcagtta 1500
atggcttctg ttacatggag ctcaatggac gtgcccagga atgcttttgg ctgttatttt 1560
gcagttaata cctcctgtaa ctaaagcatt tgtttatgag ttgacttgag agaagggtg 1620
atctcagagc cgtcttgagc taagttggat tagtcacact aggaagttaa ttccacacct 1680
ttcgtctaag tctcagtatt gaggcctctc cagttctcat gcacctgat cttagggtta 1740
gaatacttga ccctgatacc tgcaccatgc ttcattggtc ctgagctctt tctcctgttt 1800
catttgagcc tccaaactac atatttggtc atattgcctg cctaccccat gcctgctgca 1860
gaaatattca tccagggtta ccttgatata cacagagatg gtcttggaga attgtgaatg 1920
tatgtactgt attgtcatca aggatactgt cccttatttg aaggcatcta aagagaaact 1980
gttttcagat ccaagtgtc agatctaaag cctctgcaac aagtcagggt gtggtcatgt 2040
ttcccttcta gttttggctg acaggaagct cagttcagta ccataactac agaacctgtc 2100
atctgtatgt tttgttctca ccccgttttt gttattttgt ttctggtttt ttatattgag 2160
gtatgtttta gatatagttt acaaaaataa aacgcacaga tcttaaaaaa aaaaaaaaaa 2220
aaaaaaaaa 2227

```

&lt;210&gt; 148

&lt;211&gt; 116

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 148

```

Met Met Arg Ser Thr Ala Gln Gln Ile Pro Trp Ser Ser Thr Trp Asn
  1              5              10              15

Ser Ser Thr Arg Ser Ala Ser Ser Ser Arg Cys Arg Trp Leu Gly Phe
      20              25              30

Val Ser Asp Glu Gln Leu Pro Ala Glu Leu Cys His Gln Gly Tyr Ser
      35              40              45

```

Thr Gly Gly Ala Gly Ala Asp Phe Gln Val Pro Gly Pro Gln Lys Pro  
50 55 60

Arg Val Asp Met Asp Ser Thr Leu Leu Leu Trp Phe Ser Ala Val Ala  
65 70 75 80

Phe Val Leu Gly Leu Ser Pro Leu Pro Asn Pro Leu Thr Leu Thr His  
85 90 95

Ser Pro Pro Ser Ala Val Leu Leu Arg Ala Leu Leu Asp Leu Ala Cys  
100 105 110

Pro Gln Glu Trp  
115

<210> 149

<211> 983

<212> DNA

<213> Homo sapiens

<400> 149

ggcagcaggg cccggctcct gcagacgctc ggccctccgct cattcctgac cccgcagtgg 60  
gcgcgatggc ggaggctgta ctgagggtcg cccggcgcca gctgagccag cgcggcgggt-120  
ctggagcccc catcctcctg cggcagatgt tgcagcctgt gagctgcacc ttcacgtacc 180  
tgctgggtga cagagagtcc cgggaggccg ttctgatcga cccagtccctg gaaacagcgc 240  
ctcgggatgc ccagctgac aaggagctgg ggctgctggct gctctatgct gtgaataccc 300  
actgccacgc ggaccacatt acaggctcgg ggctgctcgg ttcctcctc cctggctgcc 360  
agtctgtcat ctcccgctt agtggggccc aggtgactt acacattgag gatggagact 420  
ccatccgctt cgggcgcttc gcgttgagga ccagggccag ccttgccac accccaggct 480  
gtgtcacctt cgtcctgaat gaccacagca tggccttcac tggagatgcc ctgttgatcc 540  
gtgggtgtgg gcggacagac ttccagcaag gctgtgcca gacctgtac cactcggtcc 600  
atgaaaagat cttcacactt ccaggagact gtctgatcta cctgtctac gattaccatg 660  
ggttcacagt gtccaccgtg gaggaggaga ggactctgaa cctcggctc accctcagct 720  
gtgaggagtt tgtcaaaatc atgggcaacc tgaacttgcc taaacctcag cagatagact 780  
ttgctgttcc agccaacatg cgctgtgggg tgcagacacc cactgctga tctcacttct 840  
gtcagatgct cccatccact attaatgcac taggtgggag gagaggcg caatgacact 900  
gcacctctcc tttccaccg cattccctgg agctccctaa ataaaacttt ttttaacgtg 960  
aaaaaaaaa aaaaaaaaaa aaa 983

<210> 150

<211> 254

<212> PRT

<213> Homo sapiens

<400> 150

Met Ala Glu Ala Val Leu Arg Val Ala Arg Arg Gln Leu Ser Gln Arg  
1 5 10 15

Gly Gly Ser Gly Ala Pro Ile Leu Leu Arg Gln Met Phe Glu Pro Val  
20 25 30

Ser Cys Thr Phe Thr Tyr Leu Leu Gly Asp Arg Glu Ser Arg Glu Ala  
35 40 45

Val Leu Ile Asp Pro Val Leu Glu Thr Ala Pro Arg Asp Ala Gln Leu  
50 55 60

Ile Lys Glu Leu Gly Leu Arg Leu Leu Tyr Ala Val Asn Thr His Cys  
65 70 75 80

His Ala Asp His Ile Thr Gly Ser Gly Leu Leu Arg Ser Leu Leu Pro  
85 90 95

Gly Cys Gln Ser Val Ile Ser Arg Leu Ser Gly Ala Gln Ala Asp Leu  
100 105 110

His Ile Glu Asp Gly Asp Ser Ile Arg Phe Gly Arg Phe Ala Leu Glu  
115 120 125

Thr Arg Ala Ser Pro Gly His Thr Pro Gly Cys Val Thr Phe Val Leu  
130 135 140

Asn Asp His Ser Met Ala Phe Thr Gly Asp Ala Leu Leu Ile Arg Gly  
145 150 155 160

Cys Gly Arg Thr Asp Phe Gln Gln Gly Cys Ala Lys Thr Leu Tyr His  
165 170 175

Ser Val His Glu Lys Ile Phe Thr Leu Pro Gly Asp Cys Leu Ile Tyr  
180 185 190

Pro Ala His Asp Tyr His Gly Phe Thr Val Ser Thr Val Glu Glu Glu  
195 200 205

Arg Thr Leu Asn Pro Arg Leu Thr Leu Ser Cys Glu Glu Phe Val Lys  
210 215 220

Ile Met Gly Asn Leu Asn Leu Pro Lys Pro Gln Gln Ile Asp Phe Ala  
225 230 235 240

Val Pro Ala Asn Met Arg Cys Gly Val Gln Thr Pro Thr Ala  
245 250

&lt;210&gt; 151

&lt;211&gt; 1254

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 151

tgggctggaa cgcgccggaa tctgaggtgt gagtagagcc tgggggagag tggatccagg 60  
tgaagggggc agaggactgg gagttttcgt cctcttgaat aagaactcga caacagagtg 120  
ggaactttct gtcttgtgat ccattgacctg gtgagtcaca gctcacacca tggatttaac 180  
ctgagagctt caacttctgc ttggccctg gagttcccat gccctggtgt cttctaccag 240  
ttcttagtgt gtcgcaactgg agcacagagg acaactcgatc gtgcggcgcg cagggcgggg 300  
ggcgcgcgct gctcccccgc gggatggctg gcaactgtgct cggagtccgt gcgggcgtgt 360  
tcatcttagc cctgctctgg gtggcagtg tctgctgtg tgtgctgctg tccagagcct 420  
ccggggcggc gaggttctct gtcatttttt tattcttcgg tgcgtgcatc atcacattag 480  
ttctgttget ttcccgcga gctgggtgaat tcccagcccc agaagtggaa gtttaagattg 540  
tggatgactt ttccattggc cgctatgtcc tgcctgcttt ccttagtgcc atcttccttg 600  
gaggctcttt cttgggttta atccattatg ttctggagcc gatgtatgcc aaaccaactgc 660  
actcctactg accactcttc aggaaaacga aaacatgttc tctccttcat tgtgatgaca 720  
ttgatgagca ggaaggcact attcagagcc ttgttttgac agccctcatg ccttaaggtt 780  
agaggagtat ctgtccatca ctaagacaaa tctctggagt cctggcttcc agaaacagga 840  
ttgccaaatt gtccctgtgg ggctagattc ttaccagctt aagaaggata ttgctatctt 900  
cttagtaccg gtaccttagg atttccaaact gttttgaaag ggaaatagta acagtgatct 960  
gcttagagtg gattttcact caagtcctta gtaagtggat tggggaaaaa agcacatggg 1020  
cttctgtgtc tttttgataa tacataaaat tattcattat gaggttcgag ttgtttgcaa 1080  
aggagaggca ctcaaatttg aaaggttatt ttaatgtgat aatttggaag acttactcag 1140

argttggtca ttgaccactc tgtgcatata tttctgcaga gctctgtgaa ggcaatgagt 1200  
gtcacttccc tctgctctaa taaagcaata aataataaaa aaaaaaaaaa aaaa 1254

<210> 152  
<211> 150  
<212> PRT  
<213> Homo sapiens

<400> 152  
Met Pro Trp Cys Leu Leu Pro Val Leu Ser Val Ser His Trp Ser Thr  
1 5 10 15

Glu Asp Thr Arg Ser Cys Gly Ala Gln Gly Gly Gly Pro Pro Leu Pro  
20 25 30

Pro Arg Gly Met Ala Gly Thr Val Leu Gly Val Arg Ala Gly Val Phe  
35 40 45

Ile Leu Ala Leu Leu Trp Val Ala Val Leu Leu Leu Cys Val Leu Leu  
50 55 60

Ser Arg Ala Ser Gly Ala Ala Arg Phe Ser Val Ile Phe Leu Phe Phe  
65 70 75 80

Gly Ala Val Ile Ile Thr Leu Val Leu Leu Leu Phe Pro Arg Ala Gly  
85 90 95

Glu Phe Pro Ala Pro Glu Val Glu Val Lys Ile Val Asp Asp Phe Phe  
100 105 110

Ile Gly Arg Tyr Val Leu Leu Ala Phe Leu Ser Ala Ile Phe Leu Gly  
115 120 125

Gly Leu Phe Leu Val Leu Ile His Tyr Val Leu Glu Pro Met Tyr Ala  
130 135 140

Lys Pro Leu His Ser Tyr  
145 150

<210> 153  
<211> 1803  
<212> DNA  
<213> Homo sapiens

<400> 153  
aaaatcattc ctgatggact tcatgttgag atacatgtac aaccaggaat cagttgattg 60  
ggttggagac tacaatgaac cattgactgg ttttcatgg agaggtggat ctgaacgaga 120  
gaccacagga attcagatat ggagtgaat ctcccttacc aataaacctg atggtaaaaa 180  
ggttgacagt ttattgatgg atactcaggg aacctttgat agtcagtcaa ctttgagaga 240  
ttcagccaca gtatttggcc ttagcacaat gatcagctca atacaggtat ataacttacc 300  
ccaaaatgtc caggaggatg atcttcagca cctccagctt ttcactgagt atggcagact 360  
ggcaatggag gaaacattcc tgaagccatt tcagagtctg atattttctg ttcgagactg 420  
gagtttccca tacgaatttt catatggagc cgatgggtgt gccaaattct tggaaaaacg 480  
cctcaaggtc tcagggaacc agcatgaaga actacagaac gtcagaaaac acatccattc 540  
ctgtttcacc aacatttctt gttttctgct acctcaccct ggcttaaaag tagctaccaa 600  
tccaaacttt gatggaaaat tgaaagaaat agatgatgaa ttcacaaaaa acttgaaaat 660  
actgattcct tggctactta gcccagagag cctagatatt aaagagatca atgggaataa 720  
aatcacctgc cggggtctgg tggagtactt caaggcttat ataaagatct atcaagggtg 780  
agaattacca catcccaaat ccattgttaca ggccacagca gaagctaaca atttagcagc 840

```

cgtggcaact gccaggaca cacaacaaa aaaaatggaa gagatttgtg gtggtgacaa 900
accatttctg gcccacaaatg acttgcagac caaacacctg caacttaagg aagaatctgt 960
gaagctattc cgaggggtga agaagatggg tggggaagaa tttagccggc gttacctgca 1020
gcagttggag agtgaaatag atgaacttta catccaatat atcaagcaca atgatagcaa 1080
aaatatcttc catgcagctc gtacccagc cacactgttt gtagtcatct ttatcacata 1140
tgtgattgct ggtgtgactg gattcattgg tttggacatc atagctagcc tatgcaatat 1200
gataatggga ctgaccctta tcacctgtg cacttgggca tatatccggt actctggaga 1260
ataccgagag ctgggagctg taatagacca ggtggctgca gctctgtggg accaggggaag 1320
tacaatgag gctttgtaca agctttacag tgcagcagca acccacagac atctgtatca 1380
tcaagctttc cctacaccaa agtcggaatc tactgaacaa tcagaaaaga aaaaaatgta 1440
atgcaaatTT taagaaatac aggtgcata ccaattgtca attaaatatt cagttttatg 1500
tctccatgca aacattcaaa gtgcttccat cagaacggag taaaatacta aacacctctg 1560
aagactgcaa actggattag ttcttttact tcagtgttta ataagcagat gtatgtatgc 1620
atggttatac tattttgtta acatgtacaa ttctctgatt tttcttcaaa aatgctgtta 1680
taaagtattt gtctatttat gataacagta cacgtgttct gcttgaattt actaaattct 1740
actatgggt tataattaaa tcatgtgata ttccaaaaaa aaaaaaaaaa aaaaaaaaaa 1800
aaa

```

&lt;210&gt; 154

&lt;211&gt; 475

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 154

```

Met Asp Phe Met Leu Arg Tyr Met Tyr Asn Gln Glu Ser Val Asp Trp
  1             5             10             15

```

```

Val Gly Asp Tyr Asn Glu Pro Leu Thr Gly Phe Ser Trp Arg Gly Gly
      20             25             30

```

```

Ser Glu Arg Glu Thr Thr Gly Ile Gln Ile Trp Ser Glu Ile Phe Leu
      35             40             45

```

```

Ile Asn Lys Pro Asp Gly Lys Lys Val Ala Val Leu Leu Met Asp Thr
      50             55             60

```

```

Gln Gly Thr Phe Asp Ser Gln Ser Thr Leu Arg Asp Ser Ala Thr Val
      65             70             75             80

```

```

Phe Ala Leu Ser Thr Met Ile Ser Ser Ile Gln Val Tyr Asn Leu Ser
      85             90             95

```

```

Gln Asn Val Gln Glu Asp Asp Leu Gln His Leu Gln Leu Phe Thr Glu
      100            105            110

```

```

Tyr Gly Arg Leu Ala Met Glu Glu Thr Phe Leu Lys Pro Phe Gln Ser
      115            120            125

```

```

Leu Ile Phe Leu Val Arg Asp Trp Ser Phe Pro Tyr Glu Phe Ser Tyr
      130            135            140

```

```

Gly Ala Asp Gly Gly Ala Lys Phe Leu Glu Lys Arg Leu Lys Val Ser
      145            150            155            160

```

```

Gly Asn Gln His Glu Glu Leu Gln Asn Val Arg Lys His Ile His Ser
      165            170            175

```

```

Cys Phe Thr Asn Ile Ser Cys Phe Leu Leu Pro His Pro Gly Leu Lys
      180            185            190

```

WO 00/37630

PCT/US99/31005

Val Ala Thr Asn Pro Asn Phe Asp Gly Lys Leu Lys Glu Ile Asp Asp  
195 200 205

Glu Phe Ile Lys Asn Leu Lys Ile Leu Ile Pro Trp Leu Leu Ser Pro  
210 215 220

Glu Ser Leu Asp Ile Lys Glu Ile Asn Gly Asn Lys Ile Thr Cys Arg  
225 230 235 240

Gly Leu Val Glu Tyr Phe Lys Ala Tyr Ile Lys Ile Tyr Gln Gly Glu  
245 250 255

Glu Leu Pro His Pro Lys Ser Met Leu Gln Ala Thr Ala Glu Ala Asn  
260 265 270

Asn Leu Ala Ala Val Ala Thr Ala Lys Asp Thr Tyr Asn Lys Lys Met  
275 280 285

Glu Glu Ile Cys Gly Gly Asp Lys Pro Phe Leu Ala Pro Asn Asp Leu  
290 295 300

Gln Thr Lys His Leu Gln Leu Lys Glu Glu Ser Val Lys Leu Phe Arg  
305 310 315 320

Gly Val Lys Lys Met Gly Gly Glu Glu Phe Ser Arg Arg Tyr Leu Gln  
325 330 335

Gln Leu Glu Ser Glu Ile Asp Glu Leu Tyr Ile Gln Tyr Ile Lys His  
340 345 350

Asn Asp Ser Lys Asn Ile Phe His Ala Ala Arg Thr Pro Ala Thr Leu  
355 360 365

Phe Val Val Ile Phe Ile Thr Tyr Val Ile Ala Gly Val Thr Gly Phe  
370 375 380

Ile Gly Leu Asp Ile Ile Ala Ser Leu Cys Asn Met Ile Met Gly Leu  
385 390 395 400

Thr Leu Ile Thr Leu Cys Thr Trp Ala Tyr Ile Arg Tyr Ser Gly Glu  
405 410 415

Tyr Arg Glu Leu Gly Ala Val Ile Asp Gln Val Ala Ala Ala Leu Trp  
420 425 430

Asp Gln Gly Ser Thr Asn Glu Ala Leu Tyr Lys Leu Tyr Ser Ala Ala  
435 440 445


Ala Thr His Arg His Leu Tyr His Gln Ala Phe Pro Thr Pro Lys Ser  
450 455 460

Glu Ser Thr Glu Gln Ser Glu Lys Lys Lys Met  
465 470 475

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/31005

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC(7) : C12N 15/00; C07K 14/00 US CL : +35/69.1; 536/23.5; 530/350 According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) U.S. : +35/69.1; 536/23.5; 530/350 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) GENBANK, EST, Swissprot		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97/39123 A2 (GENETICS INSTITUTE, INC.) 23 OCTOBER 1997, page 87, SEQ ID No:23.	1-3, 5, 7
X	Database on EST, AN AA430259, HILLIER et al. 'WashU-Merck EST Project 1997,' sequence listing, 16 October 1997, see entire document.	1-3, 5, 7
X, P	WO 99/00405 A1 (GENETICS INSTITUTE, INC.) 07 January 1999, page 47, SEQ ID No:2 at position 1-179 (full length).	1-3, 5-7
X	Database on Genbank, AN AB018272, NAGASE et al., 'Prediction of the coding sequences of unidentified human genes. XI. The complete sequences of 100 new cDNA clones from brain which code for large proteins in vitro,' sequence listing, DNA Res., 1998, 5(5), pages 277-286, see entire document.	1-3, 5, 7
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document published on or after the international filing date "L" document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "Z" document member of the same patent family		
Date of the actual completion of the international search 23 MARCH 2000		Date of mailing of the international search report 17 APR 2000
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer ELIANE LAZAR-WESLEY  Telephone No. (703) 308-0196

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/31005

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
  
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
  
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-7

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.



## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/31005

### BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claims 1-7, drawn to an isolated protein of SEQ ID No:2 encoded by the polynucleotide of SEQ ID No:1, a polynucleotide sequence encoding the full length protein encoded by the cDNA insert of clone AK296\_li, and a composition.

Group II, claims 8-9, drawn to an isolated protein of SEQ ID No:22 encoded by the polynucleotide of SEQ ID No:21, and a polynucleotide sequence encoding the full length protein encoded by the cDNA insert of clone AS34\_li.

The inventions listed as Groups I-II do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: the special technical feature of Group I is the polypeptide of SEQ ID No:2 and the polynucleotide of SEQ ID No:1. The polypeptide of SEQ ID No:22 and polynucleotide of SEQ ID No:21 of Group II do not share the special technical feature of Group I, as they have different structures and functions.